

09/76231

Attorney Docket No.: 46.US2.PC

Page 1

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 USC 371

International Application No.:

PCT/IB99/01444

International Filing Date:

June 8, 1999

Priority Date Claimed:

August 7, 1998

Title of Invention:

NUCLEIC ACIDS ENCODING HUMAN TBC-1 PROTEIN AND

POLYMORPHIC MARKERS THEREOF.

Applicant(s) for DO/EO/US:

Blumenfeld, M., et al

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

- 1. (X) This is a **FIRST** submission of items concerning a filing under 35 USC 371.
- This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 USC 371.
- 3. (X) This express request to begin national examination procedures (35 USC 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 USC 371(b) and PCT Articles 22 and 39(1).
- 4. (X) A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
- 5. (X) A copy of the International Application as filed (35 USC 371(c)(2))
 - a) () is transmitted herewith (required only if not transmitted by the International Bureau).
 - b) (X) has been transmitted by the International Bureau.
 - c) is not required, as the application was filed in the United States Receiving Office (RO/US).
- 6. () A translation of the International Application into English (35 USC 371(c)(2)).
- 7. (X) Amendments to the claims of the International Application under PCT Article 19 (35 USC 371(c)(3))
 - a) () are transmitted herewith (required only if not transmitted by the International Bureau).
 - b) (X) have been transmitted by the International Bureau.
 - c) () have not been made; however, the time limit for making such amendments has NOT expired.
 - d) () have not been made and will not be made.
- 8. () A translation of the amendments to the claims under PCT Article 19 (35 USC 371(c)(3)).
- 9. () An oath or declaration of the inventor(s) (35 USC 371(c)(4)).
- 10. (X) A copy of the International Preliminary Examination Report with any annexes thereto, such as any amendments made under PCT Article 34.
- 11. () A translation of the annexes, such as any amendments made under PCT Article 34, to the International Preliminary Examination Report under PCT Article 36 (35 USC 371(c)(5)).

Items 12. to 23. below concern other document(s) or information included:

09/762311 JC02 Rec'd PCT/PTO 02 FEB 20\

Attorney Docket No.: 46.US2.PCT

February 1, 200	Page 2
12. ()	An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
13. ()	An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
14. (X)	A FIRST preliminary amendment. A SECOND or SUBSEQUENT preliminary amendment.
15. ()	A substitute specification.
16. ()	A power of attorney and/or address letter.
17. (X)	International Application as published.
18. ()	Small Entity Statement.
19. (X)	Sequence Submission Statement in 1 page.
20. (X)	Paper Copy of the Sequence Listing in 52 pages.
21. (X)	Sequence Listing in Computer Readable Format.
22. (X)	A return prepaid postcard.
23. (X)	The following fees are submitted:

				FEES
	BASIC FEE		7	\$860.00
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	22 - 20 =	2 ×	\$18	\$ 36.00
Independent Claims	6 - 3 =	3 ×	\$80	\$240.00
Multiple dependent claims(s) (if applicable) \$270				
	TOTAL OF AB	OVE CALCULATION	NS \$1,136.00	
Reduction by 1/2 for filing by statement must also be filed.	small entity (if applicable (NOTE 37 CFR 1.9, 1.27,	e). Verified Small Entity 1.28)	y \$ 0	
	TOTAL NATIO	NAL FEE		\$1,136.00
TOTAL FEES ENCLOSED				\$1,136.00
		Please credit depos Amount to be char		\$0 \$0

a. (X) A check in the amount of \$1,136.00 to cover the above fees is enclosed.



09/762311

JC02 Rec'd PCT/PTO 0 2 FEB 2001

Jun 1 Feb 0/

Attorney Docket No.: 46.US2.PCT

February 1, 2001

Page 3

- b. () Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40 per property.
- c. (X) The Commissioner is hereby authorized to charge only those additional fees which may be required, now or in the future, to avoid abandonment of the application, or credit any overpayment to Deposit Account No. 501181. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

GENSET CORPORATION 875Prospect Street, Suite 206 La Jolla CA 92037 Signature

John Lucas, Ph.D., J.D. Registration No. 43,373

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant

Blumenfeld, M., et al

App. No.

To Be Assigned

Filed

Herewith

For

Hall find

U

NUCLEIC ACIDS ENCODING HUMAN TBC-1 PROTEIN

AND POLYMORPHIC MARKERS THEREOF

Examiner

Unknown

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents

Washington, D.C. 20231

Dear Sir:

Please amend the application as follows.

IN THE SPECIFICAITON:

After the Title, but before "FIELD OF THE INVENTION", please add:

--This application claims benefits under 35 U.S.C. § 119(e) to U.S. Provisional Applications Serial No. 60/095,653 and under 35 U.S.C. § 371 to International Application No. PCT/IB99/01444, both of which are hereby incorporated by reference herein in their entireties.--

IN THE CLAIMS

Please cancel claims 2 to 4, 9, 10, 14, 15, 17 to 21, 28 to 32, 34, and 36 to 38 without prejudice.

Please amend the following claims:

- --1. (amended) A composition comprising: [A] an isolated, purified, or recombinant polynucleotide comprising a nucleotide sequence selected from the group consisting of the nucleotide sequences [a contiguous span of at least 12 nucleotides] of SEQ ID No 1, 2, 3 and 4, or the complements thereof.
- 5. (amended) A composition comprising: [A] an isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 12 [8 to 50] nucleotides of any one of SEQ ID Nos 1 and 2 or the complement thereof, wherein said span includes a TBC-1-related biallelic marker in said sequence.
- 6. (amended) The [A] polynucleotide according to claim 5, wherein said TBC-1-related biallelic marker is selected from the group consisting of [A1 to A19] the biallelic markers in positions 9494 of SEQ ID No 1, and 1443, 5247, 6223, 14723, 19186, 18997, 19891, 29617, 42519, 69324, 69181, 69146, 76458, 78595, 82159, 84522, 84810, and 89967 of SEQ ID No 2.
- 7. (amended) The [A] polynucleotide according to [any one of claims 5 or 6] claim 5, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide.
- 8. (amended) The [A] polynucleotide according to claim 7, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said polynucleotide.
- 11. (amended) The [A] polynucleotide according to [any one of claims 5 or 6] claim 5, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide.
- 12. (amended) A composition comprising: [A] an isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides of anyone of SEQ ID Nos 1 and 2 or the complement thereof, wherein the 3' end of said contiguous span is

located at the 3' end of said polynucleotide, and wherein the 3' end of said polynucleotide is located within 20 nucleotides upstream of a *TBC-1*-related biallelic marker in said sequence.

- 13. (amended) The [A] polynucleotide according to claim 12, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *TBC-1*-related biallelic marker in said sequence.
- 16. (amended) A composition comprising: [A] an isolated, purified, or recombinant polynucleotide which encodes a polypeptide comprising a contiguous span of at least 6 amino acids of SEQ ID No 5.
- 22. (amended) <u>The [A]</u> polynucleotide according to [any one of claims 1 to 21] <u>claim 5</u> attached to a solid support.
- 25. (amended) <u>The [A]</u> polynucleotide according to [any one of claims 1 to 21] <u>claim 5</u> further comprising a label.
- 26. (amended) A <u>composition comprising</u>: a recombinant vector comprising a polynucleotide according to [any one of claims 1 to 4 and 16] <u>claim 5</u>.
- 27. (amended) A <u>composition comprising: a host cell comprising a recombinant vector according to claim 26.</u>
- 33. (amended) A method according to claim [29] <u>44</u>, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said determining step.
- 35. (amended) A method according to claim[29] <u>43</u>, wherein said determining is performed by a hybridization assay, <u>sequencing assay</u>, <u>microsequencing assay or enzyme-based</u> mismatch detection assay.

39. (amended) A method according to [any one of claims 29 to 38] claim 44, wherein said TBC-1-related biallelic marker is selected from the group consisting of [A1 to A9 and the complements thereof] the biallelic markers in positions 9494 of the SEQ ID No 1, and 1443,

5247, 6223, 14723, 19186, 18997, 19891, 29617, 42519, 69324, 69181, 69146, 76458, 78595, 82159, 84522, 84810, and 89967 of the SEQ ID No 2.

Please add the following claims:

- 40. (new) A composition comprising: an isolated, purified, or recombinant polypeptide comprising a continuous span of at least 8 amino acids of SEQ ID No 5.
- 41. (new) A composition comprising: an isolated or purified antibody composition capable of selectively binding to an epitope-containing fragment of a polypeptide according to claim 35.
- 42. (new) A method of making a purified or isolated TBC-1 polypeptide encoded by a polynucleotide of claim 1; wherein said method comprises the steps of:
 - (i) obtaining a cell capable of expressing said polypeptide;
 - (ii) growing said cell under conditions suitable to produce said polypeptide; and
 - (iii) isolating said polypeptide.
- 43. (new) A method of genotyping comprising the steps of:
 - (a) obtaining a nucleic acid sample from an individual; and
- (b) determining the identity of a polymorphic base at a TBC-1-related biallelic marker or the complement thereof in said nucleic acid sample, wherein the identity of the polymorphic base determines the genotype of the individual at said TBC-1-related biallelic marker and, wherein said TBC-1-related biallelic marker is positioned in SEQ ID NO: 1 or SEQ ID NO:2.--

<u>REMARKS</u>

Amendments to the claims:

Claims 2 to 4, 9, 10, 14, 15, 17 to 21, 28 to 32, 34, and 36 to 38 have been canceled without prejudice. New Claims 40 to 43 have been added. Support for the new Claims is found throughout the specification

Accordingly, claims 1, 5 to 8, 11 to 13, 16, 22 to 27, 33, 35, and 39 to 43 are pending in this application. Please charge any additional fees, including any fees for additional extension of time, or credit overpayment to Deposit Account No. 501181.

Dated: 1 Feb 0

Respectfully submitted,

By:

John Lucas
Registration No. 43,373

Genset Corporation

875 Prospect Street, Suite 206

La Jolla, CA 92037

PCT/IB99/01

Nucleic acids encoding human TBC-1 protein and polymorphic markers

thereof.

FIELD OF THE INVENTION

The invention concerns genomic and cDNA sequences of the human *TBC-1* gene. The invention also concerns polypeptides encoded by the *TBC-1* gene. The invention also deals with antibodies directed specifically against such polypeptides that are useful as diagnostic reagents. The invention further encompasses biallelic markers of the *TBC-1* gene useful in genetic analysis.

BACKGROUND OF THE INVENTION

The incidence of prostate cancer has dramatically increased over the last decades. It averages 30-50/100,000 males in Western European countries as well as within the US White male population. In these countries, it has recently become the most commonly diagnosed malignancy, being one of every four cancers diagnosed in American males. Prostate cancer's incidence is very much population specific, since it varies from 2/100,000 in China, to over 80/100,000 among African-American males.

In France, the incidence of prostate cancer is 35/100,000 males and it is increasing by 10/100,000 per decade. Mortality due to prostate cancer is also growing accordingly. It is the second cause of cancer death among French males, and the first one among French males aged over 70. This makes prostate cancer a serious burden in terms of public health.

Prostate cancer is a latent disease. Many men carry prostate cancer cells without overt signs of disease. Autopsies of individuals dying of other causes show prostate cancer cells in 30 % of men at age 50 and in 60 % of men at age 80. Furthermore, prostate cancer can take up to 10 years to kill a patient after the initial diagnosis.

The progression of the disease usually goes from a well-defined mass within the prostate to a breakdown and invasion of the lateral margins of the prostate, followed by metastasis to regional lymph nodes, and metastasis to the bone marrow. Cancer metastasis to bone is common and often associated with uncontrollable pain.

Unfortunately, in 80 % of cases, diagnosis of prostate cancer is established when the disease has already metastasized to the bones. Of special interest is the observation that prostate cancers frequently grow more rapidly in sites of metastasis than within the prostate itself.

Early-stage diagnosis of prostate cancer mainly relies today on Prostate Specific Antigen (PSA) dosage, and allows the detection of prostate cancer seven years before clinical symptoms become apparent. The effectiveness of PSA dosage diagnosis is however limited, due to its inability to discriminate between malignant and non-malignant affections of the organ and because not all prostate cancers give rise to an elevated serum PSA concentration. Furthermore, PSA dosage and

5

WO 00/08209 PCT/IB99/01444

other currently available approaches such as physical examination, tissue biopsy and bone scans are of limited value in predicting disease progression.

Therefore, there is a strong need for a reliable diagnostic procedure which would enable a more systematic early-stage prostate cancer prognosis.

Although an early-stage prostate cancer prognosis is important, the possibility of measuring the period of time during which treatment can be deferred is also interesting as currently available medicaments are expensive and generate important adverse effects. However, the aggressiveness of prostate tumors varies widely. Some tumors are relatively aggressive, doubling every six months whereas others are slow-growing, doubling once every five years. In fact, the majority of prostate 10 cancers grows relatively slowly and never becomes clinically manifest. Very often, affected patients are among the elderly and die from another disease before prostate cancer actually develops. Thus, a significant question in treating prostate carcinoma is how to discriminate between tumors that will progress and those that will not progress during the expected lifetime of the patient.

Hence, there is also a strong need for detection means which may be used to evaluate the aggressiveness or the development potential of prostate cancer tumors once diagnosed.

Furthermore, at the present time, there is no means to predict prostate cancer susceptibility. It would also be very beneficial to detect individual susceptibility to prostate cancer. This could allow preventive treatment and a careful follow up of the development of the tumor.

A further consequence of the slow growth rate of prostate cancer is that few cancer cells are actively dividing at any one time, rendering prostate cancer generally resistant to radiation and chemotherapy. Surgery is the mainstay of treatment but it is largely ineffective and removes the ejaculatory ducts, resulting in impotence. Oral oestrogens and luteinizing releasing hormone analogs are also used for treatment of prostate cancer. These hormonal treatments provide marked improvement for many patients, but they only provide temporary relief. Indeed, most of these 25 cancers soon relapse with the development of hormone-resistant tumor cells and the oestrogen treatment can lead to serious cardiovascular complications. Consequently, there is a strong need for preventive and curative treatment of prostate cancer.

Efficacy/tolerance prognosis could be precious in prostate cancer therapy. Indeed, hormonal therapy, the main treatment currently available, presents important side effects. The use of 30 chemotherapy is limited because of the small number of patients with chemosensitive tumors. Furthermore the age profile of the prostate cancer patient and intolerance to chemotherapy make the systematic use of this treatment very difficult.

Therefore, a valuable assessment of the eventual efficacy of a medicament to be administered to a prostate cancer patent as well as the patent's eventual tolerance to it may permit to 35 enhance the benefit/risk ratio of prostate cancer treatment.

It is known today that there is a familial risk of prostate cancer. Clinical studies in the 1950s had already demonstrated a familial aggregation in prostate cancer. Control-case clinical studies

5

WO 00/08209 PCT/IB99/01444 3

have been conducted more recently to attempt to evaluate the incidence of the genetic risk factors in the disease. Thus Steinberg et al., 1990, and McWhorter et al., 1992 confirm that the risk of prostate cancer is increased in subjects having one or more relatives already affected by the disease and when forms of early diagnosis in the relatives exist.

It is now well established that cancer is a disease caused by the deregulation of the expression of certain genes. In fact, the development of a tumor necessitates an important succession of steps. Each of these steps comprises the deregulation of an important gene intervening in the normal metabolism of the cell and the emergence of an abnormal cellular sub-clone which overwhelms the other cell types because of a proliferative advantage. The genetic origin of this 10 concept has found confirmation in the isolation and the characterization of genes which could be responsible. These genes, commonly called "cancer genes", have an important role in the normal metabolism of the cell and are capable of intervening in carcinogenesis following a change.

Recent studies have identified three groups of genes which are frequently mutated in cancer. The first group of genes, called oncogenes, are genes whose products activate cell proliferation. The normal non-mutant versions are called protooncogenes. The mutated forms are excessively or inappropriately active in promoting cell proliferation, and act in the cell in a dominant way in that a single mutant allele is enough to affect the cell phenotype. Activated oncogenes are rarely transmitted as germline mutations since they may probably be lethal when expressed in all the cells. Therefore oncogenes can only be investigated in tumor tissues.

The second group of genes which are frequently mutated in cancer, called tumor suppressor genes, are genes whose products inhibit cell growth. Mutant versions in cancer cells have lost their normal function, and act in the cell in a recessive way in that both copies of the gene must be inactivated in order to change the cell phenotype. Most importantly, the tumor phenotype can be rescued by the wild type allele, as shown by cell fusion experiments first described by Harris and 25 colleagues (1969). Germline mutations of tumor suppressor genes may be transmitted and thus studied in both constitutional and tumor DNA from familial or sporadic cases. The current family of tumor suppressors includes DNA-binding transcription factors (i.e., p53, WT1), transcription regulators (i.e., RB, APC, probably BRCA1), protein kinase inhibitors (i.e., p16), among others (for review, see Haber D & Harlow E, 1997).

The third group of genes which are frequently mutated in cancer, called mutator genes, are responsible for maintaining genome integrity and/or low mutation rates. Loss of function of both alleles increases cell mutation rates, and as a consequence, proto-oncogenes and tumor suppressor genes may be mutated. Mutator genes can also be classified as tumor suppressor genes, except for the fact that tumorigenesis caused by this class of genes cannot be suppressed simply by restoration 35 of a wild-type allele, as described above. Genes whose inactivation may lead to a mutator phenotype include mismatch repair genes (i.e., MLH1, MSH2), DNA helicases (i.e., BLM, WRN)

25

WO 00/08209 PCT/IB99/01444

or other genes involved in DNA repair and genomic stability (i.e., p53, possibly BRCA1 and BRCA2) (For review see Haber D & Harlow E, 1997; Fishel R & Wilson T. 1997; Ellis NA,1997).

There is growing evidence that a critical event in the progression of a tumor cell from a non-metastatic to metastatic phenotype is the loss of function of metastasis-suppressor genes. These genes specifically suppress the ability of a cell to metastasize. Work from several groups has demonstrated that human chromosomes 8, 10, 11 and 17 encode prostate cancer metastasis suppressor activities. However, other human chromosomes such as chromosomes 1, 7, 13, 16, and 18 may also be associated to prostate cancer.

It thus remains to localize and to identify the genes specifically involved in the development and the progression of prostate cancers starting from the genetic analysis of the hereditary and the non-hereditary forms and to define their clinical implications in terms of prognosis and therapeutic innovations.

SUMMARY OF THE INVENTION

The present invention pertains to nucleic acid molecules comprising the genomic sequence of a novel human gene which encodes a TBC-1 protein. The *TBC-1* genomic sequences comprise regulatory sequence located upstream (5'-end) and downstream (3'-end) of the transcribed portion of said gene, these regulatory sequences being also part of the invention. The human *TBC-1* genomic sequence is included in a previously unknown candidate region of prostate cancer located on chromosome 4.

The invention also deals with the two complete cDNA sequences encoding the TBC-1 protein, as well as with the corresponding translation product.

Oligonucleotide probes or primers hybridizing specifically with a *TBC-1* genomic or cDNA sequence are also part of the present invention, as well as DNA amplification and detection methods using said primers and probes.

A further object of the invention consists of recombinant vectors comprising any of the nucleic acid sequences described above, and in particular of recombinant vectors comprising a *TBC-1* regulatory sequence or a sequence encoding a TBC-1 protein, as well as of cell hosts and transgenic non human animals comprising said nucleic acid sequences or recombinant vectors.

The invention also concerns a TBC-1-related biallelic marker and the use thereof.

Finally, the invention is directed to methods for the screening of substances or molecules that inhibit the expression of *TBC-1*, as well as with methods for the screening of substances or molecules that interact with a TBC-1 polypeptide.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: An amino acid alignment of a portion of the amino acid sequence of the TBC-1 protein of SEQ ID No 5 with other proteins sharing amino acid homology with TBC-1. The amino acid numbering refers to the murine TBC-1.

5 Brief Description of the sequences provided in the Sequence Listing

SEQ ID No 1 contains a first part of the *TBC-1* genomic sequence comprising the 5' regulatory sequence and the exons 1, 1bis, and 2.

SEQ ID No 2 contains a second part of the *TBC-1* genomic sequence comprising the 12 last exons of the *TBC-1* gene and the 3'regulatory sequence.

SEO ID No 3 contains a first cDNA sequence of the TBC-1 gene.

SEQ ID No 4 contains a second cDNA sequence of the TBC-1 gene.

SEQ ID No 5 contains the amino acid sequence encoded by the cDNAs of SEQ ID Nos 3 and 4.

SEQ ID No 6 contains a primer containing the additional PU 5' sequence described further in Example 3.

SEQ ID No 7 contains a primer containing the additional RP 5' sequence described further in Example 3.

In accordance with the regulations relating to Sequence Listings, the following codes have been used in the Sequence Listing to indicate the locations of biallelic markers within the sequences and to identify each of the alleles present at the polymorphic base. The code "r" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is an adenine. The code "y" in the sequences, indicates that one allele of the polymorphic base is a thymine, while the other allele is a cytosine. The code "m" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an cytosine. The code "k" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a thymine. The code "s" in the sequences indicates that one allele of the polymorphic base is a guanine, while the other allele is a cytosine. The code "w" in the sequences indicates that one allele of the polymorphic base is an adenine, while the other allele is an thymine. The nucleotide code of the original allele for each biallelic marker is the following:

30	Biallelic marker	Original allele	
	99-430-352	G	
	99-20508-456	C	
	99-20469-213	С	
	5-254-227	Α	
35	5-257-353	С	
	99-20511-32	T	

12507
Ę
4
1
77
ļ.,,
ļ.,,
315
F
Ļ

	WO 00/08209		PCT/IB99/01444
	•	. 6	
	99-20511-221	Α	
	99-20504-90	G	
	99-20493-238	Α	
	99-20499-221	G	
5	99-20499-364	Α	
	99-20499-399	Α	
	5-249-304	G	
	99-20485-269	Α	·
	99-20481-131	G	•
10	99-20481-419	T	
	99-20480-233	Α	

DETAILED DESCRIPTION OF THE INVENTION

The present invention concerns polynucleotides and polypeptides related to the human *TBC-1* gene (also termed "*TBC-1* gene" throughout the present specification), which is potentially involved in the regulation of the differentiation of various cell types in mammals. A deregulation or an alteration of *TBC-1* expression, or alternatively an alteration in the amino acid sequence of the TBC-1 protein may be involved in the generation of a pathological state related to cell differentiation in a patient, more particularly to abnormal cell proliferation leading to cancer states, such as prostate cancer.

20 Definitions

Before describing the invention in greater detail, the following definitions are set forth to illustrate and define the meaning and scope of the terms used to describe the invention herein.

The term "<u>TBC-1 gene</u>", when used herein, encompasses mRNA and cDNA sequences encoding the TBC-1 protein. In the case of a genomic sequence, the TBC-1 gene also includes native regulatory regions which control the expression of the coding sequence of the TBC-1 gene.

The term "<u>functionally active fragment</u>" of the TBC-1 protein is intended to designate a polypeptide carrying at least one of the structural features of the TBC-1 protein involved in at least one of the biological functions and/or activity of the TBC-1 protein.

A "heterologous" or "exogenous" polynucleotide designates a purified or isolated nucleic acid that has been placed, by genetic engineering techniques, in the environment of unrelated nucleotide sequences, such as the final polynucleotide construct does not occur naturally. An illustrative, but not limitative, embodiment of such a polynucleotide construct may be represented by a polynucleotide comprising (1) a regulatory polynucleotide derived from the TBC-1 gene sequence and (2) a polynucleotide encoding a cytokine, for example GM-CSF. The polypeptide

PCT/IB99/01444

encoded by the heterologous polynucleotide will be termed an heterologous polypeptide for the purpose of the present invention.

By a "biologically active fragment or variant" of a regulatory polynucleotide according to the present invention is intended a polynucleotide comprising or alternatively consisting in a fragment of said polynucleotide which is functional as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host.

For the purpose of the invention, a nucleic acid or polynucleotide is "<u>functional</u>" as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said regulatory polynucleotide contains nucleotide sequences which contain transcriptional and translational regulatory information, and such sequences are "operatively linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide. An operable linkage is a linkage in which the regulatory nucleic acid and the DNA sequence sought to be expressed are linked in such a way as to permit gene expression.

A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell required to initiate the specific transcription of a gene.

A sequence which is "operably linked" to a regulatory sequence such as a promoter means that said regulatory element is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the nucleic acid of interest.

As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in a functional relationship. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence. More precisely, two DNA molecules (such as a polynucleotide containing a promoter region and a polynucleotide encoding a desired polypeptide or polynucleotide) are said to be "operably linked" if the nature of the linkage between the two polynucleotides does not (1) result in the introduction of a frame-shift mutation or (2) interfere with the ability of the polynucleotide containing the promoter to direct the transcription of the coding polynucleotide. The promoter polynucleotide would be operably linked to a polynucleotide encoding a desired polypeptide or a desired polynucleotide if the promoter is capable of effecting transcription of the polynucleotide of interest.

The term "primer" denotes a specific oligonucleotide sequence which is complementary to a target nucleotide sequence and used to hybridize to the target nucleotide sequence. A primer serves as an initiation point for nucleotide polymerization catalyzed by either DNA polymerase, RNA polymerase or reverse transcriptase.

The term "probe" denotes a defined nucleic acid segment (or nucleotide analog segment, e.g., polynucleotide as defined hereinbelow) which can be used to identify a specific polynucleotide sequence present in samples, said nucleic acid segment comprising a nucleotide sequence complementary of the specific polynucleotide sequence to be identified.

WO 00/08209 PCT/IB99/01444

The terms "sample" or "material sample" are used herein to designate a solid or a liquid material suspected to contain a polynucleotide or a polypeptide of the invention. A solid material may be, for example, a tissue slice or biopsy within which is searched the presence of a polynucleotide encoding a TBC-1 protein, either a DNA or RNA molecule or within which is searched the presence of a native or a mutated TBC-1 protein, or alternatively the presence of a desired protein of interest the expression of which has been placed under the control of a TBC-1 regulatory polynucleotide. A liquid material may be, for example, any body fluid like serum, urine etc., or a liquid solution resulting from the extraction of nucleic acid or protein material of interest from a cell suspension or from cells in a tissue slice or biopsy. The term "biological sample" is also used and is more precisely defined within the Section dealing with DNA extraction.

As used herein, the term "<u>purified</u>" does not require absolute purity; rather, it is intended as a relative definition. Purification if starting material or natural material to at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated. As an example, purification from 0.1% concentration to 10% concentration is two orders of magnitude.

The term "isolated" requires that the material be removed from its original environment (e.g. the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or DNA or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotide could be part of a vector and/or such polynucleotide or polypeptide could be part of a composition and still be isolated in that the vector or composition is not part of its natural environment.

The term "polypeptide" refers to a polymer of amino acids without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term also does not specify or exclude post-expression modifications of polypeptides, for example, polypeptides which include the covalent attachment of glycosyl groups, acetyl groups, phosphate groups, lipid groups and the like are expressly encompassed by the term polypeptide. Also included within the definition are polypeptides which contain one or more analogs of an amino acid (including, for example, non-naturally occurring amino acids, amino acids which only occur naturally in an unrelated biological system, modified amino acids from mammalian systems etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

The term "<u>recombinant polypeptide</u>" is used herein to refer to polypeptides that have been artificially designed and which comprise at least two polypeptide sequences that are not found as contiguous polypeptide sequences in their initial natural environment, or to refer to polypeptides which have been expressed from a recombinant polynucleotide.

رقسة,

10

WO 00/08209 PCT/IB99/01444

The term "purified" is used herein to describe a polypeptide of the invention which has been separated from other compounds including, but not limited to nucleic acids, lipids, carbohydrates and other proteins. A polypeptide is substantially pure when at least about 50%, preferably 60 to 75% of a sample exhibits a single polypeptide sequence. A substantially pure polypeptide typically comprises about 50%, preferably 60 to 90% weight/weight of a protein sample, more usually about 95%, and preferably is over about 99% pure. Polypeptide purity or homogeneity is indicated by a number of means well known in the art, such as polyacrylamide gel electrophoresis of a sample, followed by visualizing a single polypeptide band upon staining the gel. For certain purposes higher resolution can be provided by using HPLC or other means well known in the art.

As used herein, the term "non-human animal" refers to any non-human vertebrate, birds and more usually mammals, preferably primates, farm animals such as swine, goats, sheep, donkeys, and horses, rabbits or rodents, more preferably rats or mice. As used herein, the term "animal" is used to refer to any vertebrate, preferable a mammal. Both the terms "animal" and "mammal" expressly embrace human subjects unless preceded with the term "non-human".

As used herein, the term "antibody" refers to a polypeptide or group of polypeptides which are comprised of at least one binding domain, where an antibody binding domain is formed from the folding of variable domains of an antibody molecule to form three-dimensional binding spaces with an internal surface shape and charge distribution complementary to the features of an antigenic determinant of an antigen, which allows an immunological reaction with the antigen. Antibodies include recombinant proteins comprising the binding domains, as wells as fragments, including Fab, Fab', F(ab)₂, and F(ab')₂ fragments.

As used herein, an "antigenic determinant" is the portion of an antigen molecule, in this case a TBC-1 polypeptide, that determines the specificity of the antigen-antibody reaction. An "epitope" refers to an antigenic determinant of a polypeptide. An epitope can comprise as few as 3 amino acids in a spatial conformation which is unique to the epitope. Generally an epitope consists of at least 6 such amino acids, and more usually at least 8-10 such amino acids. Methods for determining the amino acids which make up an epitope include x-ray crystallography, 2-dimensional nuclear magnetic resonance, and epitope mapping e.g. the Pepscan method described by Geysen et al. 1984; PCT Publication No. WO 84/03564; and PCT Publication No. WO 84/03506.

Throughout the present specification, the expression "nucleotide sequence" may be employed to designate indifferently a polynucleotide or an oligonucleotide or a nucleic acid. More precisely, the expression "nucleotide sequence" encompasses the nucleic material itself and is thus not restricted to the sequence information (i.e. the succession of letters chosen among the four base letters) that biochemically characterizes a specific DNA or RNA molecule.

As used interchangeably herein, the term "oligonucleotides", and "polynucleotides" include RNA, DNA, or RNA/DNA hybrid sequences of more than one nucleotide in either single chain or

WO 00/08209 PCT/IB99/01444

duplex form. The term "nucleotide" as used herein as an adjective to describe molecules comprising RNA, DNA, or RNA/DNA hybrid sequences of any length in single-stranded or duplex form. The term "nucleotide" is also used herein as a noun to refer to individual nucleotides or varieties of nucleotides, meaning a molecule, or individual unit in a larger nucleic acid molecule, comprising a purine or pyrimidine, a ribose or deoxyribose sugar moiety, and a phosphate group, or phosphodiester linkage in the case of nucleotides within an oligonucleotide or polynucleotide. Although the term "nucleotide" is also used herein to encompass "modified nucleotides" which comprise at least one modification (a) an alternative linking group, (b) an analogous form of purine, (c) an analogous form of pyrimidine, or (d) an analogous sugar, for examples of analogous linking groups, purine, pyrimidines, and sugars see for example PCT publication No WO 95/04064. However, the polynucleotides of the invention are preferably comprised of greater than 50% conventional deoxyribose nucleotides, and most preferably greater than 90% conventional deoxyribose nucleotides. The polynucleotide sequences of the invention may be prepared by any known method, including synthetic, recombinant, *ex vivo* generation, or a combination thereof, as well as utilizing any purification methods known in the art.

The term "heterozygosity rate" is used herein to refer to the incidence of individuals in a population which are heterozygous at a particular allele. In a biallelic system, the heterozygosity rate is on average equal to $2P_a(1-P_a)$, where P_a is the frequency of the least common allele. In order to be useful in genetic studies, a genetic marker should have an adequate level of heterozygosity to allow a reasonable probability that a randomly selected person will be heterozygous.

The term "genotype" as used herein refers the identity of the alleles present in an individual or a sample. In the context of the present invention a genotype preferably refers to the description of the biallelic marker alleles present in an individual or a sample. The term "genotyping" a sample or an individual for a biallelic marker consists of determining the specific allele or the specific nucleotide carried by an individual at a biallelic marker.

The term "polymorphism" as used herein refers to the occurrence of two or more alternative genomic sequences or alleles between or among different genomes or individuals. "Polymorphic" refers to the condition in which two or more variants of a specific genomic sequence can be found in a population. A "polymorphic site" is the locus at which the variation occurs. A single nucleotide polymorphism is a single base pair change. Typically a single nucleotide polymorphism is the replacement of one nucleotide by another nucleotide at the polymorphic site. Deletion of a single nucleotide or insertion of a single nucleotide, also give rise to single nucleotide polymorphisms. In the context of the present invention "single nucleotide polymorphism" preferably refers to a single nucleotide substitution. However, the polymorphism can also involve an insertion or a deletion of at least one nucleotide, preferably between 1 and 5 nucleotides. Typically, between different genomes or between different individuals, the polymorphic site may be occupied by two different nucleotides.

In the first from the first first first for

10

The term "biallelic polymorphism" and "biallelic marker" are used interchangeably herein to refer to a single nucleotide polymorphism having two alleles at a fairly high frequency in the population. A "biallelic marker allele" refers to the nucleotide variants present at a biallelic marker site. Typically, the frequency of the less common allele of the biallelic markers of the present 5 invention has been validated to be greater than 1%, preferably the frequency is greater than 10%, more preferably the frequency is at least 20% (i.e. heterozygosity rate of at least 0.32), even more preferably the frequency is at least 30% (i.e. heterozygosity rate of at least 0.42). A biallelic marker wherein the frequency of the less common allele is 30% or more is termed a "high quality biallelic marker".

The location of nucleotides in a polynucleotide with respect to the center of the polynucleotide are described herein in the following manner. When a polynucleotide has an odd number of nucleotides, the nucleotide at an equal distance from the 3' and 5' ends of the polynucleotide is considered to be "at the center" of the polynucleotide, and any nucleotide immediately adjacent to the nucleotide at the center, or the nucleotide at the center itself is considered to be "within I nucleotide of the center." With an odd number of nucleotides in a polynucleotide any of the five nucleotides positions in the middle of the polynucleotide would be considered to be within 2 nucleotides of the center, and so on. When a polynucleotide has an even number of nucleotides, there would be a bond and not a nucleotide at the center of the polynucleotide. Thus, either of the two central nucleotides would be considered to be "within 1 nucleotide of the center" and any of the four nucleotides in the middle of the polynucleotide would be considered to be "within 2 nucleotides of the center", and so on. For polymorphisms which involve the substitution, insertion or deletion of 1 or more nucleotides, the polymorphism, allele or biallelic marker is "at the center" of a polynucleotide if the difference between the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 3' end of the 25 polynucleotide, and the distance from the substituted, inserted, or deleted polynucleotides of the polymorphism and the 5' end of the polynucleotide is zero or one nucleotide. If this difference is 0 to 3, then the polymorphism is considered to be "within 1 nucleotide of the center." If the difference is 0 to 5, the polymorphism is considered to be "within 2 nucleotides of the center." If the difference is 0 to 7, the polymorphism is considered to be "within 3 nucleotides of the center," and 30 so on.

As used herein the terminology "defining a biallelic marker" means that a sequence includes a polymorphic base from a biallelic marker. The sequences defining a biallelic marker may be of any length consistent with their intended use, provided that they contain a polymorphic base from a biallelic marker. The sequence has between 1 and 500 nucleotides in length, preferably 35 between 5, 10, 15, 20, 25, or 40 and 200 nucleotides and more preferably between 30 and 50 nucleotides in length. Each biallelic marker therefore corresponds to two forms of a polynucleotide sequence included in a gene, which, when compared with one another, present a nucleotide

WO 00/08209 PCT/IB99/01444 12

modification at one position. Preferably, the sequences defining a biallelic marker include a polymorphic base selected from the group consisting of the biallelic markers A1 to A19 and the complements thereof. In some embodiments the sequences defining a biallelic marker comprise one of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19 and the 5 complementary sequences thereto. Likewise, the term "marker" or "biallelic marker" requires that the sequence is of sufficient length to practically (although not necessarily unambiguously) identify the polymorphic allele, which usually implies a length of at least 4, 5, 6, 10, 15, 20, 25, or 40 nucleotides.

The term "upstream" is used herein to refer to a location which is toward the 5' end of the polynucleotide from a specific reference point.

The terms "base paired" and "Watson & Crick base paired" are used interchangeably herein to refer to nucleotides which can be hydrogen bonded to one another be virtue of their sequence identities in a manner like that found in double-helical DNA with thymine or uracil residues linked to adenine residues by two hydrogen bonds and cytosine and guanine residues linked by three hydrogen bonds (See Stryer, L., Biochemistry, 4th edition, 1995).

The terms "complementary" or "complement thereof" are used herein to refer to the sequences of polynucleotides which is capable of forming Watson & Crick base pairing with another specified polynucleotide throughout the entirety of the complementary region. For the purpose of the present invention, a first polynucleotide is deemed to be complementary to a second polynucleotide when each base in the first polynucleotide is paired with its complementary base. Complementary bases are, generally, A and T (or A and U), or C and G. "Complement" is used herein as a synonym from "complementary polynucleotide", "complementary nucleic acid" and "complementary nucleotide sequence". These terms are applied to pairs of polynucleotides based solely upon their sequences and not any particular set of conditions under which the two 25 polynucleotides would actually bind.

Variants and fragments

1. Polynucleotides

The invention also relates to variants and fragments of the polynucleotides described herein, particularly of a TBC-1 gene containing one or more biallelic markers according to the invention.

Variants of polynucleotides, as the term is used herein, are polynucleotides that differ from a reference polynucleotide. A variant of a polynucleotide may be a naturally occurring variant such as a naturally occurring allelic variant, or it may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the polynucleotide may be made by mutagenesis techniques, including those applied to polynucleotides, cells or organisms. Generally, differences 35 are limited so that the nucleotide sequences of the reference and the variant are closely similar overall and, in many regions, identical.

WO 00/08209 PCT/IB99/01444

Variants of polynucleotides according to the invention include, without being limited to, nucleotide sequences that are at least 95% identical to any of SEQ ID Nos 1-4 or the sequences complementary thereto or to any polynucleotide fragment of at least 8 consecutive nucleotides of any of SEQ ID Nos 1-4 or the sequences complementary thereto, and preferably at least 98% identical, more particularly at least 99.5% identical, and most preferably at least 99.9% identical to any of SEQ ID Nos 1-4 or the sequences complementary thereto or to any polynucleotide fragment of at least 8 consecutive nucleotides of any of SEQ ID Nos 1-4 or the sequences complementary thereto.

Changes in the nucleotide of a variant may be silent, which means that they do not alter the 10 amino acids encoded by the polynucleotide.

However, nucleotide changes may also result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence. The substitutions, deletions or additions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions.

In the context of the present invention, particularly preferred embodiments are those in which the polynucleotides encode polypeptides which retain substantially the same biological function or activity as the mature TBC-1 protein.

A polynucleotide fragment is a polynucleotide having a sequence that entirely is the same as part but not all of a given nucleotide sequence, preferably the nucleotide sequence of a TBC-1 gene, and variants thereof. The fragment can be a portion of an exon or of an intron of a TBC-1 gene. It can also be a portion of the regulatory sequences of the TBC-1 gene. Preferably, such fragments comprise the polymorphic base of a biallelic marker selected from the group consisting of the biallelic markers A1 to/A19 and the complements thereof.

Such fragments may be "free-standing", i.e. not part of or fused to other polynucleotides, or they may be comprised within a single larger polynucleotide of which they form a part or region. However, several fragments may be comprised within a single larger polynucleotide.

As representative examples of polynucleotide fragments of the invention, there may be mentioned those which have from about 4, 6, 8, 15, 20, 25, 40, 10 to 20, 10 to 30, 30 to 55, 50 to 100, 75 to 100 or 100 to 200 nucleotides in length. Preferred are those fragments having about 49 nucleotides in length, such as those of P1 to P7, P9 to P13, P15 to P19 or the sequences complementary thereto and containing at least one of the biallelic markers of a *TBC-1* gene which are described herein.

2. Polypeptides.

The invention also relates to variants, fragments, analogs and derivatives of the polypeptides described herein, including mutated TBC-1 proteins.

30

10

WO 00/08209 PCT/IB99/01444 14

The variant may be 1) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or 2) one in which one or more of the amino acid residues includes a substituent group, or 3) one in which the mutated TBC-1 is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or 4) one in which the additional amino acids are fused to the mutated TBC-1, such as a leader or secretory sequence or a sequence which is employed for purification of the mutated TBC-1 or a preprotein sequence. Such variants are deemed to be within the scope of those skilled in the art.

More particularly, a variant TBC-1 polypeptide comprises amino acid changes ranging from 1, 2, 3, 4, 5, 10 to 20 substitutions, additions or deletions of one aminoacid, preferably from 1 to 10, more preferably from 1 to 5 and most preferably from 1 to 3 substitutions, additions or deletions of one amino acid. The preferred amino acid changes are those which have little or no influence on the biological activity or the capacity of the variant TBC-1 polypeptide to be recognized by antibodies raised against a native TBC-1 protein.

By homologous peptide according to the present invention is meant a polypeptide containing one or several aminoacid additions, deletions and/or substitutions in the amino acid sequence of a TBC-1 polypeptide. In the case of an aminoacid substitution, one or several consecutive or non-consecutive- aminoacids are replaced by « equivalent » aminoacids.

The expression "equivalent" amino acid is used herein to designate any amino acid that may be substituted for one of the amino acids having similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. Generally, the following groups of amino acids represent equivalent changes: (1) Ala, Pro, Gly, Glu, Asp, Gln, Asn, Ser, Thr; (2) Cys, Ser, Tyr, Thr; (3) Val, Ile, Leu, 25 Met, Ala, Phe; (4) Lys, Arg, His; (5) Phe, Tyr, Trp, His.

By an equivalent aminoacid according to the present invention is also meant the replacement of a residue in the L-form by a residue in the D form or the replacement of a Glutamic acid (E) residue by a Pyro-glutamic acid compound. The synthesis of peptides containing at least one residue in the D-form is, for example, described by Koch (1977).

A specific, but not restrictive, embodiment of a modified peptide molecule of interest according to the present invention, which consists in a peptide molecule which is resistant to proteolysis, is a peptide in which the -CONH- peptide bond is modified and replaced by a (CH₂NH) reduced bond, a (NHCO) retro inverso bond, a (CH₂-O) methylene-oxy bond, a (CH₂-S) thiomethylene bond, a (CH₂CH₂) carba bond, a (CO-CH₂) cetomethylene bond, a (CHOH-CH₂) 35 hydroxyethylene bond), a (N-N) bound, a E-alcene bond or also a -CH=CH- bond.

The polypeptide according to the invention could have post-translational modifications. For example, it can present the following modifications: acylation, disulfide bond formation, prenylation, carboxymethylation and phosphorylation.

A polypeptide fragment is a polypeptide having a sequence that entirely is the same as part but not all of a given polypeptide sequence, preferably a polypeptide encoded by a TBC-1 gene and variants thereof. Preferred fragments include those regions possessing antigenic properties and which can be used to raise antibodies against the TBC-1 protein.

Such fragments may be "free-standing", i.e. not part of or fused to other polypeptides, or they may be comprised within a single larger polypeptide of which they form a part or region.

10 However, several fragments may be comprised within a single larger polypeptide.

As representative examples of polypeptide fragments of the invention, there may be mentioned those which comprise at least about 5, 6, 7, 8, 9 or 10 to 15, 10 to 20, 15 to 40, or 30 to 55 amino acids of the TBC-1. In some embodiments, the fragments contain at least one amino acid mutation in the TBC-1 protein.

Identity Between Nucleic Acids Or Polypeptides

The terms "percentage of sequence identity" and "percentage homology" are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result 25 by 100 to yield the percentage of sequence identity. Homology is evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman, 1988; Altschul et al., 1990; Thompson et al., 1994; Higgins et al., 1996; Altschul et al., 1993). In a particularly preferred embodiment, protein and nucleic acid 30 sequence homologies are evaluated using the Basic Local Alignment Search Tool ("BLAST") which is well known in the art (see, e.g., Karlin and Altschul, 1990; Altschul et al., 1990, 1993, 1997). In particular, five specific BLAST programs are used to perform the following task:

- (1) BLASTP and BLAST3 compare an amino acid query sequence against a protein sequence database;
- 35 (2) BLASTN compares a nucleotide query sequence against a nucleotide sequence database;

WO 00/08209 PCT/IB99/01444 16

(3) BLASTX compares the six-frame conceptual translation products of a query nucleotide sequence (both strands) against a protein sequence database;

- (4) TBLASTN compares a query protein sequence against a nucleotide sequence database translated in all six reading frames (both strands); and
- 5 (5) TBLASTX compares the six-frame translations of a nucleotide query sequence against the six-frame translations of a nucleotide sequence database.

The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as "high-scoring segment pairs," between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database.

10 High-scoring segment pairs are preferably identified (i.e., aligned) by means of a scoring matrix. many of which are known in the art. Preferably, the scoring matrix used is the BLOSUM62 matrix (Gonnet et al., 1992; Henikoff and Henikoff, 1993). Less preferably, the PAM or PAM250 matrices may also be used (see, e.g., Schwartz and Dayhoff, eds., 1978). The BLAST programs evaluate the statistical significance of all high-scoring segment pairs identified, and preferably 15 selects those segments which satisfy a user-specified threshold of significance, such as a userspecified percent homology. Preferably, the statistical significance of a high-scoring segment pair is evaluated using the statistical significance formula of Karlin (see, e.g., Karlin and Altschul, 1990).

Stringent Hybridization Conditions

By way of example and not limitation, procedures using conditions of high stringency are as follows: Prehybridization of filters containing DNA is carried out for 8 h to overnight at 65°C in buffer composed of 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 μg/ml denatured salmon sperm DNA. Filters are hybridized for 48 h at 65°C, the preferred hybridization temperature, in prehybridization mixture containing 100 µg/ml 25 denatured salmon sperm DNA and 5-20 X 10⁶ cpm of ³²P-labeled probe. Alternatively, the hybridization step can be performed at 65°C in the presence of SSC buffer, 1 x SSC corresponding to 0.15M NaCl and 0.05 M Na citrate. Subsequently, filter washes can be done at 37°C for 1 h in a solution containing 2 x SSC, 0.01% PVP, 0.01% Ficoll, and 0.01% BSA, followed by a wash in 0.1 X SSC at 50°C for 45 min. Alternatively, filter washes can be performed in a solution containing 2 30 x SSC and 0.1% SDS, or 0.5 x SSC and 0.1% SDS, or 0.1 x SSC and 0.1% SDS at 68°C for 15 minute intervals. Following the wash steps, the hybridized probes are detectable by autoradiography. Other conditions of high stringency which may be used are well known in the art and as cited in Sambrook et al., 1989; and Ausubel et al., 1989, are incorporated herein in their entirety. These hybridization conditions are suitable for a nucleic acid molecule of about 20 35 nucleotides in length. There is no need to say that the hybridization conditions described above are to be adapted according to the length of the desired nucleic acid, following techniques well known

to the one skilled in the art. The suitable hybridization conditions may for example be adapted

WO 00/08209 PCT/IB99/01444

according to the teachings disclosed in the book of Hames and Higgins (1985) or in Sambrook et al.(1989).

Candidate Region On The Chromosome 4 (Linkage Analysis).

In order to localize the prostate cancer gene(s) starting from families, a systematic familial study of genetic link research is carried out using markers of the microsatellite type described at the Genethon laboratory by the Jean Weissenbach team (Dib et al., 1996).

The studies of genetic link or of "linkage" are based on the principle according to which two neighboring sequences on a chromosome do not present (or very rarely present) recombinations by crossing-over during meiosis. To do this, microsatellite DNA sequences (chromosomal markers) 10 constantly co-inherited with the disease studied are searched for in a family having a predisposition for this disease. These DNA sequences organized in the form of a repetition of di-, tri- or tetranucleotides are systematically present along the genome, and thus allow the identification of chromosomal fragments harboring them. More than 5000 microsatellite markers, have been localized with precision on the genome as a result of the first studies on the genetic map carried out at Genethon under the supervision of Jean Weissenbach, and on the physical map (using the "Yeast Artificial Chromosomes"), work conducted by Daniel Cohen at C.E.P.H. and at Genethon (Chumakov et al., 1995). Genetic link analysis calculates the probabilities of recombinations of the target gene with the chromosomal markers used, according to the genealogical tree, the transmission of the disease, and the transmission of the markers. Thus if a particular allele of a given marker is transmitted with the disease more often than chance would have it (recombination level of between 0 and 0.5), it is possible to deduce that the target gene in question is found in the neighborhood of the marker. Using this technique, it has been possible to localize several genes of genetic predisposition to familial cancers. In order to be able to be included in a genetic link study, the families affected by a hereditary form of the disease must satisfy the "informativeness" criteria: 25 several affected subjects (and whose constitutional DNA is available) per generation, and at best having a large number of siblings.

By linkage analysis, the inventors have identified a candidate region for prostate cancer on chromosome 4. Indeed, the LOD scores at 2 points between the disease and the markers on a total population of approximately fifty families present a value of 2.49 for marker D4S398 which indicates a probable genetic link with this marker. The curve of the variation of the LOD score on a map of 5 markers is centered on D4S398 and the value higher than 3.3 indicates that a gene involved in familial prostate cancer is probably found in the region located between markers D4S2978 and D4S3018, or a space of approximately 9.7 cM.

WO 00/08209 PCT/IB99/01444

Homologies Of The Novel Human Gene Translation Product With A Known Murine Protein.

A novel human gene was found in this candidate region. It presents a good probability to be involved in cancer. Database homology searches have allowed the inventors to determine that the translation product of this novel human gene has significant identity with a murine protein called 5 tbc1. Therefore, the novel human gene of the invention has thus been called *TBC-1* throughout the present specification. *TBC-1* comprises an open Reading frame that encodes a novel protein, the TBC-1 protein. Based on sequence similarity, an alignment of a portion of the TBC-1 amino acid sequence with the known tbc1 murine protein, it is expected that TBC1 protein may play a role in the cell cycle and in differentiation of various tissues. Indeed, the TBC1 protein contains a 200 amino acid domain called the TBC domain that is homologous to regions in the tre2-oncogene and in the yeast regulators of mitosis BUB2 and cdc16.

The cDNA of the murine *tbc1* gene has been described in US Patent No US 5,700,927 and it encodes a putative protein product of 1141 amino acids. The N-terminus of the murine tbc1 protein contains stretches of cysteines and histidines which may form zinc finger structures in the mature polypeptides. The N-terminus also comprises short stretches of basic amino acids which may be involved in a nuclear localization signal. The TBC domain of the murine tbc1 protein contains several tyrosine residues which are conserved in BUB2 and cdc16. The C-terminus of the murine tbc1 protein contains a long stretch of evenly spaced leucine residues which are susceptible to form a leucine zipper motif.

The murine *tbc1* gene has been shown to be highly expressed in testis and kidney. However, lower levels of expression have also be identified in lung, spleen, brain, and heart. Moreover, murine tbc1 is a nuclear protein which is expressed in a cell- and stage-specific manner.

Studies of murine bone marrow have demonstrated that erythroid cells and megakaryocytes expressed substantial levels of the murine tbc1 protein, but none was detected in mature neutrophils.

25 Similarly, spermatogonia do not express murine tbc1, but primary and secondary spermatocytes express abundant tbc1. Later in the differentiation of the germ cells, the tbc1 levels appear to decrease in spermatids and active sperm. The differentiation program of spermatogonia to spermatocytes therefore involves a significant upregulation of murine tbc1 expression.

The general distribution of murine *tbc1* is not tissue-specific, but is cell-specific within individual tissues and intimately linked to tissue differentiation. The developmental expression of murine *tbc1*, particularly in hematopoietic and germ cells, suggests that this gene plays a role in the terminal differentiation program of several tissues.

Consequently, an alteration in the expression of the *TBC-1* gene or in the amino acid sequence of the *TBC-1* protein leading to an altered biological activity of the latter is likely to cause, directly or indirectly, cell proliferation disorders and thus diseases related to an abnormal cell proliferation such as cancer, particularly prostate cancer.

WO 00/08209 PCT/IB99/01444

Genomic Sequence Of TBC-1

The present invention concerns the genomic sequence of *TBC-1*. The present invention encompasses the *TBC-1* gene, or *TBC-1* genomic sequences consisting of, consisting essentially of, or comprising a sequence selected from the group consisting of SEQ ID Nos 1 and 2, a sequence complementary thereto, as well as fragments and variants thereof. These polynucleotides may be purified, isolated, or recombinant.

The inventors have sequenced two portions of the *TBC-1* genomic sequence. The first portion of the *TBC-1* gene sequence contains the three first exons of the *TBC-1* gene, designated as Exon 1, Exon 1bis and Exon 2, and the 5' regulatory sequence located upstream of the transcribed sequences. The sequence of the first portion of the genomic sequence is disclosed in SEQ ID No 1. The second portion contains the twelve last exons of the *TBC-1* gene, designated as exons A, B, C, D, E, F, G, H, I, J, K, and L, and the 3' regulatory sequence which is located downstream of the transcribed sequences.

The exon positions in SEQ ID Nos 1 and 2 are detailed below in Table A.

Table A

Exon	Position in SEQ ID No 1		Intron	Position in S	EQ ID No 1
	Beginning	End		Beginning	End
1	2001	2077	1	2078	12739
1bis	12292	12373	1 bis	12374	12739
2	12740	13249	2	13250	at least
					17590
Exon	Position in SEQ ID No 2		Intron	Position in S	EQ ID No 2
	Beginning	End		Beginning	End
Α	4661	4789	Α	4790	6115
В `	6116	6202	В	6203	9918
С	/9919	10199	C	10200	14520
D	14521	14660	D	14661	50256
E	50257	50442	Е	50443	56255
F	56256	56417	F	56418	63325
G	63326	63484	G	63485	76035
H	76036	76280	Н	76281	78363
I	78364	78523	I	78524	85294
J	85295	85464	J	85465	93416
K	93417	93590	K	93591	97475
I.	97476	97960			

Intron 1 refers to the nucleotide sequence located between Exon 1 and Exon 2; Intron 1 bis refers to the nucleotide sequence located between Exon 1 bis and Exon 2; Intron A refers to the nucleotide sequence located between Exon A and Exon B; and so on. The position of the introns is detailed in Table A.

30

WO 00/08209 PCT/IB99/01444

The TBC-1 introns defined hereinafter for the purpose of the present invention are not exactly what is generally understood as "introns" by the one skilled in the art and will consequently be further defined below.

Generally, an intron is defined as a nucleotide sequence that is present both in the genomic

5 DNA and in the unspliced mRNA molecule, and which is absent from the mRNA molecule which
has already gone through splicing events. In the case of the *TBC-1* gene, the inventors have found
that at least two different spliced mRNA molecules are produced when this gene is transcribed, as it
will be described in detail in a further section of the specification. The first spliced mRNA molecule
comprises Exons 1 and 2. Thus, the genomic nucleotide sequence comprised between Exon 1 and
10 Exon 2 is an intronic sequence as regards to this first mRNA molecule, despite the fact that this
intronic sequence contains Exon 1bis. In contrast, Exon 1bis is of course an exonic nucleotide
sequence as regards to the second *TBC-1* mRNA molecule.

For the purpose of the present invention and in order to make a clear and unambiguous designation of the different nucleic acids encompassed, it has been postulated that the

15 polynucleotides contained both in any of the nucleotide sequences of SEQ ID Nos 1 or 2 and in any of the nucleotide sequences of SEQ ID Nos 3 or 4 are considered as exonic sequences. Conversely, the polynucleotides contained in any of the nucleotide sequences of SEQ ID Nos 1 or 2 but which are absent both from the nucleotide sequence of SEQ ID No 3 and from the nucleotide sequence of SEQ ID No 4 are considered as intronic sequences.

The nucleic acids defining the *TBC-1* introns described above, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a copy of the *TBC-1* gene in a test sample, or alternatively in order to amplify a target nucleotide sequence within the *TBC-1* intronic sequences.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the 15 exons of the *TBC-1* gene which are described in the present invention, or a sequence complementary thereto. The invention also deals with purified, isolated, or recombinant nucleic acids comprising a combination of at least two exons of the *TBC-1* gene, wherein the polynucleotides are arranged within the nucleic acid, from the 5'-end to the 3'-end of said nucleic acid, in the same order as in SEQ ID Nos 1 and 2.

Thus, the invention embodies purified, isolated, or recombinant polynucleotides comprising a nucleotide sequence selected from the group consisting of the introns of the *TBC-1* gene, or a sequence complementary thereto.

The invention also encompasses a purified, isolated, or recombinant polynucleotide comprising a nucleotide sequence having at least 70, 75, 80, 85, 90, or 95% nucleotide identity with a sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a fragment thereof. The nucleotide differences as regards to the nucleotide sequence of SEQ ID Nos 1 or 2 may be generally randomly distributed throughout the entire nucleic acid.

Nevertheless, preferred nucleic acids are those wherein the nucleotide differences as regards to the nucleotide sequence of SEQ ID Nos 1 or 2 are predominantly located outside the coding sequences contained in the exons. These nucleic acids, as well as their fragments and variants, may be used as oligonucleotide primers or probes in order to detect the presence of a copy of the *TBC-1* gene in a test sample, or alternatively in order to amplify a target nucleotide sequence within the *TBC-1* sequences.

Another object of the invention consists of a purified, isolated, or recombinant nucleic acid that hybridizes with a sequence selected from the group consisting of SEQ ID Nos 1 and 2 or a complementary sequence thereto or a variant thereof, under the stringent hybridization conditions as defined above.

Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 1 and 2, or the complements thereof. Additionally preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001-9000, 9001-10000, 10001-11000, 11001-12000, 12001-13000, 13001-14000, 14001-15000, 15001-16000, 16001-17000, and 17001-17590. Other preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the 25 following nucleotide positions of SEO ID No 2: 1-5000, 5001-10000, 10001-15000, 15001-20000, 20001-25000, 25001-30000, 30001-35000, 35001-40000, 40001-45000, 45001-50000, 50001-55000, 55001-60000, 60001-65000, 65001-70000, 70001-75000, 75001-80000, 80001-85000, 85001-90000, 90001-95000, and 95001-99960.

While this section is entitled "Genomic Sequences of *TBC-1*," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of *TBC-1* on either side or between two or more such genomic sequences.

TBC-1 cDNA Sequences

The inventors have discovered that the expression of the *TBC-1* gene leads to the production of at least two mRNA molecules, respectively a first and a second *TBC-1* transcription

WO 00/08209 PCT/IB99/01444 22

product, as the results of alternative splicing events. They result from two distinct first exons, namely Exon 1 and Exon 1 bis.

The first transcription product comprises Exons 1, 2, A, B, C, D, E, F, G, H, I, J, K, and L. This cDNA of SEQ ID No 3 includes a 5'-UTR region, spanning the whole Exon 1 and part of

5 Exon 2. This 5'-UTR region starts from the nucleotide at position 1 and ends at the nucleotide at position 170 of the nucleotide sequence of SEQ ID No 3. The cDNA of SEQ ID No 3 includes a 3'
UTR region starting from the nucleotide at position 3726 and ending at the nucleotide at position 3983 of the nucleotide sequence of SEQ ID No 3. This first transcription product harbors a polyadenylation signal located between the nucleotide at position 3942 and the nucleotide at position 3947 of the nucleotide sequence of SEQ ID No 3.

The second *TBC-1* transcription product comprises Exons 1*bis*, 2, A, B, C, D, E, F, G, H, I, J, K, and L. This cDNA of SEQ ID No 4 includes a 5'-UTR region starting from the nucleotide at position 1 and ending at the nucleotide at position 175 of the nucleotide sequence of SEQ ID No 4. This second cDNA also includes a 3'-UTR region starting from the nucleotide at position 3731 and ending at the nucleotide at position 3988 of the nucleotide sequence of SEQ ID No 4. This second transcription product harbors a polyadenylation signal located between the nucleotide at position 3947 and the nucleotide at position 3952 of the nucleotide sequence of SEQ ID No 4.

The 5'-end sequence of this second *TBC-1* mRNA, more particularly the nucleotide sequence comprised between the nucleotide in position 1 and the nucleotide in position 458 of the nucleic acid of SEQ ID No 4 molecule corresponds to the nucleotide sequence of a 5'-EST that has been obtained from a human pancreas cDNA library and characterized following the teachings of the PCT Application No WO 96/34981. This 5'-EST is also part of the invention.

Another object of the invention consists of a purified or isolated nucleic acid comprising a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 3 and 25 4 and to nucleic acid fragments thereof.

Preferred nucleic acid fragments of the nucleotide sequences of SEQ ID Nos 3 and 4 consist in polynucleotides comprising their respective Open Reading Frames encoding the TBC-1 protein.

Other preferred nucleic acid fragments of the nucleotide sequences of SEQ ID Nos 3 and 4 consist in polynucleotides comprising at least a part of their respective 5'-UTR or 3'-UTR regions.

The invention also pertains to a purified or isolated nucleic acid having at least a 95% of nucleotide identity with any one of the nucleotide sequences of SEQ ID Nos 3 and 4, or a fragment thereof.

Another object of the invention consists of purified, isolated or recombinant nucleic acids comprising a polynucleotide that hybridizes, under the stringent hybridization conditions defined herein, with any one of the nucleotide sequences of SEQ ID Nos 3 and 4, or a sequence complementary thereto or a fragment thereof.

WO 00/08209 PCT/IB99/01444 23

The invention also relates to isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 3 and 4, or the complements thereof. Particularly preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3983. Additionally preferred nucleic acids of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 4: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3988. Such a nucleic acid is notably useful as polynucleotide probe or primer specific for the *TBC-1* gene or the *TBC-1* mRNAs and cDNAs.

While this section is entitled "TBC-1 cDNA Sequences," it should be noted that nucleic acid fragments of any size and sequence may also be comprised by the polynucleotides described in this section, flanking the genomic sequences of TBC-1 on either side or between two or more such genomic sequences.

Coding Regions

The *TBC-1* open reading frame is contained in the two *TBC-1* mRNA molecules of about 4 kilobases isolated by the inventors.

More precisely, the effective *TBC-1* coding sequence is comprised between the nucleotide at position 171 and the nucleotide at position 3725 of SEQ ID No 3, and between the nucleotide at position 176 and the nucleotide at position 3730 of the nucleotide sequence of SEQ ID No 4.

The invention further provides a purified or isolated nucleic acid comprising a polynucleotide selected from the group consisting of a polynucleotide comprising a nucleic acid sequence located between the nucleotide at position 171 and the nucleotide at position 3725 of SEQ ID No 3, and a polynucleotide comprising a nucleic acid sequence located between the nucleotide at position 176 and the nucleotide at position 3730 of SEQ ID No 4 or a variant or fragment thereof or a sequence complementary thereto.

The present invention concerns a purified or isolated nucleic acid encoding a human TBC-1 protein, wherein said TBC-1 protein comprises an amino acid sequence of SEQ ID No 5, a nucleotide sequence complementary thereto, a fragment or a variant thereof. The present invention also embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids,

WO 00/08209 PCT/IB99/01444

more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 5. In a preferred embodiment, the present invention embodies isolated, purified, and recombinant polynucleotides which encode a polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 or 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, or 100 amino acids of SEQ ID No 5 wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the following amino acid positions in SEQ ID No 5: 1-300, 301-600, 601-900, and 901-1168.

The above disclosed polynucleotide that contains only coding sequences derived from the *TBC-1* ORF may be expressed in a desired host cell or a desired host organism, when said polynucleotide is placed under the control of suitable expression signals. Such a polynucleotide, when placed under the suitable expression signals, may be inserted in a vector for its expression.

Regulatory Sequences Of TBC-1

The invention further deals with a purified or isolated nucleic acid comprising the nucleotide sequence of a regulatory region which is located either upstream of the first exon of the *TBC-1* gene and which is contained in the *TBC-1* genomic sequence of SEQ ID No 1, or downstream of the last exon of the *TBC-1* gene and which is contained in the *TBC-1* genomic sequence of SEQ ID No 2.

The 5'-regulatory sequence of the *TBC-1* gene is localized between the nucleotide in position 1 and the nucleotide in position 2000 of the nucleotide sequence of SEQ ID No 1. The 3'-regulatory sequence of the *TBC-1* gene is localized between nucleotide position 97961 and nucleotide position 99960 of SEQ ID No 2.

Polynucleotides derived from the 5' and 3' regulatory regions are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1 or 2 or a fragment thereof in a test sample.

The promoter activity of the 5' regulatory regions contained in *TBC-1* can be assessed as described below.

Genomic sequences lying upstream of the *TBC-1* Exons are cloned into a suitable promoter reporter vector, such as the pSEAP-Basic, pSEAP-Enhancer, pβgal-Basic, pβgal-Enhancer, or pEGFP-1 Promoter Reporter vectors available from Clontech. Briefly, each of these promoter reporter vectors include multiple cloning sites positioned upstream of a reporter gene encoding a readily assayable protein such as secreted alkaline phosphatase, beta galactosidase, or green fluorescent protein. The sequences upstream of the *TBC-1* coding region are inserted into the cloning sites upstream of the reporter gene in both orientations and introduced into an appropriate host cell. The level of reporter protein is assayed and compared to the level obtained from a vector which lacks an insert in the cloning site. The presence of an elevated expression level in the vector containing the insert with respect to the control vector indicates the presence of a promoter in the insert. If necessary, the upstream sequences can be cloned into vectors which contain an enhancer

WO 00/08209 PCT/IB99/01444

for increasing transcription levels from weak promoter sequences. A significant level of expression above that observed with the vector lacking an insert indicates that a promoter sequence is present in the inserted upstream sequence.

Promoter sequences within the upstream genomic DNA may be further defined by

5 constructing nested deletions in the upstream DNA using conventional techniques such as

Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter

reporter vector to determine whether the deletion has reduced or obliterated promoter activity. In

this way, the boundaries of the promoters may be defined. If desired, potential individual regulatory

sites within the promoter may be identified using site directed mutagenesis or linker scanning to

10 obliterate potential transcription factor binding sites within the promoter, individually or in

combination. The effects of these mutations on transcription levels may be determined by inserting

the mutations into the cloning sites in the promoter reporter vectors.

Thus, the minimal size of the promoter of the *TBC-1* gene can be determined through the measurement of *TBC-1* expression levels. For this assay, an expression vector comprising decreasing sizes from the promoter generally ranging from 2 kb to 100 bp, with a 3' end which is constant, operably linked to *TBC-1* coding sequence or to a reporter gene is used. Cells, which are preferably prostate cells and more preferably prostate cancer cells, are transfected with this vector and the expression level of the gene is assessed.

The strength and the specificity of the promoter of the *TBC-1* gene can be assessed through the expression levels of the gene operably linked to this promoter in different types of cells and tissues. In one embodiment, the efficacy of the promoter of the *TBC-1* gene is assessed in normal and cancer cells. In a preferred embodiment, the efficacy of the promoter of the *TBC-1* gene is assessed in normal prostate cells and in prostate cancer cells which can present different degrees of malignancy.

Polynucleotides carrying the regulatory elements located both at the 5' end and at the 3' end of the *TBC-1* cDNAs may be advantageously used to control the transcriptional and translational activity of an heterologous polynucleotide of interest.

Thus, the present invention also concerns a purified or isolated nucleic acid comprising a polynucleotide which is selected from the group consisting of the 5' and 3' regulatory regions, or a sequence complementary thereto or a biologically active fragment or variant thereof. "5' regulatory region" refers to the nucleotide sequence located between positions 1 and 2000 of SEQ ID No 1. "3' regulatory region" refers to the nucleotide sequence located between positions 97961 and 99960 of SEQ ID No 2.

The invention also pertains to a purified or isolated nucleic acid comprising a

35 polynucleotide having at least 95% nucleotide identity with a polynucleotide selected from the
group consisting of the 5' and 3' regulatory regions, advantageously 99 % nucleotide identity,
preferably 99.5% nucleotide identity and most preferably 99.8% nucleotide identity with a

25

WO 00/08209 PCT/IB99/01444 26

polynucleotide selected from the group consisting of the 5' and 3' regulatory regions, or a sequence complementary thereto or a variant thereof or a biologically active fragment thereof.

Another object of the invention consists of purified, isolated or recombinant nucleic acids comprising a polynucleotide that hybridizes, under the stringent hybridization conditions defined 5 herein, with a polynucleotide selected from the group consisting of the nucleotide sequences of the 5'- and 3' regulatory regions, or a sequence complementary thereto or a variant thereof or a biologically active fragment thereof.

The 5'UTR and 3'UTR regions of a gene are of particular importance in that they often comprise regulatory elements which can play a role in providing appropriate expression levels, particularly through the control of mRNA stability.

A 5' regulatory polynucleotide of the invention may include the 5'-UTR located between the nucleotide at position 1 and the nucleotide at position 170 of SEQ ID No 3, or a biologically active fragment or variant thereof.

Alternatively, a 5'-regulatory polynucleotide of the invention may include the 5'-UTR

15 located between the nucleotide at position 1 and the nucleotide at position 175 of SEQ ID No 4, or a biologically active fragment or variant thereof.

A 3' regulatory polynucleotide of the invention may include the 3'-UTR located between the nucleotide at position 3726 and the nucleotide at position 3983 of SEQ ID No 4, or a biologically active fragment or variant thereof.

Thus, the invention also pertains to a purified or isolated nucleic acid which is selected from the group consisting of:

- a) a nucleic acid comprising the nucleotide sequence of the 5' regulatory region;
- b) a nucleic acid comprising a biologically active fragment or variant of the nucleic acid of the 5' regulatory region.

Preferred fragments of the nucleic acid of the 5' regulatory region have a length of about 1000 nucleotides, more particularly of about 400 nucleotides, more preferably of about 200 nucleotides and most preferably about 100 nucleotides. More particularly, the invention further includes specific elements within this regulatory region, these elements preferably including the promoter region.

30 Preferred fragments of the 3' regulatory region are at least 50, 100, 150, 200, 300 or 400 bases in length.

By a "biologically active fragment or variant" of a *TBC-1* regulatory polynucleotide according to the present invention is intended a polynucleotide comprising or alternatively consisting in a fragment of said polynucleotide which is functional as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide in a recombinant cell host.

For the purpose of the invention, a nucleic acid or polynucleotide is "functional" as a regulatory region for expressing a recombinant polypeptide or a recombinant polynucleotide if said

WO 00/08209 PCT/IB99/01444

regulatory polynucleotide contains nucleotide sequences which contain transcriptional and translational regulatory information, and if such sequences are "operatively linked" to nucleotide sequences which encode the desired polypeptide or the desired polynucleotide. An operable linkage is a linkage in which the regulatory nucleic acid and the DNA sequence sought to be expressed are linked in such a way as to permit gene expression.

In order, to identify the relevant biologically active polynucleotide derivatives of the 5' or 3' regulatory region, the one skill in the art will refer to the book of Sambrook et al. (Sambrook, 1989) in order to use a recombinant vector carrying a marker gene (i.e. beta galactosidase, chloramphenicol acetyl transferase, etc.) the expression of which will be detected when placed under the control of a biologically active derivative polynucleotide of the 5' or 3' regulatory region.

Regulatory polynucleotides of the invention may be prepared from any of the nucleotide sequences of SEQ ID Nos 1 or 2 by cleavage using the suitable restriction enzymes, the one skill in the art being guided by the book of Sambrook et al. (1989). Regulatory polynucleotides may also be prepared by digestion of any of the nucleotide sequences of SEQ ID Nos 1 or 2 by an exonuclease enzyme, such as Bal31 (Wabiko et al., 1986). These regulatory polynucleotides can also be prepared by chemical synthesis, as described elsewhere in the specification, when the synthesis of oligonucleotide probes or primers is disclosed.

The regulatory polynucleotides according to the invention may be advantageously part of a recombinant expression vector that may be used to express a coding sequence in a desired host cell or host organism. The recombinant expression vectors according to the invention are described elsewhere in the specification.

The invention also encompasses a polynucleotide comprising:

- a) a nucleic acid comprising a regulatory nucleotide sequence of the 5' regulatory region, or a biologically active fragment or variant thereof;
- b) a polynucleotide encoding a desired polypeptide or nucleic acid, operably linked to the nucleic acid comprising a regulatory nucleotide sequence of the 5' regulatory region, or its biologically active fragment or variant.
 - c) Optionally, a nucleic acid comprising a 3' regulatory polynucleotide, preferably a 3'regulatory polynucleotide of the invention.

The desired polypeptide encoded by the above described nucleic acid may be of various nature or origin, encompassing proteins of prokaryotic or eukaryotic origin. Among the polypeptides expressed under the control of a *TBC-1* regulatory region, it may be cited bacterial, fungal or viral antigens. Are also encompassed eukaryotic proteins such as intracellular proteins, such as "house keeping" proteins, membrane-bound proteins, like receptors, and secreted proteins like the numerous endogenous mediators such as cytokines.

WO 00/08209 PCT/IB99/01444

The desired nucleic acid encoded by the above described polynucleotide, usually a RNA molecule, may be complementary to a *TBC-1* coding sequence and thus useful as an antisense polynucleotide.

Such a polynucleotide may be included in a recombinant expression vector in order to

5 express a desired polypeptide or a desired polynucleotide in host cell or in a host organism. Suitable recombinant vectors that contain a polynucleotide such as described hereinbefore are disclosed elsewhere in the specification.

TBC-1 Polypeptide And Peptide Fragments Thereof

It is now easy to produce proteins in high amounts by genetic engineering techniques
through expression vectors such as plasmids, phages or phagemids. The polynucleotide that code
for one the polypeptides of the present invention is inserted in an appropriate expression vector in
order to produce the polypeptide of interest *in vitro*.

Thus, the present invention also concerns a method for producing one of the polypeptides described herein, and especially a polypeptide of SEQ ID No 5 or a fragment or a variant thereof, wherein said method comprises the steps of:

- a) culturing, in an appropriate culture medium, a cell host previously transformed or transfected with the recombinant vector comprising a nucleic acid encoding a TBC-1 polypeptide, or a fragment or a variant thereof;
- b) harvesting the culture medium thus conditioned or lyse the cell host, for example by sonication or by an osmotic shock;
- c) separating or purifying, from the said culture medium, or from the pellet of the resultant host cell lysate the thus produced polypeptide of interest.
 - d) Optionally characterizing the produced polypeptide of interest.

In a specific embodiment of the above method, step a) is preceded by a step wherein the

25 nucleic acid coding for a TBC-1 polypeptide, or a fragment or a variant thereof, is inserted in an
appropriate vector, optionally after an appropriate cleavage of this amplified nucleic acid with one
or several restriction endonucleases. The nucleic acid coding for a TBC-1 polypeptide or a fragment
or a variant thereof may be the resulting product of an amplification reaction using a pair of primers
according to the invention (by SDA, TAS, 3SR NASBA, TMA etc.).

The polypeptides according to the invention may be characterized by binding onto an immunoaffinity chromatography column on which polyclonal or monoclonal antibodies directed to a polypeptide of SEQ ID No 5, or a fragment or a variant thereof, have previously been immobilized.

Purification of the recombinant proteins or peptides according to the present invention may be carried out by passage onto a Nickel or Cupper affinity chromatography column. The Nickel chromatography column may contain the Ni-NTA resin (Porath et al., 1975).

The polypeptides or peptides thus obtained may be purified, for example by high performance liquid chromatography, such as reverse phase and/or cationic exchange HPLC, as described by Rougeot et al. (1994). The reason to prefer this kind of peptide or protein purification is the lack of byproducts found in the elution samples which renders the resultant purified protein or peptide more suitable for a therapeutic use.

Another object of the present invention consists in a purified or isolated TBC-1 polypeptide or a fragment or a variant thereof.

In a preferred embodiment, the TBC-1 polypeptide comprises an amino acid sequence of SEQ ID No 5 or a fragment or a variant thereof. The present invention also embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5. The present invention also embodies isolated, purified, and recombinant polypeptides comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5, wherein said contiguous span includes at least 1, 2, 3, 5 or 10 of the following amino acid positions: 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The invention also encompasses a purified, isolated, or recombinant polypeptides comprising an amino acid sequence having at least 90, 95, 98 or 99% amino acid identity with the amino acid sequence of SEQ ID No 5 or a fragment thereof.

The TBC-1 polypeptide of the invention possesses amino acid homologies as regards to the murine TBC-1 protein of 1141 amino acids in length which is described in US Patent No US 5,700,927. The TBC-1 protein of the invention also possesses some homologies with two other proteins: the Pollux drosophila protein (Zhang et al., 1996) and the CDC16 protein from Caenorhabditis elegans (Wilson et al., 1994). Figure 1 represents an amino acid alignment of a portion of the amino acid sequence of the TBC-1 protein of SEQ ID No 5 with other proteins sharing amino acid homology with TBC-1. The upper line shows the whole amino acid sequence of the murine tbc-1 protein described in US Patent No US 5,700,927; the second line represents part of the amino acid sequence of the TBC-1 protein of SEQ ID No 5; the third line (Genbank access No: dmu50542) depicts the amino acid sequence of the Pollux protein mentioned above; the fourth line (Genbank access No: celf35h12) shows the amino acid sequence of the C. elegans protein mentioned above; the fifth line presents positions in which consensus amino acids are identified, i.e. amino acids shared by the sequences presented in the four upper lines, when present.

The TBC-1 polypeptide of the amino acid sequence of SEQ ID No 5 has 1168 amino acids in length. The TBC-1 polypeptide includes a "TBC domain" which is spanning from the amino acid in position 786 to the amino acid in position 974 of the amino acid sequence of SEQ ID No 5. This TBC domain is represented in Figure 1 as a grey area spanning from the amino acid numbered 758 to the amino acid numbered 949. This TBC domain is likely to regulate protein-protein interactions.

WO 00/08209 PCT/IB99/01444

Moreover, the TBC-1 TBC domain includes the amino acid sequence EVGYCQGL, spanning from the amino acid in position 886 to the amino acid in position 893 of the amino acid sequence of SEQ ID No 5. The EVGYCQGL amino acid sequence spans from the amino acid numbered 861 to the amino acid numbered 868 of Figure 1. This site may interact with a kinase. Based on the structural similarity to cdc16, a yeast regulator of mitosis, TBC-1 is likely to regulate mitosis and cytokinesis by interacting with other proteins which also participate with the regulation of mitosis, cytokinesis and septum formation.

Preferred polypeptides of the invention comprise the TBC domain of TBC-1, or alternatively at least the EVGYCQGL amino acid sequence motif.

A further object of the present invention concerns a purified or isolated polypeptide which is encoded by a nucleic acid comprising a nucleotide sequence selected from the group consisting of SEQ ID Nos 1, 2, 3, and 4 or fragments or variants thereof.

A single variant molecule of the TBC-1 protein is explicitly excluded from the scope of the present invention, which is a polypeptide having the same amino acid sequence than the murine tbc1 protein described in the US Patent No 5,700,927.

Amino acid deletions, additions or substitutions in the TBC-1 protein are preferably located outside of the TBC domain as defined above. Most preferably, a mutated TBC-1 protein has an intact "EVGYCQGL" amino acid motif.

Such a mutated TBC-1 protein may be the target of diagnostic tools, such as specific monoclonal or polyclonal antibodies, useful for detecting the mutated TBC-1 protein in a sample.

The invention also encompasses a TBC-1 polypeptide or a fragment or a variant thereof in which at least one peptide bound has been modified as described in the "Definitions" section.

Antibodies That Bind TBC-1 Polypeptides of the Invention

Any TBC-1 polypeptide or whole protein may be used to generate antibodies capable of specifically binding to an expressed TBC-1 protein or fragments thereof as described.

One antibody composition of the invention is capable of specifically binding or specifically bind to the variant of the TBC-1 protein of SEQ ID No 5. For an antibody composition to specifically bind to TBC-1, it must demonstrate at least a 5%, 10%, 15%, 20%, 25%, 50%, or 100% greater binding affinity for TBC-1 protein than for another protein in an ELISA, RIA, or other antibody-based binding assay.

In a preferred embodiment, the invention concerns antibody compositions, either polyclonal or monoclonal, capable of selectively binding, or selectively bind to an epitope-containing a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5; Optionally said epitope comprises at least 1, 2, 3, 5 or 10 of the following amino acid positions: 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The invention also concerns a purified or isolated antibody capable of specifically binding to a mutated TBC-1 protein or to a fragment or variant thereof comprising an epitope of the mutated TBC-1 protein. In another preferred embodiment, the present invention concerns an antibody capable of binding to a polypeptide comprising at least 10 consecutive amino acids of a TBC-1 protein and including at least one of the amino acids which can be encoded by the trait causing mutations.

In a preferred embodiment, the invention concerns the use in the manufacture of antibodies of a polypeptide comprising a contiguous span of at least 6 amino acids, preferably at least 8 to 10 amino acids, more preferably at least 12, 15, 20, 25, 30, 40, 50, 100, 150 or 200 amino acids of SEQ ID No 5; Optionally said polypeptide comprises at least 1, 2, 3, 5 or 10 of the following amino acid positions: 1-200, 201-400, 401-600, 601-800, 801-1000, 1001-1168.

The antibodies of the invention may be labeled by any one of the radioactive, fluorescent or enzymatic labels known in the art.

The TBC-1 polypeptide of SEQ ID No 5 or a fragment thereof can be used for the preparation of polyclonal or monoclonal antibodies.

The TBC-1 polypeptide expressed from a DNA sequence comprising at least one of the nucleic acid sequences of SEQ ID Nos 1, 2, 3 and 4 may also be used to generate antibodies capable of specifically binding to the TBC-1 polypeptide of SEQ ID No 5 or a fragment thereof.

Preferred antibodies according to the invention are prepared using TBC-1 peptide fragments that do not comprise the EVGYCQGL amino acid motif.

Other preferred antibodies of the invention are prepared using TBC-1 peptide fragments that do not comprise the TBC domain defined elsewhere in the specification.

The antibodies may be prepared from hybridomas according to the technique described by Kohler and Milstein in 1975. The polyclonal antibodies may be prepared by immunization of a 25 mammal, especially a mouse or a rabbit, with a polypeptide according to the invention that is combined with an adjuvant of immunity, and then by purifying of the specific antibodies contained in the serum of the immunized animal on a affinity chromatography column on which has previously been immobilized the polypeptide that has been used as the antigen.

The present invention also includes, chimeric single chain Fv antibody fragments (Martineau et al., 1998), antibody fragments obtained through phage display libraries (Ridder et al., 1995; Vaughan et al., 1995) and humanized antibodies (Reinmann et al., 1997; Leger et al., 1997).

Antibody preparations prepared according to either protocol are useful in quantitative immunoassays which determine concentrations of antigen-bearing substances in biological samples; they are also used semi-quantitatively or qualitatively to identify the presence of antigen in a biological sample. The antibodies may also be used in therapeutic compositions for killing cells expressing the protein or reducing the levels of the protein in the body.

10

WO 00/08209 PCT/IB99/01444

Consequently, the invention is also directed to a method for detecting specifically the presence of a TBC-1 polypeptide according to the invention in a biological sample, said method comprising the following steps:

- a) bringing into contact the biological sample with a polyclonal or monoclonal antibody that
 5 specifically binds a TBC-1 polypeptide comprising an amino acid sequence of SEQ ID No 5, or to a peptide fragment or variant thereof; and
 - b) detecting the antigen-antibody complex formed.

The invention also concerns a diagnostic kit for detecting *in vitro* the presence of a TBC-1 polypeptide according to the present invention in a biological sample, wherein said kit comprises:

- a) a polyclonal or monoclonal antibody that specifically binds a TBC-1 polypeptide comprising an amino acid sequence of SEQ ID No 5, or to a peptide fragment or variant thereof, optionally labeled;
- b) a reagent allowing the detection of the antigen-antibody complexes formed, said reagent carrying optionally a label, or being able to be recognized itself by a labeled reagent, more
 particularly in the case when the above-mentioned monoclonal or polyclonal antibody is not labeled by itself.

TBC-1-Related Biallelic Markers

The inventors have discovered nucleotide polymorphisms located within the genomic DNA containing the *TBC-1* gene, and among them SNP that are also termed biallelic markers. The biallelic markers of the invention can be used for example for the generation of genetic map, the linkage analysis, the association studies.

A- Identification Of TBC-1-related Biallelic Markers

There are two preferred methods through which the biallelic markers of the present invention can be generated. In a first method, DNA samples from unrelated individuals are pooled together, following which the genomic DNA of interest is amplified and sequenced. The nucleotide sequences thus obtained are then analyzed to identify significant polymorphisms.

One of the major advantages of this method resides in the fact that the pooling of the DNA samples substantially reduces the number of DNA amplification reactions and sequencing which must be carried out. Moreover, this method is sufficiently sensitive so that a biallelic marker obtained therewith usually shows a sufficient degree of informativeness for conducting association studies.

In a second method for generating biallelic markers, the DNA samples are not pooled and are therefore amplified and sequenced individually. The resulting nucleotide sequences obtained are then also analyzed to identify significant polymorphisms.

It will readily be appreciated that when this second method is used, a substantially higher number of DNA amplification reactions must be carried out. It will further be appreciated that

including such potentially less informative biallelic markers in association studies to identify potential genetic associations with a trait may allow in some cases the direct identification of causal mutations, which may, depending on their penetrance, be rare mutations. This method is usually preferred when biallelic markers need to be identified in order to perform association studies within candidate genes.

In both methods, the genomic DNA samples from which the biallelic markers of the present invention are generated are preferably obtained from unrelated individuals corresponding to a heterogeneous population of known ethnic background, or from familial cases.

The number of individuals from whom DNA samples are obtained can vary substantially,
10 preferably from about 10 to about 1000, preferably from about 50 to about 200 individuals. It is
usually preferred to collect DNA samples from at least about 100 individuals in order to have
sufficient polymorphic diversity in a given population to generate as many markers as possible and
to generate statistically significant results.

As for the source of the genomic DNA to be subjected to analysis, any test sample can be foreseen without any particular limitation. The preferred source of genomic DNA used in the context of the present invention is the peripheral venous blood of each donor.

The techniques of DNA extraction are well-known to the skilled technician. Details of a preferred embodiment are provided in Example 2.

DNA samples can be pooled or unpooled for the amplification step. DNA amplification techniques are well-known to those skilled in the art.

Amplification techniques that can be used in the context of the present invention include, but are not limited to, the ligase chain reaction (LCR) described in EP-A- 320 308, WO 9320227 and EP-A-439 182, the polymerase chain reaction (PCR, RT-PCR) and techniques such as the nucleic acid sequence based amplification (NASBA) described in Guatelli J.C., et al.(1990) and in Compton J.(1991), Q-beta amplification as described in European Patent Application No 4544610, strand displacement amplification as described in Walker et al.(1996) and EP A 684 315 and, target mediated amplification as described in PCT Publication WO 9322461.

LCR and Gap LCR are exponential amplification techniques, both depend on DNA ligase to join adjacent primers annealed to a DNA molecule. In Ligase Chain Reaction (LCR), probe pairs are used which include two primary (first and second) and two secondary (third and fourth) probes, all of which are employed in molar excess to target. The first probe hybridizes to a first segment of the target strand and the second probe hybridizes to a second segment of the target strand, the first and second segments being contiguous so that the primary probes abut one another in 5' phosphate-3'hydroxyl relationship, and so that a ligase can covalently fuse or ligate the two probes into a fused product. In addition, a third (secondary) probe can hybridize to a portion of the first probe and a fourth (secondary) probe can hybridize to a portion of the second probe in a similar abutting fashion. Of course, if the target is initially double stranded, the secondary probes also will

hybridize to the target complement in the first instance. Once the ligated strand of primary probes is separated from the target strand, it will hybridize with the third and fourth probes, which can be ligated to form a complementary, secondary ligated product. It is important to realize that the ligated products are functionally equivalent to either the target or its complement. By repeated cycles of hybridization and ligation, amplification of the target sequence is achieved. A method for multiplex LCR has also been described (WO 9320227). Gap LCR (GLCR) is a version of LCR where the probes are not adjacent but are separated by 2 to 3 bases.

For amplification of mRNAs, it is within the scope of the present invention to reverse transcribe mRNA into cDNA followed by polymerase chain reaction (RT-PCR); or, to use a single enzyme for both steps as described in U.S. Patent No. 5,322,770 or, to use Asymmetric Gap LCR (RT-AGLCR) as described by Marshall et al.(1994). AGLCR is a modification of GLCR that allows the amplification of RNA.

The PCR technology is the preferred amplification technique used in the present invention. A variety of PCR techniques are familiar to those skilled in the art. For a review of PCR technology, see White (1997) and the publication entitled "PCR Methods and Applications" (1991, Cold Spring Harbor Laboratory Press). In each of these PCR procedures, PCR primers on either side of the nucleic acid sequences to be amplified are added to a suitably prepared nucleic acid sample along with dNTPs and a thermostable polymerase such as Taq polymerase, Pfu polymerase, or Vent polymerase. The nucleic acid in the sample is denatured and the PCR primers are specifically hybridized to complementary nucleic acid sequences in the sample. The hybridized primers are extended. Thereafter, another cycle of denaturation, hybridization, and extension is initiated. The cycles are repeated multiple times to produce an amplified fragment containing the nucleic acid sequence between the primer sites. PCR has further been described in several patents including US Patents 4,683,195; 4,683,202; and 4,965,188.

The PCR technology is the preferred amplification technique used to identify new biallelic markers. A typical example of a PCR reaction suitable for the purposes of the present invention is provided in Example 3.

One of the aspects of the present invention is a method for the amplification of a *TBC-1* gene, particularly the genomic sequences of SEQ ID Nos 1 and 2 or of the cDNA sequence of SEQ 30 ID Nos 3 or 4 or a fragment or variant thereof in a test sample, preferably using the PCR technology. The method comprises the steps of contacting a test sample suspected of containing the target *TBC-1* sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers.

Thus, the present invention also relates to a method for the amplification of a TBC-1 gene sequence, particularly of a fragment of the genomic sequence of SEQ ID No 1 or of the cDNA sequence of SEQ ID No 2 or 3, or a fragment or a variant thereof in a test sample, said method comprising the steps of:

WO 00/08209 PCT/IB99/01444

a) contacting a test sample suspected of containing the targeted *TBC-1* gene sequence or portion thereof with amplification reaction reagents comprising a pair of amplification primers located on either side of the *TBC-1* region to be amplified, and

b) optionally, detecting the amplification products.

The invention also concerns a kit for the amplification of a TBC-1 gene sequence, particularly of a portion of the genomic sequence of SEQ ID Nos 1 or 2, or of the cDNA sequence of SEQ ID Nos 3 or 4, or a variant thereof in a test sample, wherein said kit comprises:

- a) a pair of oligonucleotide primers located on either side of the TBC-1 region to be amplified;
- b) optionally, the reagents necessary for performing the amplification reaction.

In one embodiment of the above amplification method and kit, the amplification product is detected by hybridization with a labeled probe having a sequence which is complementary to the amplified region. In another embodiment of the above amplification method and kit, primers comprise a sequence which is selected from the group consisting of B1 to B15, C1 to C15, D1 to D19, and E1 to E19.

In a first embodiment of the present invention, biallelic markers are identified using genomic sequence information generated by the inventors. Sequenced genomic DNA fragments are used to design primers for the amplification of 500 bp fragments. These 500 bp fragments are amplified from genomic DNA and are scanned for biallelic markers. Primers may be designed using the OSP software (Hillier L. and Green P., 1991). All primers may contain, upstream of the specific target bases, a common oligonucleotide tail that serves as a sequencing primer. Those skilled in the art are familiar with primer extensions, which can be used for these purposes.

Preferred primers, useful for the amplification of genomic sequences encoding the candidate genes, focus on promoters, exons and splice sites of the genes. A biallelic marker presents a higher probability to be an eventual causal mutation if it is located in these functional regions of the gene. Preferred amplification primers of the invention include the nucleotide sequences of B1 to B15 and C1 to C15 further detailed in Example 3.

The amplification products generated as described above with the primers of the invention are then sequenced using methods known and available to the skilled technician. Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dyeprimer cycle sequencing protocol. Following gel image analysis and DNA sequence extraction, sequence data are automatically processed with adequate software to assess sequence quality.

A polymorphism analysis software is used that detects the presence of biallelic sites among individual or pooled amplified fragment sequences. Polymorphism search is based on the presence of superimposed peaks in the electrophoresis pattern. These peaks which present distinct colors correspond to two different nucleotides at the same position on the sequence. The polymorphism has to be detected on both strands for validation.

WO 00/08209 PCT/IB99/01444 36

19 biallelic markers were found in the TBC-1 gene. They are detailed in the Table 2. They are located in intronic regions.

B- Genotyping Of TBC-1-Related Biallelic Markers

The polymorphisms identified above can be further confirmed and their respective

5 frequencies can be determined through various methods using the previously described primers and probes. These methods can also be useful for genotyping either new populations in association studies or linkage analysis or individuals in the context of detection of alleles of biallelic markers which are known to be associated with a given trait. The genotyping of the biallelic markers is also important for the mapping. Those skilled in the art should note that the methods described below

10 can be equally performed on individual or pooled DNA samples.

Once a given polymorphic site has been found and characterized as a biallelic marker as described above, several methods can be used in order to determine the specific allele carried by an individual at the given polymorphic base.

The identification of biallelic markers described previously allows the design of appropriate oligonucleotides, which can be used as probes and primers, to amplify a *TBC-1* gene containing the polymorphic site of interest and for the detection of such polymorphisms.

The biallelic markers according to the present invention may be used in methods for the identification and characterization of an association between alleles for one or several biallelic markers of the sequence of the *TBC-1* gene and a trait.

The identified polymorphisms, and consequently the biallelic markers of the invention, may be used in methods for the detection in an individual of *TBC-1* alleles associated with a trait, more particularly a trait related to a cell differentiation or abnormal cell proliferation disorders, and most particularly a trait related to cancer diseases, specifically prostate cancer.

In one embodiment the invention encompasses methods of genotyping comprising

determining the identity of a nucleotide at a TBC-1-related biallelic marker or the complement thereof in a biological sample; optionally, wherein said TBC-1-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said biological sample is derived from a single subject; optionally, wherein the identity of the nucleotides at said biallelic marker is

determined for both copies of said biallelic marker present in said individual's genome; optionally, wherein said biological sample is derived from multiple subjects; Optionally, the genotyping methods of the invention encompass methods with any further limitation described in this disclosure, or those following, specified alone or in any combination; Optionally, said method is performed in vitro; optionally, further comprising amplifying a portion of said sequence

comprising the biallelic marker prior to said determining step; Optionally, wherein said amplifying is performed by PCR, LCR, or replication of a recombinant vector comprising an origin of replication and said fragment in a host cell; optionally, wherein said determining is performed by a

30

35

WO 00/08209 PCT/IB99/01444

hybridization assay, a sequencing assay, a microsequencing assay, or an enzyme-based mismatch detection assay.

Source of Nucleic Acids for genotyping

Any source of nucleic acids, in purified or non-purified form, can be utilized as the starting nucleic acid, provided it contains or is suspected of containing the specific nucleic acid sequence desired. DNA or RNA may be extracted from cells, tissues, body fluids and the like as described above. While nucleic acids for use in the genotyping methods of the invention can be derived from any mammalian source, the test subjects and individuals from which nucleic acid samples are taken are generally understood to be human.

10 Amplification Of DNA Fragments Comprising Biallelic Markers

Methods and polynucleotides are provided to amplify a segment of nucleotides comprising one or more biallelic marker of the present invention. It will be appreciated that amplification of DNA fragments comprising biallelic markers may be used in various methods and for various purposes and is not restricted to genotyping. Nevertheless, many genotyping methods, although not all, require the previous amplification of the DNA region carrying the biallelic marker of interest. Such methods specifically increase the concentration or total number of sequences that span the biallelic marker or include that site and sequences located either distal or proximal to it. Diagnostic assays may also rely on amplification of DNA segments carrying a biallelic marker of the present invention. Amplification of DNA may be achieved by any method known in the art. Amplification techniques are described above in the section entitled, "Identification of TBC-1-related biallelic markers."

Some of these amplification methods are particularly suited for the detection of single nucleotide polymorphisms and allow the simultaneous amplification of a target sequence and the identification of the polymorphic nucleotide as it is further described below.

The identification of biallelic markers as described above allows the design of appropriate oligonucleotides, which can be used as primers to amplify DNA fragments comprising the biallelic markers of the present invention. Amplification can be performed using the primers initially used to discover new biallelic markers which are described herein or any set of primers allowing the amplification of a DNA fragment comprising a biallelic marker of the present invention.

In some embodiments the present invention provides primers for amplifying a DNA fragment containing one or more biallelic markers of the present invention. Preferred amplification primers are listed in Example 2. It will be appreciated that the primers listed are merely exemplary and that any other set of primers which produce amplification products containing one or more biallelic markers of the present invention are also of use.

The spacing of the primers determines the length of the segment to be amplified. In the context of the present invention, amplified segments carrying biallelic markers can range in size from at least about 25 bp to 35 kbp. Amplification fragments from 25-3000 bp are typical,

fragments from 50-1000 bp are preferred and fragments from 100-600 bp are highly preferred. It will be appreciated that amplification primers for the biallelic markers may be any sequence which allow the specific amplification of any DNA fragment carrying the markers. Amplification primers may be labeled or immobilized on a solid support as described in "Oligonucleotide probes and 5 primers".

Methods of Genotyping DNA samples for Biallelic Markers

Any method known in the art can be used to identify the nucleotide present at a biallelic marker site. Since the biallelic marker allele to be detected has been identified and specified in the present invention, detection will prove simple for one of ordinary skill in the art by employing any of a number of techniques. Many genotyping methods require the previous amplification of the DNA region carrying the biallelic marker of interest. While the amplification of target or signal is often preferred at present, ultrasensitive detection methods which do not require amplification are also encompassed by the present genotyping methods. Methods well-known to those skilled in the art that can be used to detect biallelic polymorphisms include methods such as, conventional dot blot analyzes, single strand conformational polymorphism analysis (SSCP) described by Orita et al.(1989), denaturing gradient gel electrophoresis (DGGE), heteroduplex analysis, mismatch cleavage detection, and other conventional techniques as described in Sheffield et al.(1991), White et al.(1992), Grompe et al.(1989 and 1993). Another method for determining the identity of the nucleotide present at a particular polymorphic site employs a specialized exonuclease-resistant nucleotide derivative as described in US patent 4,656,127.

Preferred methods involve directly determining the identity of the nucleotide present at a biallelic marker site by sequencing assay, enzyme-based mismatch detection assay, or hybridization assay. The following is a description of some preferred methods. A highly preferred method is the microsequencing technique. The term "sequencing" is generally used herein to refer to polymerase extension of duplex primer/template complexes and includes both traditional sequencing and microsequencing.

1) Sequencing Assays

The nucleotide present at a polymorphic site can be determined by sequencing methods. In a preferred embodiment, DNA samples are subjected to PCR amplification before sequencing as described above. DNA sequencing methods are described in "Sequencing Of Amplified Genomic DNA And Identification Of Single Nucleotide Polymorphisms".

Preferably, the amplified DNA is subjected to automated dideoxy terminator sequencing reactions using a dye-primer cycle sequencing protocol. Sequence analysis allows the identification of the base present at the biallelic marker site.

In microsequencing methods, the nucleotide at a polymorphic site in a target DNA is detected by a single nucleotide primer extension reaction. This method involves appropriate microsequencing primers which, hybridize just upstream of the polymorphic base of interest in the target nucleic acid. A polymerase is used to specifically extend the 3' end of the primer with one single ddNTP (chain terminator) complementary to the nucleotide at the polymorphic site. Next the identity of the incorporated nucleotide is determined in any suitable way.

Typically, microsequencing reactions are carried out using fluorescent ddNTPs and the extended microsequencing primers are analyzed by electrophoresis on ABI 377 sequencing machines to determine the identity of the incorporated nucleotide as described in EP 412 883, the disclosure of which is incorporated herein by reference in its entirety. Alternatively capillary electrophoresis can be used in order to process a higher number of assays simultaneously. An example of a typical microsequencing procedure that can be used in the context of the present invention is provided in Example 4.

Different approaches can be used for the labeling and detection of ddNTPs. A

15 homogeneous phase detection method based on fluorescence resonance energy transfer has been described by Chen and Kwok (1997) and Chen et al.(1997). In this method, amplified genomic DNA fragments containing polymorphic sites are incubated with a 5'-fluorescein-labeled primer in the presence of allelic dye-labeled dideoxyribonucleoside triphosphates and a modified Taq polymerase. The dye-labeled primer is extended one base by the dye-terminator specific for the allele present on the template. At the end of the genotyping reaction, the fluorescence intensities of the two dyes in the reaction mixture are analyzed directly without separation or purification. All these steps can be performed in the same tube and the fluorescence changes can be monitored in real time. Alternatively, the extended primer may be analyzed by MALDI-TOF Mass Spectrometry. The base at the polymorphic site is identified by the mass added onto the microsequencing primer (see Haff and Smirnov, 1997).

Microsequencing may be achieved by the established microsequencing method or by developments or derivatives thereof. Alternative methods include several solid-phase microsequencing techniques. The basic microsequencing protocol is the same as described previously, except that the method is conducted as a heterogeneous phase assay, in which the primer or the target molecule is immobilized or captured onto a solid support. To simplify the primer separation and the terminal nucleotide addition analysis, oligonucleotides are attached to solid supports or are modified in such ways that permit affinity separation as well as polymerase extension. The 5' ends and internal nucleotides of synthetic oligonucleotides can be modified in a number of different ways to permit different affinity separation approaches, e.g., biotinylation. If a single affinity group is used on the oligonucleotides, the oligonucleotides can be separated from the incorporated terminator regent. This eliminates the need of physical or size separation. More than one oligonucleotide can be separated from the terminator reagent and analyzed simultaneously if

more than one affinity group is used. This permits the analysis of several nucleic acid species or more nucleic acid sequence information per extension reaction. The affinity group need not be on the priming oligonucleotide but could alternatively be present on the template. For example, immobilization can be carried out via an interaction between biotinylated DNA and streptavidin-5 coated microtitration wells or avidin-coated polystyrene particles. In the same manner, oligonucleotides or templates may be attached to a solid support in a high-density format. In such solid phase microsequencing reactions, incorporated ddNTPs can be radiolabeled (Syvänen, 1994) or linked to fluorescein (Livak and Hainer, 1994). The detection of radiolabeled ddNTPs can be achieved through scintillation-based techniques. The detection of fluorescein-linked ddNTPs can 10 be based on the binding of antifluorescein antibody conjugated with alkaline phosphatase, followed by incubation with a chromogenic substrate (such as p-nitrophenyl phosphate). Other possible reporter-detection pairs include: ddNTP linked to dinitrophenyl (DNP) and anti-DNP alkaline phosphatase conjugate (Harju et al., 1993) or biotinylated ddNTP and horseradish peroxidaseconjugated streptavidin with o-phenylenediamine as a substrate (WO 92/15712). As yet another 15 alternative solid-phase microsequencing procedure, Nyren et al.(1993) described a method relying on the detection of DNA polymerase activity by an enzymatic luminometric inorganic pyrophosphate detection assay (ELIDA).

Pastinen et al.(1997) describe a method for multiplex detection of single nucleotide polymorphism in which the solid phase minisequencing principle is applied to an oligonucleotide array format. High-density arrays of DNA probes attached to a solid support (DNA chips) are further described below.

In one aspect the present invention provides polynucleotides and methods to genotype one or more biallelic markers of the present invention by performing a microsequencing assay.

Preferred microsequencing primers include the nucleotide sequences D1 to D15 and E1 to E15. It will be appreciated that the microsequencing primers listed in Example 5 are merely exemplary and that, any primer having a 3' end immediately adjacent to the polymorphic nucleotide may be used. Similarly, it will be appreciated that microsequencing analysis may be performed for any biallelic marker or any combination of biallelic markers of the present invention. One aspect of the present invention is a solid support which includes one or more microsequencing primers listed in Example 5, or fragments comprising at least 8, 12, 15, 20, 25, 30, 40, or 50 consecutive nucleotides thereof, to the extent that such lengths are consistent with the primer described, and having a 3' terminus immediately upstream of the corresponding biallelic marker, for determining the identity of a nucleotide at a biallelic marker site.

3) Mismatch detection assays based on polymerases and ligases

In one aspect the present invention provides polynucleotides and methods to determine the allele of one or more biallelic markers of the present invention in a biological sample, by mismatch

30

WO 00/08209 PCT/IB99/01444 41

detection assays based on polymerases and/or ligases. These assays are based on the specificity of polymerases and ligases. Polymerization reactions places particularly stringent requirements on correct base pairing of the 3' end of the amplification primer and the joining of two oligonucleotides hybridized to a target DNA sequence is quite sensitive to mismatches close to the 5 ligation site, especially at the 3' end. Methods, primers and various parameters to amplify DNA fragments comprising biallelic markers of the present invention are further described above in "Amplification Of DNA Fragments Comprising Biallelic Markers".

Allele Specific Amplification Primers

Discrimination between the two alleles of a biallelic marker can also be achieved by allele 10 specific amplification, a selective strategy, whereby one of the alleles is amplified without amplification of the other allele. For allele specific amplification, at least one member of the pair of primers is sufficiently complementary with a region of a TBC-1 gene comprising the polymorphic base of a biallelic marker of the present invention to hybridize therewith and to initiate the amplification. Such primers are able to discriminate between the two alleles of a biallelic marker.

This is accomplished by placing the polymorphic base at the 3' end of one of the amplification primers. Because the extension forms from the 3'end of the primer, a mismatch at or near this position has an inhibitory effect on amplification. Therefore, under appropriate amplification conditions, these primers only direct amplification on their complementary allele. Determining the precise location of the mismatch and the corresponding assay conditions are well within the ordinary skill in the art.

Ligation/Amplification Based Methods

The "Oligonucleotide Ligation Assay" (OLA) uses two oligonucleotides which are designed to be capable of hybridizing to abutting sequences of a single strand of a target molecules. One of the oligonucleotides is biotinylated, and the other is detectably labeled. If the precise 25 complementary sequence is found in a target molecule, the oligonucleotides will hybridize such that their termini abut, and create a ligation substrate that can be captured and detected. OLA is capable of detecting single nucleotide polymorphisms and may be advantageously combined with PCR as described by Nickerson et al. (1990). In this method, PCR is used to achieve the exponential amplification of target DNA, which is then detected using OLA.

Other amplification methods which are particularly suited for the detection of single nucleotide polymorphism include LCR (ligase chain reaction), Gap LCR (GLCR) which are described above in "DNA Amplification". LCR uses two pairs of probes to exponentially amplify a specific target. The sequences of each pair of oligonucleotides, is selected to permit the pair to hybridize to abutting sequences of the same strand of the target. Such hybridization forms a 35 substrate for a template-dependant ligase. In accordance with the present invention, LCR can be performed with oligonucleotides having the proximal and distal sequences of the same strand of a biallelic marker site. In one embodiment, either oligonucleotide will be designed to include the

WO 00/08209 PCT/IB99/01444 42

biallelic marker site. In such an embodiment, the reaction conditions are selected such that the oligonucleotides can be ligated together only if the target molecule either contains or lacks the specific nucleotide that is complementary to the biallelic marker on the oligonucleotide. In an alternative embodiment, the oligonucleotides will not include the biallelic marker, such that when 5 they hybridize to the target molecule, a "gap" is created as described in WO 90/01069. This gap is then "filled" with complementary dNTPs (as mediated by DNA polymerase), or by an additional pair of oligonucleotides. Thus at the end of each cycle, each single strand has a complement capable of serving as a target during the next cycle and exponential allele-specific amplification of the desired sequence is obtained.

Ligase/Polymerase-mediated Genetic Bit AnalysisTM is another method for determining the identity of a nucleotide at a preselected site in a nucleic acid molecule (WO 95/21271). This method involves the incorporation of a nucleoside triphosphate that is complementary to the nucleotide present at the preselected site onto the terminus of a primer molecule, and their subsequent ligation to a second oligonucleotide. The reaction is monitored by detecting a specific 15 label attached to the reaction's solid phase or by detection in solution.

4) Hybridization Assay Methods

A preferred method of determining the identity of the nucleotide present at a biallelic marker site involves nucleic acid hybridization. The hybridization probes, which can be conveniently used in such reactions, preferably include the probes defined herein. Any 20 hybridization assay may be used including Southern hybridization, Northern hybridization, dot blot hybridization and solid-phase hybridization (see Sambrook et al., 1989).

Hybridization refers to the formation of a duplex structure by two single stranded nucleic acids due to complementary base pairing. Hybridization can occur between exactly complementary nucleic acid strands or between nucleic acid strands that contain minor regions of mismatch. 25 Specific probes can be designed that hybridize to one form of a biallelic marker and not to the other and therefore are able to discriminate between different allelic forms. Allele-specific probes are often used in pairs, one member of a pair showing perfect match to a target sequence containing the original allele and the other showing a perfect match to the target sequence containing the alternative allele. Hybridization conditions should be sufficiently stringent that there is a significant 30 difference in hybridization intensity between alleles, and preferably an essentially binary response, whereby a probe hybridizes to only one of the alleles. Stringent, sequence specific hybridization conditions, under which a probe will hybridize only to the exactly complementary target sequence are well known in the art (Sambrook et al., 1989). Stringent conditions are sequence dependent and will be different in different circumstances. Generally, stringent conditions are selected to be about 35 5°C lower than the thermal melting point (Tm) for the specific sequence at a defined ionic strength

and pH. Although such hybridization can be performed in solution, it is preferred to employ a

WO 00/08209 PCT/IB99/01444 43

solid-phase hybridization assay. The target DNA comprising a biallelic marker of the present invention may be amplified prior to the hybridization reaction. The presence of a specific allele in the sample is determined by detecting the presence or the absence of stable hybrid duplexes formed between the probe and the target DNA. The detection of hybrid duplexes can be carried out by a 5 number of methods. Various detection assay formats are well known which utilize detectable labels bound to either the target or the probe to enable detection of the hybrid duplexes. Typically, hybridization duplexes are separated from unhybridized nucleic acids and the labels bound to the duplexes are then detected. Those skilled in the art will recognize that wash steps may be employed to wash away excess target DNA or probe as well as unbound conjugate. Further, standard 10 heterogeneous assay formats are suitable for detecting the hybrids using the labels present on the primers and probes.

Two recently developed assays allow hybridization-based allele discrimination with no need for separations or washes (see Landegren U. et al., 1998). The TaqMan assay takes advantage of the 5' nuclease activity of Taq DNA polymerase to digest a DNA probe annealed specifically to 15 the accumulating amplification product. TaqMan probes are labeled with a donor-acceptor dye pair that interacts via fluorescence energy transfer. Cleavage of the TaqMan probe by the advancing polymerase during amplification dissociates the donor dye from the quenching acceptor dye, greatly increasing the donor fluorescence. All reagents necessary to detect two allelic variants can be assembled at the beginning of the reaction and the results are monitored in real time (see Livak et 20 al., 1995). In an alternative homogeneous hybridization based procedure, molecular beacons are used for allele discriminations. Molecular beacons are hairpin-shaped oligonucleotide probes that report the presence of specific nucleic acids in homogeneous solutions. When they bind to their targets they undergo a conformational reorganization that restores the fluorescence of an internally quenched fluorophore (Tyagi'et al., 1998).

The polynucleotides provided herein can be used to produce probes which can be used in hybridization assays for the detection of biallelic marker alleles in biological samples. These probes are characterized in that they preferably comprise between 8 and 50 nucleotides, and in that they are sufficiently complementary to a sequence comprising a biallelic marker of the present invention to hybridize thereto and preferably sufficiently specific to be able to discriminate the targeted 30 sequence for only one nucleotide variation. A particularly preferred probe is 25 nucleotides in length. Preferably the biallelic marker is within 4 nucleotides of the center of the polynucleotide probe. In particularly preferred probes, the biallelic marker is at the center of said polynucleotide. Preferred probes comprise a nucleotide sequence selected from the group consisting of amplicons listed in Table 1 and the sequences complementary thereto, or a fragment thereof, said fragment 35 comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. Preferred probes comprise a nucleotide sequence selected from the group consisting of P1 to P7, P9 to P13, P15 to

WO 00/08209

10

P19 and the sequences complementary thereto. In preferred embodiments the polymorphic base(s) are within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide.

Preferably the probes of the present invention are labeled or immobilized on a solid support.

Labels and solid supports are further described in "Oligonucleotide Probes and Primers". The probes can be non-extendable as described in "Oligonucleotide Probes and Primers".

By assaying the hybridization to an allele specific probe, one can detect the presence or absence of a biallelic marker allele in a given sample. High-Throughput parallel hybridization in array format is specifically encompassed within "hybridization assays" and are described below.

5) Hybridization To Addressable Arrays Of Oligonucleotides

Hybridization assays based on oligonucleotide arrays rely on the differences in hybridization stability of short oligonucleotides to perfectly matched and mismatched target sequence variants. Efficient access to polymorphism information is obtained through a basic structure comprising high-density arrays of oligonucleotide probes attached to a solid support (e.g., the chip) at selected positions. Each DNA chip can contain thousands to millions of individual synthetic DNA probes arranged in a grid-like pattern and miniaturized to the size of a dime.

The chip technology has already been applied with success in numerous cases. For example, the screening of mutations has been undertaken in the BRCA1 gene, in *S. cerevisiae* mutant strains, and in the protease gene of HIV-1 virus (Hacia et al., 1996; Shoemaker et al., 1996; Chips of various formats for use in detecting biallelic polymorphisms can be produced on a customized basis by Affymetrix (GeneChip™), Hyseq (HyChip and HyGnostics), and Protogene Laboratories.

In general, thèse methods employ arrays of oligonucleotide probes that are complementary to target nucleic acid sequence segments from an individual which, target sequences include a polymorphic marker. EP 785280 describes a tiling strategy for the detection of single nucleotide polymorphisms. Briefly, arrays may generally be "tiled" for a large number of specific polymorphisms. By "tiling" is generally meant the synthesis of a defined set of oligonucleotide probes which is made up of a sequence complementary to the target sequence of interest, as well as preselected variations of that sequence, e.g., substitution of one or more given positions with one or more members of the basis set of nucleotides. Tiling strategies are further described in PCT application No. WO 95/11995. In a particular aspect, arrays are tiled for a number of specific, identified biallelic marker sequences. In particular, the array is tiled to include a number of detection blocks, each detection block being specific for a specific biallelic marker or a set of biallelic markers. For example, a detection block may be tiled to include a number of probes, which span the sequence segment that includes a specific polymorphism. To ensure probes that are complementary to each allele, the probes are synthesized in pairs differing at the biallelic marker.

<u>1.1</u>

20

30

WO 00/08209 PCT/IB99/01444

In addition to the probes differing at the polymorphic base, monosubstituted probes are also generally tiled within the detection block. These monosubstituted probes have bases at and up to a certain number of bases in either direction from the polymorphism, substituted with the remaining nucleotides (selected from A, T, G, C and U). Typically the probes in a tiled detection block will include substitutions of the sequence positions up to and including those that are 5 bases away from the biallelic marker. The monosubstituted probes provide internal controls for the tiled array, to distinguish actual hybridization from artefactual cross-hybridization. Upon completion of hybridization with the target sequence and washing of the array, the array is scanned to determine the position on the array to which the target sequence hybridizes. The hybridization data from the scanned array is then analyzed to identify which allele or alleles of the biallelic marker are present in the sample. Hybridization and scanning may be carried out as described in PCT application No. WO 92/10092 and WO 95/11995 and US patent No. 5,424,186.

Thus, in some embodiments, the chips may comprise an array of nucleic acid sequences of fragments of about 15 nucleotides in length. In further embodiments, the chip may comprise an array including at least one of the sequences selected from the group consisting of amplicons listed in table 1 and the sequences complementary thereto, or a fragment thereof, said fragment comprising at least about 8 consecutive nucleotides, preferably 10, 15, 20, more preferably 25, 30, 40, 47, or 50 consecutive nucleotides and containing a polymorphic base. In preferred embodiments the polymorphic base is within 5, 4, 3, 2, 1, nucleotides of the center of the said polynucleotide, more preferably at the center of said polynucleotide. In some embodiments, the chip may comprise an array of at least 2, 3, 4, 5, 6, 7, 8 or more of these polynucleotides of the invention. Solid supports and polynucleotides of the present invention attached to solid supports are further described in "Oligonucleotide Probes And Primers".

6) Integrated Systems

Another technique, which may be used to analyze polymorphisms, includes multicomponent integrated systems, which miniaturize and compartmentalize processes such as PCR and capillary electrophoresis reactions in a single functional device. An example of such technique is disclosed in US patent 5,589,136, which describes the integration of PCR amplification and capillary electrophoresis in chips.

Integrated systems can be envisaged mainly when microfluidic systems are used. These systems comprise a pattern of microchannels designed onto a glass, silicon, quartz, or plastic wafer included on a microchip. The movements of the samples are controlled by electric, electroosmotic or hydrostatic forces applied across different areas of the microchip to create functional microscopic valves and pumps with no moving parts.

WO 00/08209 PCT/IB99/01444 46

For genotyping biallelic markers, the microfluidic system may integrate nucleic acid amplification, microsequencing, capillary electrophoresis and a detection method such as laserinduced fluorescence detection.

Association Studies With The Biallelic Markers Of The TBC-1 Gene

The identification of genes involved in suspected heterogeneous, polygenic and multifactorial traits such as cancer can be carried out through two main strategies currently used for genetic mapping: linkage analysis and association studies. Association studies examine the frequency of marker alleles in unrelated trait positive (T+) individuals compared with trait negative (T-) controls, and are generally employed in the detection of polygenic inheritance. Association 10 studies as a method of mapping genetic traits rely on the phenomenon of linkage disequilibrium.

If two genetic loci lie on the same chromosome, then sets of alleles of these loci on the same chromosomal segment (called haplotypes) tend to be transmitted as a block from generation to generation. When not broken up by recombination, haplotypes can be tracked not only through pedigrees but also through populations. The resulting phenomenon at the population level is that the 15 occurrence of pairs of specific alleles at different loci on the same chromosome is not random, and the deviation from random is called linkage disequilibrium (LD).

If a specific allele in a given gene is directly involved in causing a particular trait T, its frequency will be statistically increased in a trait positive population when compared to the frequency in a trait negative population. As a consequence of the existence of linkage disequilibrium, the frequency of all other alleles present in the haplotype carrying the trait-causing allele (TCA) will also be increased in trait positive individuals compared to trait negative individuals. Therefore, association between the trait and any allele in linkage disequilibrium with the trait-causing allele will suffice to suggest the presence of a trait-related gene in that particular allele's region. Linkage disequilibrium allows the relative frequencies in trait positive and trait 25 negative populations of a limited number of genetic polymorphisms (specifically biallelic markers) to be analyzed as an alternative to screening all possible functional polymorphisms in order to find trait-causing alleles.

The general strategy to perform association studies using biallelic markers derived from a candidate region is to scan two groups of individuals (trait positive and trait negative control 30 individuals which are characterized by a well defined phenotype as described below) in order to measure and statistically compare the allele frequencies of such biallelic markers in both groups.

If a statistically significant association with a trait is identified for at least one or more of the analyzed biallelic markers, one can assume that : either the associated allele is directly responsible for causing the trait (associated allele is the trait-causing allele), or the associated allele 35 is in linkage disequilibrium with the trait-causing allele. If the evidence indicates that the associated allele within the candidate region is most probably not the trait-causing allele but is in linkage

disequilibrium with the real trait-causing allele, then the trait-causing allele, and by consequence the gene carrying the trait-causing allele, can be found by sequencing the vicinity of the associated marker.

Collection of DNA samples from trait positive (trait +) and trait negative (trait -individuals 5 (inclusion criteria)

In order to perform efficient and significant association studies such as those described herein, the trait under study should preferably follow a bimodal distribution in the population under study, presenting two clear non-overlapping phenotypes, trait positive and trait negative.

Nevertheless, even in the absence of such a bimodal distribution (as may in fact be the case for more complex genetic traits), any genetic trait may still be analyzed by the association method proposed here by carefully selecting the individuals to be included in the trait positive and trait negative phenotypic groups. The selection procedure involves to select individuals at opposite ends of the non-bimodal phenotype spectra of the trait under study, so as to include in these trait positive and trait negative populations individuals which clearly represent extreme, preferably non-overlapping phenotypes.

The definition of the inclusion criteria for the trait positive and trait negative populations is an important aspect of the present invention. The selection of drastically different but relatively uniform phenotypes enables efficient comparisons in association studies and the possible detection of marked differences at the genetic level, provided that the sample sizes of the populations under study are significant enough.

Generally, trait positive and trait negative populations to be included in association studies such as proposed in the present invention consist of phenotypically homogenous populations of individuals each representing 100% of the corresponding trait if the trait distribution is bimodal.

A first group of between 50 and 300 trait positive individuals, preferably about 100 individuals, can be recruited according to clinical inclusion criteria.

In each case, a similar number of trait negative individuals, preferably more than 100 individuals, are included in such studies who are preferably both ethnically- and age-matched to the trait positive cases. They are checked for the absence of the clinical criteria defined above. Both trait positive and trait negative individuals should correspond to unrelated cases.

30 Genotyping of trait positive and trait negative individuals

Allelic frequencies of the biallelic markers in each of the above described population can be determined using one of the methods described above under the heading "Methods of Genotyping DNA samples for biallelic markers". Analyses are preferably performed on amplified fragments obtained by genomic PCR performed on the DNA samples from each individual in similar conditions as those described above for the generation of biallelic markers.

In a preferred embodiment, amplified DNA samples are subjected to automated microsequencing reactions using fluorescent ddNTPs (specific fluorescence for each ddNTP) and

20

WO 00/08209 PCT/IB99/01444

the appropriate microsequencing oligonucleotides which hybridize just upstream of the polymorphic base.

Genotyping is further described in Example 5.

Associations studies can be carried out by the skilled technician using the biallelic markers of the invention defined above, with different trait positive and trait negative populations. Suitable examples of association studies using biallelic markers of the *TBC-1* gene, including the biallelic markers A1 to A19, involve studies on the following populations:

- a trait positive population suffering from a cancer, preferably prostate cancer and a healthy unaffected population; or
- a trait positive population suffering from prostate cancer treated with agents acting against prostate cancer and suffering from side-effects resulting from this treatment and an trait negative population suffering from prostate cancer treated with same agents without any substantial sideeffects, or
- a trait positive population suffering from prostate cancer treated with agents acting against 15 prostate cancer showing a beneficial response and a trait negative population suffering from prostate cancer treated with same agents without any beneficial response, or
 - a trait positive population suffering from prostate cancer presenting highly aggressive prostate cancer tumors and a trait negative population suffering from prostate cancer with prostate cancer tumors devoid of aggressiveness.

It is another object of the present invention to provide a method for the identification and characterization of an association between an allele of one or more biallelic markers of a TBC-1 gene and a trait. The method comprises the steps of:

- genotyping a marker or a group of biallelic markers according to the invention in trait positive;
- genotyping a marker or a group of biallelic markers according to the invention in and trait negative individuals; and
 - establishing a statistically significant association between one allele of at least one marker and the trait.

Preferably, the trait positive and trait negative individuals are selected from nonoverlapping phenotypes as regards to the trait under study. In one embodiment, the biallelic marker are selected from the group consisting of the biallelic markers A1 to A19.

In a preferred embodiment, the trait is cancer, prostate cancer, an early onset of prostate cancer, a susceptibility to prostate cancer, the level of aggressiveness of prostate cancer tumors, a modified expression of the *TBC-1* gene, a modified production of the *TBC-1* protein, or the production of a modified TBC-1 protein.

In a further embodiment, the trait negative population can be replaced in the association studies by a random control population.

15

The step of testing for and detecting the presence of DNA comprising specific alleles of a biallelic marker or a group of biallelic markers of the present invention can be carried out as described further below.

Oligonucleotide Probes And Primers

The invention relates also to oligonucleotide molecules useful as probes or primers, wherein said oligonucleotide molecules hybridize specifically with a nucleotide sequence comprised in the TBC-1 gene, particularly the TBC-1 genomic sequence of SEQ ID Nos 1 and 2 or the TBC-1 cDNAs sequences of SEQ ID Nos 3 and 4. More particularly, the present invention also concerns oligonucleotides for the detection of alleles of biallelic markers of the TBC-1 gene. These 10 oligonucleotides are useful either as primers for use in various processes such as DNA amplification and microsequencing or as probes for DNA recognition in hybridization analyses. Polynucleotides derived from the TBC-1 gene are useful in order to detect the presence of at least a copy of a nucleotide sequence of SEQ ID Nos 1-4, or a fragment, complement, or variant thereof in a test sample.

Particularly preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEO ID Nos 1 and 2, or the complements thereof. Additionally preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 1 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 1: 1-1000, 1001-2000, 2001-3000, 3001-4000, 4001-5000, 5001-6000, 6001-7000, 7001-8000, 8001 - 9000, 9001 - 10000, 10001 - 11000, 11001 - 12000, 12001 - 13000, 13001 - 14000, 14001 - 15000, 12001 - 12000, 12000 - 12000, 12000 - 120000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 120000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 12000, 12000 - 120000, 12000 - 12000, 12000 - 120000, 120000 - 120000, 120000 - 120000, 120000 - 1200000, 120000 - 1200000, 120000 - 12000000, 120025 15001-16000, 16001-17000, and 17001-17590. Other preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEO ID No 2 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 2: 1-5000, 5001-10000, 10001-15000, 15001- $30 \quad 20000, 20001 - 25000, 25001 - 30000, 30001 - 35000, 35001 - 40000, 40001 - 45000, 45001 - 50000.$ 50001-55000, 55001-60000, 60001-65000, 65001-70000, 70001-75000, 75001-80000, 80001-85000, 85001-90000, 90001-95000, and 95001-99960.

Moreover, preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 35 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID Nos 3 and 4, or the complements thereof.. Particularly preferred

WO 00/08209 PCT/IB99/01444

probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 3 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 3: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3983. Additional preferred probes and primers of the invention include isolated, purified, or recombinant polynucleotides comprising a contiguous span of at least 12, 15, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 500, or 1000 nucleotides of SEQ ID No 4 or the complements thereof, wherein said contiguous span comprises at least 1, 2, 3, 5, or 10 of the following nucleotide positions of SEQ ID No 4: 1-500, 501-1000, 1001-1500, 1501-2000, 2001-2500, 2501-3000, 3001-3500, and 3501-3988.

Thus, the invention also relates to nucleic acid probes characterized in that they hybridize specifically, under the stringent hybridization conditions defined above, with a nucleic acid selected from the group consisting of the nucleotide sequences of SEQ ID Nos 1-4 or a variant thereof or a sequence complementary thereto.

In one embodiment the invention encompasses isolated, purified, and recombinant polynucleotides consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of any one of SEQ ID Nos 1 and 2 and the complement thereof, wherein said span includes a *TBC-1*-related biallelic marker in said sequence; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith; optionally, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide; optionally, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said polynucleotide; optionally, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide; and optionally, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide. In a preferred embodiment, said probes comprises, consists of, or consists essentially of a sequence selected from the following sequences: P1 to P7, P9 to P13, P15 to P19 and the complementary sequences thereto.

In another embodiment the invention encompasses isolated, purified and recombinant polynucleotides comprising, consisting of, or consisting essentially of a contiguous span of 8 to 50 nucleotides of SEQ ID Nos 1 and 2, or the complements thereof, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide, and wherein the 3' end of said polynucleotide is located within 20 nucleotides upstream of a TBC-1-related biallelic marker in said sequence; optionally, wherein said TBC-1-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage

WO 00/08209 PCT/IB99/01444

disequilibrium therewith; optionally, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *TBC-1*-related biallelic marker in said sequence; and optionally, wherein said polynucleotide consists essentially of a sequence selected from the following sequences: D1 to D19 and E1 to E19.

In a further embodiment, the invention encompasses isolated, purified, or recombinant polynucleotides comprising, consisting of, or consisting essentially of a sequence selected from the following sequences: B1 to B15 and C1 to C15.

In an additional embodiment, the invention encompasses polynucleotides for use in hybridization assays, sequencing assays, and enzyme-based mismatch detection assays for determining the identity of the nucleotide at a *TBC-1*-related biallelic marker in SEQ ID Nos 1 and 2, or the complements thereof, as well as polynucleotides for use in amplifying segments of nucleotides comprising a *TBC-1*-related biallelic marker in SEQ ID Nos 1 and 2, or the complements thereof; optionally, wherein said *TBC-1*-related biallelic marker is selected from the group consisting of A1 to A19, and the complements thereof, or optionally the biallelic markers in linkage disequilibrium therewith.

A probe or a primer according to the invention has between 8 and 1000 nucleotides in length, or is specified to be at least 12, 15, 18, 20, 25, 35, 40, 50, 60, 70, 80, 100, 250, 500 or 1000 nucleotides in length. More particularly, the length of these probes and primers can range from 8, 10, 15, 20, or 30 to 100 nucleotides, preferably from 10 to 50, more preferably from 15 to 30 nucleotides. Shorter probes and primers tend to lack specificity for a target nucleic acid sequence and generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. Longer probes and primers are expensive to produce and can sometimes self-hybridize to form hairpin structures. The appropriate length for primers and probes under a particular set of assay conditions may be empirically determined by one of skill in the art. A preferred probe or primer consists of a nucleic acid comprising a polynucleotide selected from the group of the nucleotide sequences of P1 to P7, P9 to P13, P15 to P19 and the complementary sequence thereto, B1 to B15, C1 to C15, D1 to D19, E1 to E19, for which the respective locations in the sequence listing are provided in Tables 2, 3 and 4.

The formation of stable hybrids depends on the melting temperature (Tm) of the DNA. The 30 Tm depends on the length of the primer or probe, the ionic strength of the solution and the G+C content. The higher the G+C content of the primer or probe, the higher is the melting temperature because G:C pairs are held by three H bonds whereas A:T pairs have only two. The GC content in the probes of the invention usually ranges between 10 and 75 %, preferably between 35 and 60 %, and more preferably between 40 and 55 %.

The primers and probes can be prepared by any suitable method, including, for example, cloning and restriction of appropriate sequences and direct chemical synthesis by a method such as the phosphodiester method of Narang et al.(1979), the phosphodiester method of Brown et

WO 00/08209 PCT/IB99/01444 52

al.(1979), the diethylphosphoramidite method of Beaucage et al.(1981) and the solid support method described in EP 0 707 592.

Detection probes are generally nucleic acid sequences or uncharged nucleic acid analogs such as, for example peptide nucleic acids which are disclosed in International Patent Application 5 WO 92/20702, morpholino analogs which are described in U.S. Patents Numbered 5,185,444; 5,034,506 and 5,142,047. The probe may have to be rendered "non-extendable" in that additional dNTPs cannot be added to the probe. In and of themselves analogs usually are non-extendable and nucleic acid probes can be rendered non-extendable by modifying the 3' end of the probe such that the hydroxyl group is no longer capable of participating in elongation. For example, the 3' end of 10 the probe can be functionalized with the capture or detection label to thereby consume or otherwise block the hydroxyl group. Alternatively, the 3' hydroxyl group simply can be cleaved, replaced or modified, U.S. Patent Application Serial No. 07/049,061 filed April 19, 1993 describes modifications, which can be used to render a probe non-extendable.

Any of the polynucleotides of the present invention can be labeled, if desired, by incorporating any label known in the art to be detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include radioactive substances (including, 32P, 35S, 3H, 125I), fluorescent dyes (including, 5-bromodesoxyuridin, fluorescein, acetylaminofluorene, digoxigenin) or biotin. Preferably, polynucleotides are labeled at their 3' and 5' ends. Examples of non-radioactive labeling of nucleic acid fragments are described 20 in the French patent No. FR-7810975 or by Urdea et al (1988) or Sanchez-Pescador et al (1988). In addition, the probes according to the present invention may have structural characteristics such that they allow the signal amplification, such structural characteristics being, for example, branched DNA probes as those described by Urdea et al. in 1991 or in the European patent No. EP 0 225 807 (Chiron).

A label can also be used to capture the primer, so as to facilitate the immobilization of either the primer or a primer extension product, such as amplified DNA, on a solid support. A capture label is attached to the primers or probes and can be a specific binding member which forms a binding pair with the solid's phase reagent's specific binding member (e.g. biotin and streptavidin). Therefore depending upon the type of label carried by a polynucleotide or a probe, it 30 may be employed to capture or to detect the target DNA. Further, it will be understood that the polynucleotides, primers or probes provided herein, may, themselves, serve as the capture label. For example, in the case where a solid phase reagent's binding member is a nucleic acid sequence, it may be selected such that it binds a complementary portion of a primer or probe to thereby immobilize the primer or probe to the solid phase. In cases where a polynucleotide probe itself 35 serves as the binding member, those skilled in the art will recognize that the probe will contain a sequence or "tail" that is not complementary to the target. In the case where a polynucleotide

WO 00/08209 PCT/IB99/01444 53

primer itself serves as the capture label, at least a portion of the primer will be free to hybridize with a nucleic acid on a solid phase. DNA Labeling techniques are well known to the skilled technician.

The probes of the present invention are useful for a number of purposes. They can be notably used in Southern hybridization to genomic DNA. The probes can also be used to detect 5 PCR amplification products. They may also be used to detect mismatches in the *TBC-1* gene or mRNA using other techniques.

Any of the polynucleotides, primers and probes of the present invention can be conveniently immobilized on a solid support. Solid supports are known to those skilled in the art and include the walls of wells of a reaction tray, test tubes, polystyrene beads, magnetic beads, 10 nitrocellulose strips, membranes, microparticles such as latex particles, sheep (or other animal) red blood cells, duracytes and others. The solid support is not critical and can be selected by one skilled in the art. Thus, latex particles, microparticles, magnetic or non-magnetic beads, membranes, plastic tubes, walls of microtiter wells, glass or silicon chips, sheep (or other suitable animal's) red blood cells and duracytes are all suitable examples. Suitable methods for immobilizing nucleic acids on solid phases include ionic, hydrophobic, covalent interactions and the like. A solid support, as used herein, refers to any material which is insoluble, or can be made insoluble by a subsequent reaction. The solid support can be chosen for its intrinsic ability to attract and immobilize the capture reagent. Alternatively, the solid phase can retain an additional receptor which has the ability to attract and immobilize the capture reagent. The additional receptor can include a charged substance that is oppositely charged with respect to the capture reagent itself or to a charged substance conjugated to the capture reagent. As yet another alternative, the receptor molecule can be any specific binding member which is immobilized upon (attached to) the solid support and which has the ability to immobilize the capture reagent through a specific binding reaction. The receptor molecule enables the indirect binding of the capture reagent to a solid 25 support material before the performance of the assay or during the performance of the assay. The solid phase thus can be a plastic, derivatized plastic, magnetic or non-magnetic metal, glass or silicon surface of a test tube, microtiter well, sheet, bead, microparticle, chip, sheep (or other suitable animal's) red blood cells, duracytes® and other configurations known to those of ordinary skill in the art. The polynucleotides of the invention can be attached to or immobilized on a solid 30 support individually or in groups of at least 2, 5, 8, 10, 12, 15, 20, or 25 distinct polynucleotides of the invention to a single solid support. In addition, polynucleotides other than those of the invention may be attached to the same solid support as one or more polynucleotides of the invention.

Consequently, the invention also deals with a method for detecting the presence of a nucleic acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto in a sample, said method comprising the following steps of:

WO 00/08209 PCT/IB99/01444 54

a) bringing into contact a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a nucleotide sequence included in a nucleic acid selected form the group consisting of the nucleotide sequences of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto and the sample to be assayed; and

b) detecting the hybrid complex formed between the probe and a nucleic acid in the sample.

The invention further concerns a kit for detecting the presence of a nucleic acid comprising a nucleotide sequence selected from a group consisting of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto in a sample, said kit comprising:

- a) a nucleic acid probe or a plurality of nucleic acid probes which can hybridize with a
 nucleotide sequence included in a nucleic acid selected form the group consisting of the nucleotide sequences of SEQ ID Nos 1-4, a fragment or a variant thereof and a complementary sequence thereto; and
 - b) optionally, the reagents necessary for performing the hybridization reaction.

In a first preferred embodiment of this detection method and kit, said nucleic acid probe or the plurality of nucleic acid probes are labeled with a detectable molecule. In a second preferred embodiment of said method and kit, said nucleic acid probe or the plurality of nucleic acid probes has been immobilized on a substrate. In a third preferred embodiment, the nucleic acid probe or the plurality of nucleic acid probes comprise either a sequence which is selected from the group consisting of the nucleotide sequences of P1 to P7, P9 to P13, P15 to P19 and the complementary sequence thereto, B1 to B15, C1 to C15, D1 to D19, E1 to E19 or a biallelic marker selected from the group consisting of A1 to A19 and the complements thereto.

Oligonucleotide Arrays

A substrate comprising a plurality of oligonucleotide primers or probes of the invention may be used either for detecting or amplifying targeted sequences in the TBC-1 gene and may also be used for detecting mutations in the coding or in the non-coding sequences of the TBC-1 gene.

Any polynucleotide provided herein may be attached in overlapping areas or at random locations on the solid support. Alternatively the polynucleotides of the invention may be attached in an ordered array wherein each polynucleotide is attached to a distinct region of the solid support which does not overlap with the attachment site of any other polynucleotide. Preferably, such an ordered array of polynucleotides is designed to be "addressable" where the distinct locations are recorded and can be accessed as part of an assay procedure. Addressable polynucleotide arrays typically comprise a plurality of different oligonucleotide probes that are coupled to a surface of a substrate in different known locations. The knowledge of the precise location of each polynucleotides location makes these "addressable" arrays particularly useful in hybridization assays. Any addressable array technology known in the art can be employed with the polynucleotides of the invention. One particular embodiment of these polynucleotide arrays is known as the Genechips™, and has been generally described in US Patent 5,143,854; PCT

WO 00/08209 PCT/IB99/01444 55

publications WO 90/15070 and 92/10092. These arrays may generally be produced using mechanical synthesis methods or light directed synthesis methods which incorporate a combination of photolithographic methods and solid phase oligonucleotide synthesis (Fodor et al., 1991). The immobilization of arrays of oligonucleotides on solid supports has been rendered possible by the development of a technology generally identified as "Very Large Scale Immobilized Polymer Synthesis" (VLSIPS™) in which, typically, probes are immobilized in a high density array on a solid surface of a chip. Examples of VLSIPS™ technologies are provided in US Patents 5,143,854; and 5,412,087 and in PCT Publications WO 90/15070, WO 92/10092 and WO 95/11995, which describe methods for forming oligonucleotide arrays through techniques such as light-directed synthesis techniques. In designing strategies aimed at providing arrays of nucleotides immobilized on solid supports, further presentation strategies were developed to order and display the oligonucleotide arrays on the chips in an attempt to maximize hybridization patterns and sequence information. Examples of such presentation strategies are disclosed in PCT Publications WO 94/12305, WO 94/11530, WO 97/29212 and WO 97/31256.

In another embodiment of the oligonucleotide arrays of the invention, an oligonucleotide probe matrix may advantageously be used to detect mutations occurring in the *TBC-1* gene and preferably in its regulatory region. For this particular purpose, probes are specifically designed to have a nucleotide sequence allowing their hybridization to the genes that carry known mutations (either by deletion, insertion or substitution of one or several nucleotides). By known mutations, it is meant, mutations on the *TBC-1* gene that have been identified according, for example to the technique used by Huang et al.(1996) or Samson et al.(1996).

Another technique that is used to detect mutations in the *TBC-1* gene is the use of a high-density DNA array. Each oligonucleotide probe constituting a unit element of the high density DNA array is designed to match a specific subsequence of the *TBC-1* genomic DNA or cDNA.

25 Thus, an array consisting of oligonucleotides complementary to subsequences of the target gene sequence is used to determine the identity of the target sequence with the wild gene sequence, measure its amount, and detect differences between the target sequence and the reference wild gene sequence of the *TBC-1* gene. In one such design, termed 4L tiled array, is implemented a set of four probes (A, C, G, T), preferably 15-nucleotide oligomers. In each set of four probes, the perfect complement will hybridize more strongly than mismatched probes. Consequently, a nucleic acid target of length L is scanned for mutations with a tiled array containing 4L probes, the whole probe set containing all the possible mutations in the known wild reference sequence. The hybridization signals of the 15-mer probe set tiled array are perturbed by a single base change in the target sequence. As a consequence, there is a characteristic loss of signal or a "footprint" for the probes

55 flanking a mutation position. This technique was described by Chee et al. in 1996.

Consequently, the invention concerns an array of nucleic acid molecules comprising at least one polynucleotide described above as probes and primers. Preferably, the invention concerns an

30

WO 00/08209 PCT/IB99/01444 56

array of nucleic acid comprising at least two polynucleotides described above as probes and primers.

A further object of the invention consists of an array of nucleic acid sequences comprising either at least one of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19, B1 to B15, C1 to C15, D1 to D19, E1 to E19, the sequences complementary thereto, a fragment thereof of at least 8, 10, 12, 15, 18, 20, 25, 30, or 40 consecutive nucleotides thereof, and at least one sequence comprising a biallelic marker selected from the group consisting of A1 to A19 and the complements thereto.

The invention also pertains to an array of nucleic acid sequences comprising either at least two of the sequences selected from the group consisting of P1 to P7, P9 to P13, P15 to P19, B1 to B15, C1 to C15, D1 to D19, E1 to E19, the sequences complementary thereto, a fragment thereof of at least 8 consecutive nucleotides thereof, and at least two sequences comprising a biallelic marker selected from the group consisting of A1 to A19 and the complements thereof.

Vectors For The Expression Of A Regulatory Or A Coding Polynucleotide Of TBC-1.

Any of the regulatory polynucleotides or the coding polynucleotides of the invention may be inserted into recombinant vectors for expression in a recombinant host cell or a recombinant host organism.

Thus, the present invention also encompasses a family of recombinant vectors that contains either a regulatory polynucleotide selected from the group consisting of any one of the regulatory polynucleotides derived from the *TBC-1* genomic sequences of SEQ ID Nos 1 and 2, or a polynucleotide comprising the *TBC-1* coding sequence, or both.

In a first preferred embodiment, a recombinant vector of the invention is used as an expression vector: (a) the TBC-1 regulatory sequence comprised therein drives the expression of a coding polynucleotide operably linked thereto; (b) the TBC-1 coding sequence is operably linked to regulation sequences allowing its expression in a suitable cell host and/or host organism.

In a second preferred embodiment, a recombinant vector of the invention is used to amplify the inserted polynucleotide derived from the *TBC-1* genomic sequences of SEQ ID Nos 1 and 2 or *TBC-1* cDNAs in a suitable cell host, this polynucleotide being amplified at every time that the recombinant vector replicates.

More particularly, the present invention relates to expression vectors which include nucleic acids encoding a TBC-1 protein, preferably the TBC-1 protein of the amino acid sequence of SEQ ID No 5 described therein, under the control of a regulatory sequence selected among the TBC-1 regulatory polynucleotides, or alternatively under the control of an exogenous regulatory sequence.

A recombinant expression vector comprising a nucleic acid selected from the group

35 consisting of 5' and 3' regulatory regions, or biologically active fragments or variants thereof, is
also part of the present invention.

WO 00/08209 PCT/IB99/01444 57

The invention also encompasses a recombinant expression vector comprising:

- a) a nucleic acid comprising the 5' regulatory polynucleotide of the nucleotide sequence SEQ ID No 1, or a biologically active fragment or variant thereof;
- b) a polynucleotide encoding a polypeptide or a polynucleotide of interest operably linked with said nucleic acid.
 - c) optionally, a nucleic acid comprising a 3'-regulatory polynucleotide, preferably a 3'-regulatory polynucleotide of the invention, or a biologically active fragment or variant thereof.

The nucleic acid comprising the 5' regulatory polynucleotide or a biologically active fragment or variant thereof may also comprises the 5'-UTR sequence from any of the two cDNA of the invention or a biologically active fragment or variant thereof.

The invention also pertains to a recombinant expression vector useful for the expression of the *TBC-1* coding sequence, wherein said vector comprises a nucleic acid selected from the group consisting of SEQ ID Nos 3 and 4 or a nucleic acid having at least 95% nucleotide identity with a polynucleotide selected from the group consisting of the nucleotide sequences of SEQ ID Nos 3 and 4.

Another recombinant expression vector of the invention consists in a recombinant vector comprising a nucleic acid comprising the nucleotide sequence beginning at the nucleotide in position 176 and ending in position 3730 of the polynucleotide of SEQ ID No 4.

Generally, a recombinant vector of the invention may comprise any of the polynucleotides described herein, including regulatory sequences, and coding sequences, as well as any *TBC-1* primer or probe as defined above. More particularly, the recombinant vectors of the present invention can comprise any of the polynucleotides described in the "*TBC-1* cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of *TBC-1*" section and the "Oligonucleotide Probes And Primers" section.

Some of the elements which can be found in the vectors of the present invention are described in further detail in the following sections.

a) Vectors

25

A recombinant vector according to the invention comprises, but is not limited to, a YAC (Yeast Artificial Chromosome), a BAC (Bacterial Artificial Chromosome), a phage, a phagemid, a cosmid, a plasmid or even a linear DNA molecule which may consist of a chromosomal, non-chromosomal and synthetic DNA. Such a recombinant vector can comprise a transcriptional unit comprising an assembly of:

- (1) a genetic element or elements having a regulatory role in gene expression, for example promoters or enhancers. Enhancers are cis-acting elements of DNA, usually from about 10 to 300
 35 bp in length that act on the promoter to increase the transcription.
 - (2) a structural or coding sequence which is transcribed into mRNA and eventually translated into a polypeptide, and

(3) appropriate transcription initiation and termination sequences. Structural units intended for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where a recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal residue.
 5 This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

Generally, recombinant expression vectors will include origins of replication, selectable markers permitting transformation of the host cell, and a promoter derived from a highly expressed gene to direct transcription of a downstream structural sequence. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably a leader sequence capable of directing secretion of the translated protein into the periplasmic space or the extracellular medium.

The selectable marker genes for selection of transformed host cells are preferably dihydrofolate reductase or neomycin resistance for eukaryotic cell culture, TRP1 for *S. cerevisiae* or tetracycline, rifampicin or ampicillin resistance in *E. coli*, or levan saccharase for mycobacteria.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and a bacterial origin of replication derived from commercially available plasmids comprising genetic elements of pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia, Uppsala, Sweden), and GEM1 (Promega Biotec, Madison, WI, USA).

Large numbers of suitable vectors and promoters are known to those of skill in the art, and commercially available, such as bacterial vectors: pQE70, pQE60, pQE-9 (Qiagen), pbs, pD10, phagescript, psiX174, pbluescript SK, pbsks, pNH8A, pNH16A, pNH18A, pNH46A (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia); or eukaryotic vectors: pWLNEO, pSV2CAT, pOG44, pXT1, pSG (Stratagene); pSVK3, pBPV, pMSG, pSVL (Pharmacia); baculovirus transfer vector pVL1392/1393 (Pharmingen); pQE-30 (QIAexpress).

A suitable vector for the expression of the TBC-1 polypeptide of SEQ ID No 5 is a baculovirus vector that can be propagated in insect cells and in insect cell lines. A specific suitable host vector system is the pVL1392/1393 baculovirus transfer vector (Pharmingen) that is used to transfect the SF9 cell line (ATCC N°CRL 1711) which is derived from Spodoptera frugiperda.

Other suitable vectors for the expression of the TBC-1 polypeptide of SEQ ID No 5 in a baculovirus expression system include those described by Chai et al. (1993), Vlasak et al. (1983) and Lenhard et al. (1996).

Mammalian expression vectors will comprise an origin of replication, a suitable promoter
and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and
acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences.

DNA sequences derived from the SV40 viral genome, for example SV40 origin, early promoter,

25

30

WO 00/08209 PCT/IB99/01444 59

enhancer, splice and polyadenylation sites may be used to provide the required nontranscribed genetic elements.

b) Promoters

The suitable promoter regions used in the expression vectors according to the present invention are chosen taking into account the cell host in which the heterologous gene has to be expressed.

A suitable promoter may be heterologous with respect to the nucleic acid for which it controls the expression or alternatively can be endogenous to the native polynucleotide containing the coding sequence to be expressed. Additionally, the promoter is generally heterologous with respect to the recombinant vector sequences within which the construct promoter/coding sequence has been inserted.

Preferred bacterial promoters are the LacI, LacZ, the T3 or T7 bacteriophage RNA polymerase promoters, the polyhedrin promoter, or the p10 protein promoter from baculovirus (Kit Novagen) (Smith et al., 1983; O'Reilly et al., 1992), the lambda P_R promoter or also the trc promoter.

Promoter regions can be selected from any desired gene using, for example, CAT (chloramphenicol transferase) vectors and more preferably pKK232-8 and pCM7 vectors.

Particularly preferred bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, PL and trp.

Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40,

LTRs from retrovirus, and mouse metallothionein-L. Selection of a convenient vector and promoter is well within the level of ordinary skill in the art.

The choice of a promoter is well within the ability of a person skilled in the field of genetic egineering. For example, one may refer to the book of Sambrook et al. (1989) or also to the procedures described by Fuller et al. (1996).

The vector containing the appropriate DNA sequence as described above, more preferably a *TBC-1* gene regulatory polynucleotide, a polynucleotide encoding the TBC-1 polypeptide of SEQ ID No 5 or both of them, can be utilized to transform an appropriate host to allow the expression of the desired polypeptide or polynucleotide.

c) Other types of vectors

The *in vivo* expression of a TBC-1 polypeptide of SEQ ID No 5 may be useful in order to correct a genetic defect related to the expression of the native gene in a host organism or to the production of a biologically inactive TBC-1 protein.

Consequently, the present invention also deals with recombinant expression vectors mainly designed for the *in vivo* production of the TBC-1 polypeptide of SEQ ID No 5 by the introduction of the appropriate genetic material in the organism of the patient to be treated. This genetic material may be introduced *in vitro* in a cell that has been previously extracted from the organism, the

6 inju

35

5

10

WO 00/08209 PCT/IB99/01444

modified cell being subsequently reintroduced in the said organism, directly in vivo into the appropriate tissue.

By « vector » according to this specific embodiment of the invention is intended either a circular or a linear DNA molecule.

One specific embodiment for a method for delivering a protein or peptide to the interior of a cell of a vertebrate in vivo comprises the step of introducing a preparation comprising a physiologically acceptable carrier and a naked polynucleotide operatively coding for the polypeptide of interest into the interstitial space of a tissue comprising the cell, whereby the naked polynucleotide is taken up into the interior of the cell and has a physiological effect.

In a specific embodiment, the invention provides a composition for the in vivo production of the TBC-1 protein or polypeptide described herein. It comprises a naked polynucleotide operatively coding for this polypeptide, in solution in a physiologically acceptable carrier, and suitable for introduction into a tissue to cause cells of the tissue to express the said protein or polypeptide.

Compositions comprising a polynucleotide are described in PCT application N° WO 90/11092 (Vical Inc.) and also in PCT application N° WO 95/11307 (Institut Pasteur, INSERM, Université d'Ottawa) as well as in the articles of Tacson et al. (1996) and of Huygen et al. (1996).

The amount of vector to be injected to the desired host organism varies according to the site of injection. As an indicative dose, it will be injected between 0,1 and 100 µg of the vector in an animal body, preferably a mammal body, for example a mouse body.

In another embodiment of the vector according to the invention, it may be introduced in vitro in a host cell, preferably in a host cell previously harvested from the animal to be treated and more preferably a somatic cell such as a muscle cell. In a subsequent step, the cell that has been transformed with the vector coding for the desired TBC-1 polypeptide or the desired fragment thereof is reintroduced into the animal body in order to deliver the recombinant protein within the 25 body either locally or systemically.

In one specific embodiment, the vector is derived from an adenovirus. Preferred adenovirus vectors according to the invention are those described by Feldman and Steg (1996) or Ohno et al. (1994). Another preferred recombinant adenovirus according to this specific embodiment of the present invention is the human adenovirus type 2 or 5 (Ad 2 or Ad 5) or an adenovirus of animal 30 origin (French patent application N° FR-93.05954).

Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery systems of choice for the transfer of exogenous polynucleotides in vivo, particularly to mammals, including humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host

Particularly preferred retroviruses for the preparation or construction of retroviral in vitro or in vitro gene delivery vehicles of the present invention include retroviruses selected from the group consisting of Mink-Cell Focus Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis virus

and Rous Sarcoma virus. Particularly preferred Murine Leukemia Viruses include the 4070A and the 1504A viruses, Abelson (ATCC No VR-999), Friend (ATCC No VR-245), Gross (ATCC No VR-590), Rauscher (ATCC No VR-998) and Moloney Murine Leukemia Virus (ATCC No VR-190; PCT Application No WO 94/24298). Particularly preferred Rous Sarcoma Viruses include Bryan high titer (ATCC Nos VR-334, VR-657, VR-726, VR-659 and VR-728). Other preferred retroviral vectors are those described in Roth et al. (Roth J.A. et al., 1996), PCT Application No WO 93/25234, PCT Application No WO 94/06920, Roux et al., 1989, Julan et al., 1992 and Neda et al., 1991.

Yet another viral vector system that is contemplated by the invention consists in the adenoassociated virus (AAV). The adeno-associated virus is a naturally occurring defective virus that
requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient
replication and a productive life cycle (Muzyczka et al., 1992). It is also one of the few viruses that
may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration
(Flotte et al., 1992; Samulski et al., 1989; McLaughlin et al., 1989). One advantageous feature of

AAV derives from its reduced efficacy for transducing primary cells relative to transformed cells.

Other compositions containing a vector of the invention advantageously comprise an oligonucleotide fragment of a nucleic sequence selected from the group consisting of SEQ ID Nos 3 or 4 as an antisense tool that inhibits the expression of the corresponding *TBC-1* gene. Preferred methods using antisense polynucleotide according to the present invention are the procedures described by Sczakiel et al. (1995) or those described in PCT Application No WO 95/24223.

Host cells

Another object of the invention consists in host cell that have been transformed or transfected with one of the polynucleotides described therein, and more precisely a polynucleotide either comprising a *TBC-1* regulatory polynucleotide or the coding sequence of the TBC-1 polypeptide having the amino acid sequence of SEQ ID No 5. Are included host cells that are transformed (prokaryotic cells) or that are transfected (eukaryotic cells) with a recombinant vector such as one of those described above.

A recombinant host cell of the invention comprises any one of the polynucleotides or the recombinant vectors described therein. More particularly, the cell hosts of the present invention can comprise any of the polynucleotides described in "TBC-1 cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of TBC-1" section and the "Oligonucleotide Probes And Primers" section.

Another preferred recombinant cell host according to the present invention is characterized in that its genome or genetic background (including chromosome, plasmids) is modified by the nucleic acid coding for the TBC-1 polypeptide of SEQ ID No 5.

Preferred host cells used as recipients for the expression vectors of the invention are the following:

- a) Prokaryotic host cells: Escherichia coli strains (I.E. DH5-α strain) or Bacillus subtilis.
- b) Eukaryotic host cells: HeLa cells (ATCC N°CCL2; N°CCL2.1; N°CCL2.2), Cv 1 cells

 5 (ATCC N°CCL70), COS cells (ATCC N°CRL1650; N°CRL1651), Sf-9 cells (ATCC N°CRL1711).

The constructs in the host cells can be used in a conventional manner to produce the gene product encoded by the recombinant sequence.

Following transformation of a suitable host and growth of the host to an appropriate cell density, the selected promoter is induced by appropriate means, such as temperature shift or chemical induction, and cells are cultivated for an additional period.

Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Microbial cells employed in the expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing

15 agents. Such methods are well known by the skill artisan.

Transgenic animals

The terms "transgenic animals" or "host animals" are used herein to designate animals that have their genome genetically and artificially manipulated so as to include one of the nucleic acids according to the invention. Preferred animals are non-human mammals and include those belonging to a genus selected from Mus (e.g. mice), Rattus (e.g. rats) and Oryctogalus (e.g. rabbits) which have their genome artificially and genetically altered by the insertion of a nucleic acid according to the invention.

The transgenic animals of the invention all include within a plurality of their cells a cloned recombinant or synthetic DNA sequence, more specifically one of the purified or isolated nucleic acids comprising a TBC-1 coding sequence, a TBC-1 regulatory polynucleotide or a DNA sequence encoding an antisense polynucleotide such as described in the present specification.

More particularly, transgenic animals according to the invention contain in their somatic cells and/or in their germ line cells any of the polynucleotides described in "TBC-1 cDNA Sequences" section, the "Coding Regions" section, "Genomic sequence of TBC-1" section, the "Oligonucleotide Probes And Primers" section and the "Vectors for the expression of a regulatory or coding polynucleotide of TBC-1" section.

The transgenic animals of the invention thus contain specific sequences of exogenous genetic material such as the nucleotide sequences described above in detail.

In a first preferred embodiment, these transgenic animals may be good experimental models in order to study the diverse pathologies related to cell differentiation, in particular concerning the

20

WO 00/08209 PCT/IB99/01444 63

transgenic animals within the genome of which has been inserted one or several copies of a polynucleotide encoding a native TBC-1 protein, or alternatively a mutant TBC-1 protein.

In a second preferred embodiment, these transgenic animals may express a desired polypeptide of interest under the control of the regulatory polynucleotides of the TBC-1 gene, 5 leading to good yields in the synthesis of this protein of interest, and eventually a tissue specific expression of this protein of interest.

Since it is possible to produce transgenic animals of the invention using a variety of different sequences, a general description will be given of the production of transgenic animals by referring generally to exogenous genetic material. This general description can be adapted by those 10 skilled in the art in order to incorporate the DNA sequences into animals. For more details regarding the production of transgenic animals, and specifically transgenic mice, it may be referred to Sandou et al. (1994) and also to US Patents Nos 4,873,191, issued Oct.10, 1989, 5,968,766, issued Dec. 16, 1997 and 5,387,742, issued Feb. 28, 1995, these documents being herein incorporated by reference to disclose methods for producing transgenic mice.

Transgenic animals of the present invention are produced by the application of procedures which result in an animal with a genome that incorporates exogenous genetic material which is integrated into the genome. The procedure involves obtaining the genetic material, or a portion thereof, which encodes either a TBC-1 coding sequence, a TBC-1 regulatory polynucleotide or a DNA sequence encoding an antisense polynucleotide such as described in the present specification.

A recombinant polynucleotide of the invention is inserted into an embryonic or ES stem cell line. The insertion is made using electroporation. The cells subjected to electroporation are screened (e.g. Southern blot analysis) to find positive cells which have integrated the exogenous recombinant polynucleotide into their genome. An illustrative positive-negative selection procedure that may be used according to the invention is described by Mansour et al. (1988). Then, the positive cells are 25 isolated, cloned and injected into 3.5 days old blastocysts from mice. The blastocysts are then inserted into a female host animal and allowed to grow to term. The offsprings of the female host are tested to determine which animals are transgenic e.g. include the inserted exogenous DNA sequence and which are wild-type.

Screening Of Agents Interacting With TBC-1

30 In a further embodiment, the present invention also concerns a method for the screening of new agents, or candidate substances interacting with TBC-1. These new agents could be useful against cancer.

In a preferred embodiment, the invention relates to a method for the screening of candidate substances comprising the following steps:

35 - providing a cell line, an organ, or a mammal expressing a TBC-1 gene or a fragment thereof, preferably the regulatory region or the promoter region of the TBC-1 gene.

25

WO 00/08209 PCT/IB99/01444

- obtaining a candidate substance preferably a candidate substance capable of inhibiting the binding of a transcription factor to the *TBC-1* regulatory region,

- testing the ability of the candidate substance to decrease the symptoms of prostate cancer and/or to modulate the expression levels of *TBC-1*.

In some embodiments, the cell line, organ or mammal expresses a heterologous protein, the coding sequence of which is operably linked to the *TBC-1* regulatory or promoter sequence. In other embodiments, they express a *TBC-1* gene comprising alleles of one or more *TBC-1*-related biallelic markers.

A candidate substance is a substance which can interact with or modulate, by binding or other intramolecular interactions, expression, stability, and function of *TBC-1*. Such substances may be potentially interesting for patients who are not responsive to existing drugs or develop side effects to them. Screening may be effected using either *in vitro* methods or *in vivo* methods.

Such methods can be carried out in numerous ways such as on transformed cells which express the considered alleles of the *TBC-1* gene, on tumors induced by said transformed cells, for example in mice, or on a *TBC-1* protein encoded by the considered allelic variant of *TBC-1*.

Screening assays of the present invention generally involve determining the ability of a candidate substance to present a cytotoxic effect, to change the characteristics of transformed cells such as proliferative and invasive capacity, to affect the tumor growth, or to modify the expression level of TBC-1.

Typically, this method includes preparing transformed cells with different forms of *TBC-1* sequences containing particular alleles of one or more biallelic markers and/or trait causing mutations described above. This is followed by testing the cells expressing the TBC-1 with a candidate substance to determine the ability of the substance to present cytotoxic effect, to affect the characteristics of transformed cells, the tumor growth, or to modify the expression level of *TBC-1*.

Typical examples of such drug screening assays are provided below. It is to be understood that the parameters set forth in these examples can be modified by the skilled person without undue experimentation.

Methods for screening substances interacting with a TBC-1 polypeptide

A method for the screening of a candidate substance according to the invention comprises 30 the following steps:

a)providing a polypeptide comprising the amino acid sequence SEQ ID No 5, or a peptide fragment or a variant thereof;

- b) obtaining a candidate substance;
- c) bringing into contact said polypeptide with said candidate substance;
- d) detecting the complexes formed between said polypeptide and said candidate substance. For the purpose of the present invention, a ligand means a molecule, such as a protein, a peptide, an antibody or any synthetic chemical compound capable of binding to the TBC-1 protein

1.11

30

35

or one of its fragments or variants or to modulate the expression of the polynucleotide coding for TBC-1 or a fragment or variant thereof.

In the ligand screening method according to the present invention, a biological sample or a defined molecule to be tested as a putative ligand of the TBC-1 protein is brought into contact with a purified TBC-1 protein, for example a purified recombinant TBC-1 protein produced by a recombinant cell host as described hereinbefore, in order to form a complex between the TBC-1 protein and the putative ligand molecule to be tested.

A. Candidate ligands obtained form random peptide libraries

In a particular embodiment of the screening method, the putative ligand is the expression product of a DNA insert contained in a phage vector (Parmley and Smith, 1988). Specifically, random peptide phages libraries are used. The random DNA inserts encode peptides of 8 to 20 aminoacids in length (Oldenburg K.R. et al., 1992,.; Valadon P., et al., 1996; Lucas A.H., 1994; Westerink M.A.J., 1995; Castagnoli L. et al., 1991). According to this particular embodiment, the recombinant phages expressing a protein that binds to the immobilized TBC-1 protein are retained and the complex formed between the TBC-1 protein and the recombinant phage may be subsequently immunoprecipitated by a polyclonal or a monoclonal antibody directed against the TBC-1 protein.

Once the ligand library in recombinant phages has been constructed, the phage population is brought into contact with the immobilized TBC-1 protein. Then the preparation of complexes is washed in order to remove the non-specifically bound recombinant phages. The phages that bind specifically to the TBC-1 protein are then eluted by a buffer (acid pH) or immunoprecipitated by the anti-TBC-1monoclonal antibody produced by a hybridoma, and this phage population is subsequently amplified by an over-infection of bacteria (for example E. coli). The selection step may be repeated several times, preferably 2-4 times, in order to select the more specific recombinant phage clones. The last step consists in characterizing the peptide produced by the selected recombinant phage clones either by expression in infected bacteria and isolation, expressing the phage insert in another host-vector system, or sequencing the insert contained in the selected recombinant phages.

B. Candidate ligands obtained through a two-hybrid screening assay.

The yeast two-hybrid system is designed to study protein-protein interactions *in vivo* (Fields and Song, 1989), and relies upon the fusion of a bait protein to the DNA binding domain of the yeast Gal4 protein. This technique is also described in US Patent N° US 5,667,973 and US Patent N° 5,283,173 (Fields et al.) the technical teachings of both patents being herein incorporated by reference.

The general procedure of library screening by the two-hybrid assay may be performed as described by Harper et al. (Harper JW et al., 1993) or as described by Cho et al. (1998) or also Fromont-Racine et al. (1997).

WO 00/08209 PCT/IB99/01444

The bait protein or polypeptide consists of a TBC-1 polypeptide or a fragment or variant thereof.

More precisely, the nucleotide sequence encoding the TBC-1 polypeptide or a fragment or variant thereof is fused to a polynucleotide encoding the DNA binding domain of the GAL4 protein, the fused nucleotide sequence being inserted in a suitable expression vector, for example pAS2 or pM3.

Then, a human cDNA library is constructed in a specially designed vector, such that the human cDNA insert is fused to a nucleotide sequence in the vector that encodes the transcriptional domain of the GAL4 protein. Preferably, the vector used is the pACT vector. The polypeptides encoded by the nucleotide inserts of the human cDNA library are termed "pray" polypeptides.

A third vector contains a detectable marker gene, such as beta galactosidase gene or CAT gene that is placed under the control of a regulation sequence that is responsive to the binding of a complete Gal4 protein containing both the transcriptional activation domain and the DNA binding domain. For example, the vector pG5EC may be used.

Two different yeast strains are also used. As an illustrative but non limiting example the two different yeast strains may be the following:

- Y190, the phenotype of which is (MATa, Leu2-3, 112 ura3-12, trp1-901, his3-D200, ade2-101, gal4Dgal180D URA3 GAL-LacZ, LYS GAL-HIS3, cyh');
- Y187, the phenotype of which is (MATa gal4 gal80 his3 trp1-901 ade2-101 ura3-52 leu2-3, -112 URA3 GAL-lacZmet), which is the opposite mating type of Y190.

Briefly, 20 μg of pAS2/TBC-1 and 20 μg of pACT-cDNA library are co-transformed into yeast strain Y190. The transformants are selected for growth on minimal media lacking histidine, leucine and tryptophan, but containing the histidine synthesis inhibitor 3-AT (50 mM). Positive colonies are screened for beta galactosidase by filter lift assay. The double positive colonies (*His*⁺,

- 25 beta-gal⁺) are then grown on plates lacking histidine, leucine, but containing tryptophan and cycloheximide (10 mg/ml) to select for loss of pAS2/TBC-1 plasmids but retention of pACT-cDNA library plasmids. The resulting Y190 strains are mated with Y187 strains expressing TBC-1 or non-related control proteins; such as cyclophilin B, lamin, or SNF1, as Gal4 fusions as described by Harper et al. (1993) and by Bram et al. (1993), and screened for beta galactosidase by filter lift
 30 assay. Yeast clones that are beta gal- after mating with the control Gal4 fusions are considered false
 - assay. Yeast clones that are beta gal- after mating with the control Gal4 fusions are considered false positives.

In another embodiment of the two-hybrid method according to the invention, the interaction between TBC-1 or a fragment or variant thereof with cellular proteins may be assessed using the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech).). As described in the manual accompanying the Matchmaker Two Hybrid System 2 (Catalog No. K1604-1, Clontech), the disclosure of which is incorporated herein by reference, nucleic acids encoding the TBC-1 protein or a portion thereof, are inserted into an expression vector such that they are in frame with DNA encoding the DNA

20

WO 00/08209 PCT/IB99/01444

binding domain of the yeast transcriptional activator GAL4. A desired cDNA, preferably human cDNA, is inserted into a second expression vector such that they are in frame with DNA encoding the activation domain of GAL4. The two expression plasmids are transformed into the yeast cells and the yeast cells are plated on selection medium which selects for expression of selectable markers on each of the expression vectors as well as GAL4 dependent expression of the HIS3 gene. Transformants capable of growing on medium lacking histidine are screened for GAL4 dependent lacZ expression. Those cells which are positive in both the histidine selection and the lacZ assay are those in which an interaction between TBC-1 and the protein or peptide encoded by the initially selected cDNA insert has taken place.

10 Method for screening ligands that modulate the expression of the TBC-1 gene.

Another subject of the present invention is a method for screening molecules that modulate the expression of the TBC-1 protein. Such a screening method comprises the steps of:

a) cultivating a prokaryotic or an eukaryotic cell that has been transfected with a nucleotide sequence encoding the TBC-1 protein, operably linked to a TBC-1 5'-regulatory sequence;

b) bringing into contact the cultivated cell with a molecule to be tested;

c) quantifying the expression of the TBC-1 protein.

Using DNA recombination techniques well known by the one skill in the art, the TBC-1 protein encoding DNA sequence is inserted into an expression vector, downstream from a TBC-1 5'-regulatory sequence that contains a TBC-1 promoter sequence.

The quantification of the expression of the TBC-1 protein may be realized either at the mRNA level or at the protein level. In the latter case, polyclonal or monoclonal antibodies may be used to quantify the amounts of the TBC-1 protein that have been produced, for example in an ELISA or a RIA assay.

In a preferred embodiment, the quantification of the TBC-1 mRNAs is realized by a quantitative PCR amplification of the cDNAs obtained by a reverse transcription of the total mRNA of the cultivated TBC-1-transfected host cell, using a pair of primers specific for TBC-1.

Expression levels and patterns of *TBC-1* may be analyzed by solution hybridization with long probes as described in International Patent Application No. WO 97/05277, the entire contents of which are incorporated herein by reference. Briefly, the *TBC-1* cDNA or the *TBC-1* genomic DNA described above, or fragments thereof, is inserted at a cloning site immediately downstream of a bacteriophage (T3, T7 or SP6) RNA polymerase promoter to produce antisense RNA. Preferably, the *TBC-1* insert comprises at least 100 or more consecutive nucleotides of the genomic DNA sequence or the cDNA sequences, particularly those comprising one of the nuceotide sequences of SEQ ID Nos 3, 4 and 6-8 or those encoding a mutated TBC-1. The plasmid is linearized and transcribed in the presence of ribonucleotides comprising modified ribonucleotides (i.e. biotin-UTP and DIG-UTP). An excess of this doubly labeled RNA is hybridized in solution with mRNA isolated from cells or tissues of interest. The hybridizations are performed under standard stringent conditions (40-50°C for 16 hours in an 80%

WO 00/08209 PCT/IB99/01444

formamide, 0.4 M NaCl buffer, pH 7-8). The unhybridized probe is removed by digestion with ribonucleases specific for single-stranded RNA (i.e. RNases CL3, T1, Phy M, U2 or A). The presence of the biotin-UTP modification enables capture of the hybrid on a microtitration plate coated with streptavidin. The presence of the DIG modification enables the hybrid to be detected and quantified by 5 ELISA using an anti-DIG antibody coupled to alkaline phosphatase.

Quantitative analysis of *TBC-1* gene expression may also be performed using arrays. As used herein, the term array means a one dimensional, two dimensional, or multidimensional arrangement of a plurality of nucleic acids of sufficient length to permit specific detection of expression of mRNAs capable of hybridizing thereto. For example, the arrays may contain a plurality of nucleic acids derived from genes whose expression levels are to be assessed. The arrays may include the *TBC-1* genomic DNA, the *TBC-1* cDNA sequences or the sequences complementary thereto or fragments thereof, particularly those comprising at least one of the biallelic markers according the present invention. Preferably, the fragments are at least 15 nucleotides in length. In other embodiments, the fragments are at least 25 nucleotides in length. In some embodiments, the fragments are at least 100 nucleotides in length. In another preferred embodiment, the fragments are more than 100 nucleotides in length. In some embodiments may be more than 500 nucleotides in length.

For example, quantitative analysis of *TBC-1* gene expression may be performed with a complementary DNA microarray as described by Schena et al. (1995). Full length *TBC-1* cDNAs or fragments thereof are amplified by PCR and arrayed from a 96-well microtiter plate onto silylated microscope slides using high-speed robotics. Printed arrays are incubated in a humid chamber to allow rehydration of the array elements and rinsed, once in 0.2% SDS for 1 min, twice in water for 1 min and once for 5 min in sodium borohydride solution. The arrays are submerged in water for 2 min at 95°C, transferred into 0.2% SDS for 1 min, rinsed twice with water, air dried and stored in the dark at 25°C.

Cell or tissue mRNA is isolated or commercially obtained and probes are prepared by a single round of reverse transcription. Probes are hybridized to 1 cm² microarrays under a 14 x 14 mm glass coverslip for 6-12 hours at 60°C. Arrays are washed for 5 min at 25°C in low stringency wash buffer (1 x SSC/0.2% SDS), then for 10 min at room temperature in high stringency wash buffer (0.1 x SSC/0.2% SDS). Arrays are scanned in 0.1 x SSC using a fluorescence laser scanning device fitted with a custom filter set. Accurate differential expression measurements are obtained by taking the average of the ratios of two independent hybridizations.

Quantitative analysis of *TBC-1* gene expression may also be performed with full length 35 *TBC-1* cDNAs or fragments thereof in complementary DNA arrays as described by Pietu et al. (1996). The full length *TBC-1* cDNA or fragments thereof is PCR amplified and spotted on membranes. Then, mRNAs originating from various tissues or cells are labeled with radioactive

30

WO 00/08209 PCT/IB99/01444

nucleotides. After hybridization and washing in controlled conditions, the hybridized mRNAs are detected by phospho-imaging or autoradiography. Duplicate experiments are performed and a quantitative analysis of differentially expressed mRNAs is then performed.

Alternatively, expression analysis using the *TBC-1* genomic DNA, the *TBC-1* cDNAs, or fragments thereof can be done through high density nucleotide arrays or chips as described by Lockhart et al. (1996) and Sosnowsky et al. (1997). Oligonucleotides of 15-50 nucleotides from the sequences of the *TBC-1* genomic DNA, the *TBC-1* cDNA sequences particularly those comprising at least one of biallelic markers according the present invention, preferably at least one of SEQ ID No 7-8 or those comprising the trait causing mutation, or the sequences complementary thereto, are synthesized directly on the chip (Lockhart et al., supra) or synthesized and then addressed to the chip (Sosnowski et al., supra). Preferably, the oligonucleotides are about 20 nucleotides in length.

TBC-1 cDNA probes labeled with an appropriate compound, such as biotin, digoxigenin or fluorescent dye, are synthesized from the appropriate mRNA population and then randomly fragmented to an average size of 50 to 100 nucleotides. The said probes are then hybridized to the chip. After washing as described in Lockhart et al., supra and application of different electric fields (Sosnowsky et al., 1997)., the dyes or labeling compounds are detected and quantified. Duplicate hybridizations are performed. Comparative analysis of the intensity of the signal originating from cDNA probes on the same target oligonucleotide in different cDNA samples indicates a differential expression of TBC-1 mRNAs.

Thus, is also part of the present invention a method for screening of a candidate substance or molecule that modulates the expression of the *TBC-1* gene according to the invention, wherein this method comprises the following steps:

- a) providing a recombinant cell host containing a nucleic acid, wherein said nucleic acid comprises the 5' regulatory région sequence or a biologically active fragment or variant thereof, the
 5' regulatory region or its biologically active fragment or variant being operably linked to a polynucleotide encoding a detectable protein;
 - b) obtaining a candidate substance, and
 - c) determining the ability of the candidate substance to modulate the expression levels of the polynucleotide encoding the detectable protein.

In a preferred embodiment of the above screening method, the nucleic acid comprising the 5' regulatory region sequence or a biologically active fragment or variant thereof also includes a 5'UTR region of one of the *TBC-1* cDNAs of SEQ ID Nos 3 and 4, or one of their biologically active fragments or variants thereof.

A second method for the screening of a candidate substance or molecule that modulates the expression of the TBC-1 gene comprises the following steps:

a) providing a recombinant cell host containing a nucleic acid, wherein said nucleic acid comprises a 5'UTR sequence of one of the TBC-1 cDNAs of SEQ ID Nos 3 and 4, or one of their

WO 00/08209 PCT/IB99/01444

biologically active fragments or variants, the 5'UTR sequence or its biologically active fragment or variant being operably linked to a polynucleotide encoding a detectable protein;

- b) obtaining a candidate substance, and
- c) determining the ability of the candidate substance to modulate the expression levels of the polynucleotide encoding the detectable protein.

In a preferred embodiment of the screening method described above, the nucleic acid that comprises a nucleotide sequence selected from the group consisting of the 5'UTR sequence of one of the TBC-1 cDNAs of SEQ ID Nos 3 and 4 or one of their biologically active fragments or variants, includes a promoter sequence, wherein said promoter sequence can be either endogenous, or in contrast exogenous with respect to the TBC-1 5'UTR sequences defined therein.

Among the preferred polynucleotides encoding a detectable protein, there may be cited polynucleotides encoding beta galactosidase, green fluorescent protein (GFP) and chloramphenicol acetyl transferase (CAT).

For the design of suitable recombinant vectors useful for performing the screening methods described above, it will be referred to the section of the present specification wherein the preferred recombinant vectors of the invention are detailed.

Screening using transgenic animals

In vivo methods can utilize transgenic animals for drug screening. Nucleic acids including at least one of the biallelic polymorphisms of interest can be used to generate genetically modified non-human animals or to generate site specific gene modifications in cell lines. The term "transgenic" is intended to encompass genetically modified animals having a deletion or other knock-out of TBC-1 gene activity, having an exogenous TBC-1 gene that is stably transmitted in the host cells, or having an exogenous TBC-1 promoter operably linked to a reporter gene. Transgenic animals may be made through homologous recombination, where the TBC-1 locus is altered.

25 Alternatively, a nucleic acid construct is randomly integrated into the genome. Vectors for stable integration include for example plasmids, retroviruses and other animal viruses, and YACs. Of interest are transgenic mammals e.g. cows, pigs, goats, horses, and particularly rodents such as rats and mice. Transgenic animals allow to study both efficacy and toxicity of the candidate drug.

Methods for inhibiting the expression of a TBC-1 gene

30 Other therapeutic compositions according to the present invention comprise advantageously an oligonucleotide fragment of the nucleic sequence of *TBC-1* as an antisense tool that inhibits the expression of the corresponding *TBC-1* gene. Preferred methods using antisense polynucleotide according to the present invention are the procedures described by Sczakiel et al. (1995).

Preferably, the antisense tools are chosen among the polynucleotides (15-200 bp long) that

35 are complementary to the 5'end of the *TBC-1* mRNA. In another embodiment, a combination of
different antisense polynucleotides complementary to different parts of the desired targetted gene
are used.

WO 00/08209 PCT/IB99/01444

Preferred antisense polynucleotides according to the present invention are complementary to a sequence of the mRNAs of *TBC-1* that contains the translation initiation codon ATG.

The antisense nucleic acid molecules to be used in gene therapy may be either DNA or RNA sequences. They comprise a nucleotide sequence complementary to the targeted sequence of the PTCA-1 genomic DNA, the sequence of which can be determined using one of the detection methods of the present invention. The targeted DNA or RNA sequence preferably comprises at least one of the biallelic markers according to the present invention. The antisense nucleic acids should have a length and melting temperature sufficient to permit formation of an intracellular duplex having sufficient stability to inhibit the expression of the *TBC-1* mRNA in the duplex. Strategies for designing antisense nucleic acids suitable for use in gene therapy are disclosed in Green et al., (1986) and Izant and Weintraub, (1984), the disclosures of which are incorporated herein by reference.

In some strategies, antisense molecules are obtained by reversing the orientation of the *TBC-1* coding region with respect to a promoter so as to transcribe the opposite strand from that which is normally transcribed in the cell. The antisense molecules may be transcribed using in vitro transcription systems such as those which employ T7 or SP6 polymerase to generate the transcript. Another approach involves transcription of *TBC-1* antisense nucleic acids in vivo by operably linking DNA containing the antisense sequence to a promoter in a suitable expression vector.

Alternatively, suitable antisense strategies are those described by Rossi et al. (1991), in the International Applications Nos. WO 94/23026, WO 95/04141, WO 92/18522 and in the European Patent Application No. EP 0 572 287 A2

An alternative to the antisense technology that is used according to the present invention consists in using ribozymes that will bind to a target sequence via their complementary polynucleotide tail and that will cleave the corresponding RNA by hydrolyzing its target site

25 (namely « hammerhead ribozymes »). Briefly, the simplified cycle of a hammerhead ribozyme consists of (1) sequence specific binding to the target RNA via complementary antisense sequences; (2) site-specific hydrolysis of the cleavable motif of the target strand; and (3) release of cleavage products, which gives rise to another catalytic cycle. Indeed, the use of long-chain antisense polynucleotide (at least 30 bases long) or ribozymes with long antisense arms are advantageous. A preferred delivery system for antisense ribozyme is achieved by covalently linking these antisense ribozymes to lipophilic groups or to use liposomes as a convenient vector. Preferred antisense ribozymes according to the present invention are prepared as described by Sczakiel et al. (1995), the specific preparation procedures being referred to in said article being herein incorporated by reference.

35

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specification

WO 00/08209 PCT/IB99/01444

referenced in this application are hereby incorporated by reference into the present disclosure to more fully describe the sate of the art to which this invention pertains.

EXAMPLES

EXAMPLE 1:

5 Analysis of the first mRNA encoding a TBC-1 polypeptide synthesized by the cells.

TBC-1 cDNA was obtained as follows: 4μl of ethanol suspension containing 1 mg of human prostate total RNA (Clontech laboratories, Inc., Palo Alto, USA; Catalogue N. 64038-1) was centrifuged, and the resulting pellet was air dried for 30 minutes at room temperature.

First strand cDNA synthesis was performed using the AdvantageTM RT-for- PCR kit

(Clontech laboratories Inc., catalogue N. K1402-1). 1 μl of 20 mM solution of a specific oligo dT primer was added to 12.5 μl of RNA solution in water, heated at 74°C for 2.5 min and rapidly quenched in an ice bath. 10 μl of 5 x RT buffer (50 mM Tris-HCl, pH 8.3, 75 mM KCl, 3 mM MgCl₂), 2.5 μl of dNTP mix (10 mM each), 1.25 μl of human recombinant placental RNA inhibitor were mixed with 1 ml of MMLV reverse transcriptase (200 units). 6.5 μl of this solution were added to RNA-primer mix and incubated at 42°C for one hour. 80 μl of water were added and the solution was incubated at 94°C for 5 minutes.

5μl of the resulting solution were used in a Long Range PCR reaction with hot start, in 50 μl final volume, using 2 units of rtTHXL, 20 pmol/μl of each of 5'-

TGACCACCATGCCCATGCT-3' (271-289 in SEQ ID No 3) and 5'-

20 GCATTTATTCACGTCCACGCC-3' (3929-3949 in SEQ ID No 3) primers with 35 cycles of elongation for 6 minutes at 67°C in thermocycler.

The amplification products corresponding to both cDNA strands were partially sequenced in order to ensure the specificity of the amplification reaction.

Results of Nothern blot analysis of prostate mRNAs supported the existence of the first 25 TBC-1 cDNA having about 4 kb in length, which is the nucleotide sequence of SEQ ID No 3.

Example 2:

Detection of TBC-1 biallelic markers: DNA extraction

Donors were unrelated and healthy. They presented a sufficient diversity for being representative of a French heterogeneous population. The DNA from 100 individuals was extracted and tested for the detection of the biallelic markers.

30 ml of peripheral venous blood were taken from each donor in the presence of EDTA. Cells (pellet) were collected after centrifugation for 10 minutes at 2000 rpm. Red cells were lysed by a lysis solution (50 ml final volume : 10 mM Tris pH7.6; 5 mM MgCl₂; 10 mM NaCl). The

10

15

solution was centrifuged (10 minutes, 2000 rpm) as many times as necessary to eliminate the residual red cells present in the supernatant, after resuspension of the pellet in the lysis solution.

The pellet of white cells was lysed overnight at 42°C with 3.7 ml of lysis solution composed of:

- 3 ml TE 10-2 (Tris-HCl 10 mM, EDTA 2 mM) / NaCl 0.4 M
- 200 µl SDS 10%
- 500 µl K-proteinase (2 mg K-proteinase in TE 10-2 / NaCl 0.4 M).

For the extraction of proteins, 1 ml saturated NaCl (6M) (1/3.5 v/v) was added. After vigorous agitation, the solution was centrifuged for 20 minutes at 10000 rpm.

For the precipitation of DNA, 2 to 3 volumes of 100% ethanol were added to the previous supernatant, and the solution was centrifuged for 30 minutes at 2000 rpm. The DNA solution was rinsed three times with 70% ethanol to eliminate salts, and centrifuged for 20 minutes at 2000 rpm. The pellet was dried at 37°C, and resuspended in 1 ml TE 10-1 or 1 ml water. The DNA concentration was evaluated by measuring the OD at 260 nm (1 unit OD = $50 \mu g/ml$ DNA).

To determine the presence of proteins in the DNA solution, the OD 260 / OD 280 ratio was determined. Only DNA preparations having a OD 260 / OD 280 ratio between 1.8 and 2 were used in the subsequent examples described below.

The pool was constituted by mixing equivalent quantities of DNA from each individual.

Example 3:

20 Detection of the biallelic markers: amplification of genomic DNA by PCR

The amplification of specific genomic sequences of the DNA samples of example 2 was carried out on the pool of DNA obtained previously. In addition, 50 individual samples were similarly amplified.

PCR assays were performed using the following protocol:

25	Final volume	25 μΙ
	DNA	2 ng/μl
	$MgCl_2$	2 mM
	dNTP (each)	200 μΜ
	primer (each)	2.9 ng/µl
30	Ampli Taq Gold DNA polymerase	0.05 unit/µl
	PCR buffer (10x = 0.1 M TrisHCl pH8.3 0.5M KCl	1x

Each pair of first primers was designed using the sequence information of the *TBC-1* gene disclosed herein and the OSP software (Hillier & Green, 1991). This first pair of primers was about 20 nucleotides in length and had the sequences disclosed in Table 1 in the columns labeled PU and 35 RP.

15

Table 1

Amplicon	Posi range ampli SEQ	of the con in	Primer name	Position range of amplification primer in SEQ ID No 1		Primer name	Complementary position range of amplification primer in SEQ ID No 1	
99-430	9391	9845	B1	9391	9408	C1	9828	9845
Amplicon	Posi range ampli SEQ	of the con in	Primer name			Primer name	position amplificat	mentary range of ion primer ID No 2
99-20508	988	1529	B2	988	1006	C2	1509	1529
99-20469	5039	5554	В3	5039	5056	C3	5534	5554
5-254	5997	6350	B4	5997	6015	C4	6332	6350
5-257	14371	14817	B5	14371	14390	C5	14798	14817
99-20511	18751	19217	B6	18751	18771	C6	19198	19217
99-20510	19605	20005	B7	19605	19625	C7	19986	20005
99-20504	29529	30061	B8	29529	29547	C8	30041	30061
99-20493	42268	42752	В9	42268	42287	C9	42732	42752
99-20499	69026	69543	B10	69026	69046	C10	69525	69543
99-20473	76323	76790	B11	76323	76343	C11	76771	· 76790
5-249	78292	78721	B12	78292	78309	C12	78704	78721
99-20485	81893	82372	B13	81893	81912	C13	82353	82372
99-20481	84392	84929	B14	84392	84412	C14	84909	84929
99-20480	89746	90198	B15	89746	89765	C15	90179	90198

Preferably, the primers contained a common oligonucleotide tail upstream of the specific bases targeted for amplification which was useful for sequencing.

Primers PU contain the following additional PU 5' sequence:
TGTAAAACGACGGCCAGT (SEQ ID No 6); primers RP contain the following RP 5' sequence:
CAGGAAACAGCTATGACC (SEQ ID No 7).

The synthesis of these primers was performed following the phosphoramidite method, on a GENSET UFPS 24.1 synthesizer.

DNA amplification was performed on a Genius II thermocycler. After heating at 95°C for 10 min, 40 cycles were performed. Each cycle comprised: 30 sec at 95°C, 54°C for 1 min, and 30 sec at 72°C. For final elongation, 10 min at 72°C ended the amplification. The quantities of the amplification products obtained were determined on 96-well microtiter plates, using a fluorometer and Picogreen as intercalant agent (Molecular Probes).

Example 4:

Detection of the biallelic markers: sequencing of amplified genomic DNA and identification of polymorphisms.

The sequencing of the amplified DNA obtained in example 3 was carried out on ABI 377 sequencers. The sequences of the amplification products were determined using automated dideoxy terminator sequencing reactions with a dye terminator cycle sequencing protocol. The products of

the sequencing reactions were run on sequencing gels and the sequences were determined using gel image analysis [ABI Prism DNA Sequencing Analysis software (2.1.2 version)].

The sequence data were further evaluated to detect the presence of biallelic markers among the pooled amplified fragments. The polymorphism search was based on the presence of superimposed peaks in the electrophoresis pattern resulting from different bases occurring at the same position as described previously.

15 fragments of amplification was analyzed. In this segment, 19 biallelic markers were detected. The localization of the biallelic marker is as shown in Table 2.

Table 2

Amplicon	BM	Marker	Marker Localization		rphism	BM position in
		Name	in TBC-1 gene	Allele 1	allele 2	SEQ ID No 1
99-430	A1	99-430-352	Intron 1	Α	G	9494
Amplicon	BM	Marker	Localization	Polymo	rphism	BM position in
		Name	in <i>TBC-1</i> gene	allele 1	allele 2	SEQ ID No 2
99-20508	A2	99-20508-456	Intron	С	T	1443
			upstream to			
			Exon A			
99-20469	A3	99-20469-213	Intron A	C	T	5247
5-254	A4	5-254-227	Intron B	A	G	6223
5-257	A.5	5-257-353	Intron D	С	T	14723
99-20511	A6	99-20511-32	Intron D	С	T	19186
99-20511	A 7	99-20511-221	Intron D	Α	G	18997
99-20510	A8	99-20510-115	Intron D	deletion of		19891
				TCT		l
99-20504	A9	99-20504-90	Intron D	Α	G	29617
99-20493	A10	99-20493-238	Intron D	Α	C	42519
99-20499	A11	99-20499-221	Intron G	Α	G	69324
99-20499	A12	99-20499-364	Intron G	A	T	69181
99-20499	A13	99-20499-399	Intron G	Α	G	69146
99-20473	A14	99-20473-138	Intron H	deletion of	-	76458
				TAACA		
5-249	A15	5-249-304	Intron I	Α	G	78595
99-20485	A16	99-20485-269	Intron I	Α	G	82159
99-20481	A17	99-20481-131	Intron I	G	С	84522
99-20481	A18	99-20481-419	Intron I	Α	T	84810
99-20480	A19	99-20480-233	Intron J	A	G	89967

BM refers to "biallelic marker". All 1 and all 2 refer respectively to allele 1 and allele 2 of the biallelic marker.

Table 3

BM	Marker Name		n range bes in	Probes
1		SEQ I	D No 1	
A1	99-430-352	9482	9506	P1

BM	Marker Name	Positio	Position range	
		of pro	bes in	
		SEQ I	D No 2	
A2	99-20508-456	1431	1455	P2
A 3	99-20469-213	5235	5259	P3
A4	5-254-227	6211	6235	P4
A5	5-257-353	14711	14735	P5
A6	99-20511-32	19174	19198	P6
A 7	99-20511-221	18985	19009	P7
A 9	99-20504-90	29605	29629	P9
A10	99-20493-238	42507	42531	P10
A11	99-20499-221	69312	69336	P11
A12	99-20499-364	69169	69193	P12
A13	99-20499-399	69134	69158	P13
A15	5-249-304	78583	78607	P15
A16	99-20485-269	82147	82171	P16
A17	99-20481-131	84510	84534	P17
A18	99-20481-419	84798	84822	P18
A19	99-20480-233	89955	89979	P19

Example 5:

Validation of the polymorphisms through microsequencing

The biallelic markers identified in example 4 were further confirmed and their respective frequencies were determined through microsequencing. Microsequencing was carried out for each individual DNA sample described in Example 2.

Amplification from genomic DNA of individuals was performed by PCR as described above for the detection of the biallelic markers with the same set of PCR primers (Table 1).

The preferred primers used in microsequencing were about 19 nucleotides in length and 10 hybridized just upstream of the considered polymorphic base. According to the invention, the primers used in microsequencing are detailed in Table 4.

Table 4

Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer mis 1 in SEQ ID No 1		ran	icing primer	
99-430-352	A1	D1	9475	9493	E1	9495	9513
Marker Name	Biallelic Marker	Mis. 1	Position range of microsequencing primer mis 1 in SEO ID No 2		Mis. 2	ran microseque	tary position ge of ncing primer EQ ID No 2
99-20508-456	A2	D2	1424	1442	E2	1444	1462
99-20469-213	A3	D3	5228	5246	E3	5248	5266
5-254-227	A4	D4	6204	6222	E4	6224	6242
5-257-353	A5	D 5	14704	14722	E 5	14724	14742
99-20511-32	A6	D6	19167	19185	E 6	19187	19205
99-20511-221	A7	D 7	18978	18996	E7	18998	19016

WO 00/08209 PCT/IB99/01444

99-20510-115	A8	D8	19872	19890	E8	19892	19910
99-20504-90	A9	D9	29598	29616	E9	29618	29636
99-20493-238	A10	D10	42500	42518	E10	42520	42538
99-20499-221	A11	D11	69305	69323	E11	69325	69343
99-20499-364	A12	D12	69162	69180	E12	69182	69200
99-20499-399	A13	D13	69127	69145	E13	69147	69165
99-20473-138	A14	D14	76439	76457	E14	76459	76477
5-249-304	A15	D15	78576	78594	E15	78596	78614
99-20485-269	A16	D16	82140	82158	E16	82160	82178
99-20481-131	A17	D17	84503	84521	E17	84523 -	84541
99-20481-419	A18	D18	84791	84809	E18	84811	84829
99-20480-233	A19	D19	89948	89966	E19	89968	89986

The microsequencing reaction was performed as follows:

After purification of the amplification products, the microsequencing reaction mixture was prepared by adding, in a 20µl final volume: 10 pmol microsequencing oligonucleotide, 1 U

Thermosequenase (Amersham E79000G), 1.25 µl Thermosequenase buffer (260 mM Tris HCl pH 9.5, 65 mM MgCl₂), and the two appropriate fluorescent ddNTPs (Perkin Elmer, Dye Terminator Set 401095) complementary to the nucleotides at the polymorphic site of each biallelic marker tested, following the manufacturer's recommendations. After 4 minutes at 94°C, 20 PCR cycles of 15 sec at 55°C, 5 sec at 72°C, and 10 sec at 94°C were carried out in a Tetrad PTC-225 thermocycler (MJ Research). The unincorporated dye terminators were then removed by ethanol precipitation. Samples were finally resuspended in formamide-EDTA loading buffer and heated for 2 min at 95°C before being loaded on a polyacrylamide sequencing gel. The data were collected by an ABI PRISM 377 DNA sequencer and processed using the GENESCAN software (Perkin Elmer).

Following gel analysis, data were automatically processed with software that allows the determination of the allèles of biallelic markers present in each amplified fragment.

The software evaluates such factors as whether the intensities of the signals resulting from the above microsequencing procedures are weak, normal, or saturated, or whether the signals are ambiguous. In addition, the software identifies significant peaks (according to shape and height criteria). Among the significant peaks, peaks corresponding to the targeted site are identified based on their position. When two significant peaks are detected for the same position, each sample is categorized classification as homozygous or heterozygous type based on the height ratio.

References

Altschul et al., 1990, J. Mol. Biol. 215(3):403-410 / Altschul et al., 1993, Nature Genetics 3:266-272 / Altschul et al., 1997, Nuc. Acids Res. 25:3389-3402 / Ausubel et al. (1989)Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. / Beaucage et al., Tetrahedron Lett 1981, 22: 1859-1862 / Bram RJ et al., 1993, Mol. Cell Biol., 13: 4760-4769. / Brown EL, Belagaje R, Ryan MJ, Khorana HG, Methods Enzymol 1979;68:109-151 / Castagnoli L. et al. (Felici F.), 1991, J. Mol. Biol., 222:301-310. /

- Chai H. et al., 1993, Biotechnol. Appl. Biochem., **18**:259-273 / Chee et al. (1996) *Science*. 274:610-614. / Chen and Kwok *Nucleic Acids Research* 25:347-353 1997 / Chen et al. *Proc. Natl. Acad. Sci. USA* 94/20 10756-10761,1997 / Cho RJ et al., 1998, Proc. Natl. Acad. Sci. USA, **95**(7): 3752-3757. / Chumakov I. et al., 1995, Nature, **377**(6547 Suppl): 175-297. / Compton J.
- 5 (1991) Nature. 350(6313):91-92. / Dib et al., 1996, Nature, 380: III-V. / Ellis NA,1997
 Curr.Op.Genet.Dev., 7: 354-363 / Feldman and Steg, 1996, Medecine/Sciences, synthese, 12:47-55 / Fields and Song, 1989, Nature, Vol. 340: 245-246. / Fishel R & Wilson T. 1997,
 Curr.Op.Genet.Dev.7: 105-113 / Flotte et al., 1992, Am. J. Respir. Cell Mol. Biol., 7: 349-356. / Fodor et al. (1991) Science 251:767-777. / Fromont-Racine M. et al., 1997, Nature Genetics,
- 16(3): 277-282. / Fuller S.A. et al., 1996, Immunology in Current Protocols in Molecular Biology, Ausubel et al. Eds, John Wiley & Sons, Inc., USA / Geysen H. Mario et al. 1984. Proc. Natl. Acad. Sci. U.S.A. 81:3998-4002 / Gonnet et al., 1992, Science 256:1443-1445 / Green et al., Ann. Rev. Biochem. 55:569-597 (1986) / Grompe, M. et al., Proc. Natl. Acad. Sci. U.S.A 1989; 86:5855-5892 / Grompe, M. Nature Genetics 1993; 5:111-117 / Guatelli J C et al. Proc. Natl.
- 15 Acad. Sci. USA. 35:273-286. / Haber D & Harlow E, 1997, Nature Genet. 16:320-322. / Hacia JG, Brody LC, Chee MS, Fodor SP, Collins FS, Nat Genet 1996;14(4):441-447 / Haff L. A. and Smirnov I. P. (1997) Genome Research, 7:378-388. / Hames B.D. and Higgins S.J. (1985) Nucleic Acid Hybridization: A Practical Approach. Hames and Higgins Ed., IRL Press, Oxford. / Harju L, et al., Clin Chem 1993;39(11Pt 1):2282-2287 / Harper JW et al., 1993, Cell, Vol. 75:
- 805-816. / Harris H et al.,1969, Nature 223:363-368. / Henikoff and Henikoff, 1993, Proteins 17:49-61 / Higgins et al., 1996, Methods Enzymol. 266:383-402 / Hillier L. and Green P. Methods Appl., 1991, 1: 124-8. / Huang L. et al. (1996) Cancer Res 56(5):1137-1141. / Huygen et al., 1996, Nature Medicine, 2(8):893-898 / Izant and Weintraub, Cell 36:1007-1015 (1984) / Julan et al., 1992, J. Gen. Virol., 73: 3251 3255. / Karlin and Altschul, 1990, Proc. Natl. Acad.
- Sci. USA 87:2267-2268 / Koch Y., 1977, Biochem. Biophys. Res. Commun., 74:488-491 / Kohler G. and Milstein C., 1975, Nature, 256: 495. / Kozal MJ, et al., Nat Med 1996;2(7):753-759 / Landegren U. et al. (1998) Genome Research, 8:769-776. / Leger OJ, et al., 1997, Hum Antibodies, 8(1): 3-16 / Lenhard T. et al., 1996, Gene, 169:187-190 / Livak et al., Nature Genetics, 9:341-342, 1995 / Livak KJ, and Hainer JW., 1994, Hum Mutat., 3(4): 379-385. /
- Lockhart et al. Nature Biotechnology 14: 1675-1680, 1996 / Lucas A.H., 1994, In: Development and Clinical Uses of Haempophilus b Conjugate. / Mansour SL et al., 1988, Nature, 336: 348-352.
 / Marshall R. L. et al. (1994) PCR Methods and Applications. 4:80-84. / Martineau P, Jones P, Winter G, 1998, J Mol Biol, 280(1):117-127 / Mc Whorter W.P., et al. A screening study of prostate cancer in high risk families. J Urol 1992;148:826-828. / McLaughlin et al., 1989, J. Virol.,
- 62: 1963 1973. / Muzyczka et al., 1992, Cuur. Topics in Micro. and Immunol., 158: 97-129. /
 Narang SA, Hsiung HM, Brousseau R, Methods Enzymol 1979;68:90-98 / Neda et al., 1991, J.
 Biol. Chem., 266: 14143 14146. / Nickerson D.A. et al. (1990) Proc. Natl. Acad. Sci. U.S.A.

1119.

79 87:8923-8927. / Nyren P, Pettersson B, Uhlen M, Anal Biochem 1993;208(1):171-175 / O'Reilly et al., 1992, Baculovirus expression vectors: a Laboratory Manual. W.H. Freeman and Co., New York / Ohno et al., 1994, Sciences, 265:781-784 / Oldenburg K.R. et al., 1992, Proc. Natl. Acad. Sci., 89:5393-5397. / Orita et al., Proc. Natl. Acad. Sci. U.S.A. 1989;86: 2776-2770 / 5 Parmley and Smith, Gene, 1988, 73:305-318. / Pastinen et al., Genome Research 1997; 7:606-614 / PCR Methods and Applications", 1991, Cold Spring Harbor Laboratory Press. / Pearson and Lipman, 1988, Proc. Natl. Acad. Sci. USA 85(8):2444-2448 / Pietu et al. Genome Research 6:492-503, 1996 / Porath J et al., 1975, Nature, 258(5536): 598-599. / Reimann KA, et al., 1997, AIDS Res Hum Retroviruses. 13(11): 933-943 / Ridder R, et al., 1995, Biotechnology (N Y), 10 13(3):255-260 / Rossi et al., Pharmacol. Ther. 50:245-254, (1991) / Roth J.A. et al., 1996, Nature Medicine, 2(9):985-991 / Rougeot, C. et al., Eur. J. Biochem. 219 (3): 765-773, 1994 / Roux et al., 1989, Proc. Natl Acad. Sci. USA, 86: 9079 - 9083. / Sambrook, et al. 1989. Molecular cloning: a laboratory manual. 2ed. Cold Spring Harbor Laboratory, Cold spring Harbor, New York. / Samson M, et al. (1996) Nature, 382(6593):722-725. / Samulski et al., 1989, J. Virol., 63: 3822-3828. / Sanchez-Pescador R., 1988, J. Clin. Microbiol., 26(10):1934-1938 / Sandou et al., 1994, Science, 265: 1875-1878. / Schena et al. Science 270:467-470, 1995 / Schwartz and Dayhoff, eds., 1978, Matrices for Detecting Distance Relationships: Atlas of Protein Sequence and Structure, Washington: National Biomedical Research Foundation / Sczakiel G. et al., 1995, Trends Microbiol., 1995, 3(6):213-217 / Sheffield, V.C. et al, Proc. Natl. Acad. Sci. U.S.A 1991; 20 49:699-706 / Shoemaker DD, et al., Nat Genet 1996;14(4):450-456 / Smith et al., 1983, Mol. Cell. Biol., 3:2156-2165. / Sosnowski RG, et al., Proc Natl Acad Sci USA 1997;94:1119-1123 / Steinberg G.D., et al. Family history and the risk of prostate cancer, The prostate 1990;17,337-347. / Stryer, L., Biochemistry, 4th edition, 1995 / Syvanen AC, et al., 1994, Hum Mutat., 3(3): 172-179. / Tacson et al., 1996, Nature Medicine, 2(8):888-892. / Thompson et al., 1994, Nucleic 25 Acids Res. 22(2):4673-4680 / Tyagi et al. (1998) Nature Biotechnology. 16:49-53. / Urdea M.S., 1988, Nucleic Acids Research, 11: 4937-4957 / Urdea MS et al., 1991, Nucleic Acids Symp Ser., 24: 197-200. / Valadon P., et al., 1996, J. Mol. Biol., Vol. 261:11-22. / Vaughan TJ, et al., 1996, Nat Biotechnol. 14(3): 309-314 / Vlasak R. et al., 1983, Eur. J. Biochem., 135:123-126 / Wabiko et al., 1986, DNA, 5(4):305-314. / Walker et al. (1996) Clin. Chem. 42:9-13. / 30 Westerink M.A.J., 1995, Proc. Natl. Acad. Sci., 92:4021-4025. / White, M.B. et al. (1992) Genomics. 12:301-306. / White, M.B. et al. (1997) Genomics. 12:301-306. / Wilson R. et al., 1994, Nature, 368(6466): 32-38. / Zhang SD et al., 1996, Genes and development, 10: 1108-

SEQUENCE LISTING FREE TEXT

The following free text appears in the accompanying Sequence Listing:
5' regulatory region

polymorphic base complement

3' regulatory region

deletion of

5 or

probe

homology with Genset 5' EST in ref sequencing oligonucleotide PrimerPU sequencing oligonucleotide PrimerRP

10

15

20

25

30

35

81

CLAIMS

- 1. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 60 nucleotides of SEQ ID No. 1 or the complements thereof.
- 2. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 60 nucleotides of SEQ ID No. 2 or the complements thereof.
- 3. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 60 nucleotides of SEQ ID No. 3 or the complements thereof.
- 4. An isolated, purified, or recombinant polynucleotide comprising a contiguous span of at least 60 nucleotides of SEQ ID No 4. or the complements thereof.
- 5. An isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides of anyone of SEQ ID Nos. 1 and 2 or the complement thereof, wherein said span includes a *TBC-1*-related biallelic marker in said sequence.
- 6. A polynucleotide according to claim 5, wherein said *TBC-I*-related biallelic marker is selected from the group consisting of the biallelic markers in positions 9494 of the SEQ ID No. 1, and 1443, 5247, 6223, 14723, 19186, 18997, 19891, 29617, 42519, 69324, 69181, 69146, 76458, 78595, 82159, 84522, 84810, and 89967 of the SEQ ID No. 2.
- 7. A polynucleotide according to any one of claims 5 or 6, wherein said contiguous span is 18 to 35 nucleotides in length and said biallelic marker is within 4 nucleotides of the center of said polynucleotide.
- 8. A polynucleotide according to claim 7, wherein said polynucleotide consists of said contiguous span and said contiguous span is 25 nucleotides in length and said biallelic marker is at the center of said polynucleotide.
- 9. A polynucleotide according to claim 8, wherein said polynucleotide consists essentially of a sequence selected from the sequences with the position range 9482-9506 in SEQ ID No. 1 and with the following position ranges in SEQ ID No. 2: 1431-1455, 5235-5259, 6211-6235, 14711-14735, 19174-19198, 18985-19009, 29605-29629, 42507-42531, 69312-69336, 69169-69193, 69134-69158, 78583-78607, 82147-82171, 84510-84534, 84798-84822, and 89955-89979, and the complementary sequences thereto.
- 10. A polynucleotide according to any one of claims 1 to 6, wherein the 3' end of said contiguous span is present at the 3' end of said polynucleotide.
 - 11. A polynucleotide according to any one of claims 5 or 6, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide and said biallelic marker is present at the 3' end of said polynucleotide.
- 12. An isolated, purified, or recombinant polynucleotide consisting essentially of a contiguous span of 8 to 50 nucleotides of anyone of SEQ ID Nos. 1 and 2 or the complement thereof, wherein the 3' end of said contiguous span is located at the 3' end of said polynucleotide,

10

15

20

25

30

and wherein the 3' end of said polynucleotide is located within 20 nucleotides upstream of a TBC-I-related biallelic marker in said sequence.

- 13. A polynucleotide according to claim 12, wherein the 3' end of said polynucleotide is located 1 nucleotide upstream of said *TBC-1*-related biallelic marker in said sequence.
- 14. A polynucleotide according to claim 13, wherein said polynucleotide consists essentially of a sequence selected from the sequences with the position range 9475-9493 in SEQ ID No. 1 and with the following position ranges in SEQ ID No 2: 1424-1442, 5228-5246, 6204-6222, 14704-14722, 19167-19185, 18978-18996, 19872-19890, 29598-29616, 42500-42518, 69305-69323, 69162-69180, 69127-69145, 76439-76457, 78576-78594, 82140-82158, 84503-84521, 84791-84809, and 89948-89966, and the complementary position range 9495-9513 in SEQ ID No. 1 and the following complementary position ranges in SEQ ID No 2: 1444-1462, 5248-5266, 6224-6242, 14724-14742, 19187-19205, 18998-19016, 19892-19910, 29618-29636, 42520-42538, 69325-69343, 69182-69200, 69147-69165, 76459-76477, 78596-78614, 82160-82178, 84523-84541, 84811-84829, and 89968-89986.
- 15. An isolated, purified, or recombinant polynucleotide consisting essentially of a sequence selected from the sequences with the position range 9391-9408 in SEQ ID No 1 and with the following position ranges in SEQ ID No 2: 988-1006, 5039-5056, 5997-6015, 14371-14390, 18751-18771, 19605-19625, 29529-29547, 42268-42287, 69026-69046, 76323-76343, 78292-78309, 81893-81912, 84392-84412, and 89746-89765, and the complementary position range 9828-9845 in SEQ ID No 1 and the following complementary position ranges in SEQ ID No 2: 1509-1529, 5534-5554, 6332-6350, 14798-14817, 19198-19217, 19986-20005, 30041-30061, 42732-42752, 69525-69543, 76771-76790, 78704-78721, 82353-82372, 84909-84929, and 90179-90198.
- 16. An isolated, purified, or recombinant polynucleotide which encodes a polypeptide comprising a contiguous span of at least 6 amino acids of SEQ ID No 5.
 - 17. A polynucleotide according to any one of claims 1 to 16 attached to a solid support.
 - 18. An array of polynucleotides comprising at least one polynucleotide according to claim 17.
 - 19. An array according to claim 18, wherein said array is addressable.
 - A polynucleotide according to any one of claims 1 to 16 further comprising a label.
 - 21. A recombinant vector comprising a polynucleotide according to any one of claims 1 to 4 and 16.
 - 22. A host cell comprising a recombinant vector according to claim 21.
- 23. A non-human host animal or mammal comprising a recombinant vector according to claim 22.
 - 24. A method of genotyping comprising determining the identity of a nucleotide at a *TBC-1*-related biallelic marker or the complement thereof in a biological sample.

10

15

20

- 25. A method according to claim 24, wherein said biological sample is derived from a single subject.
- 26. A method according to claim 25, wherein the identity of the nucleotides at said biallelic marker is determined for both copies of said biallelic marker present in said individual's genome.
- 27. A method according to claim 24, wherein said biological sample is derived from multiple subjects.
- 28. A method according to claim 24, further comprising amplifying a portion of said sequence comprising the biallelic marker prior to said determining step.
 - 29. A method according to claim 28, wherein said amplifying is performed by PCR.
- 30. A method according to claim 24, wherein said determining is performed by a hybridization assay.
- 31. A method according to claim 24, wherein said determining is performed by a sequencing assay.
- 32. A method according to claim 24, wherein said determining is performed by a microsequencing assay.
- 33. A method according to claim 24, wherein said determining is performed by an enzyme-based mismatch detection assay.
- 34. A method according to any one of claims 24 to 33 wherein said *TBC-I*-related biallelic marker is selected from the group consisting of the biallelic markers in positions 9494 of the SEQ ID No. 1, and 1443, 5247, 6223, 14723, 19186, 18997, 19891, 29617, 42519, 69324, 69181, 69146, 76458, 78595, 82159, 84522, 84810, and 89967 of the SEQ ID No. 2.
- 35. An isolated, purified, or recombinant polypeptide comprising a continuous span of at least 8 amino acids of SEQ ID No 5.
- 25 36. An isolated or purified antibody composition capable of selectively binding to an epitope-containing fragment of a polypeptide according to claim 35.

This is to be your

1/5 Figure 1

Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	mpmlpwvvae		kepvtkqvrl	WVSPSGLRCE cvspsglrce ~~~~~~	pepgrsqqwd
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus				LIKEDAAHRQ likedavhrq	sicyvfkadd
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	101 QTKVPEIISS qtkvpeiiss ~~~~~~	irqagkiarq	eelhcpsefd	DTFAKKFEVL dtfskkfevl ~~~~~~	
Mur. tbc1 TBC-1 dmu50542 Celf35h12 Consensus				QPSA pmprpqwsqe ~~~~~~	
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus				NDIENHLIGG sdienhlisg	
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	enrtmlftig	qsevylispd	tkkialeknf	KEISFCSQGI keisfcsqgi	rhvdhfgfic
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	MEDFKDFTEV	fhfvcyvfqc TQFTNVQYLG	tnealvdeim CSQLVNNDND	MTLKQAFTVA mtlkqaftva ~~~~~ NEMKALMKVL KV-	avqqtaka.p DEQKGAQTIN

PCT/IB99/01444

2/5 Figure 1 (Continued I)

		rigare i (C	Ontinued 1)		400
Mur. tbc1	351 AQLCEGCPLQ	GLHKLCERIE	GMNSSKTKLE	LQKHLTTLTN	400 QEQATIFEEV
TBC-1		sinkicerie	gmnssktkle	rdkureeren	
dmu50542	· · · · · · · · · · · · · · · · · · ·		KVLSSFSLVN		
celf35h12 Consensus	VILVVPHNIS	GIVKLIDAQG	SS	TRECINGESS	OF
Compensus	D	143	20		x. -
	401			•	450
Mur. tbc1	QKLRPRNEQR	ENELIISFLR	CLYEEKQKEH	SHTGAPKQTL	QVAAENIGSD
TBC-1	qklrprneqr	eneliisflr	clyeekqkeh	ihigemkqts	qmaaenigse
. dmu50542	~~~~~~	~~~~~~~	~~~~~~~	~~~~~~	~~~~~~~
celf35h12			HVFRTSKAET		
Consensus		-NI	E-		NS-
	451				500
Mur= tbc1	451	DSLKNRAKRS	LTESLESILS	RGNKARGLOD	
TBC-1			lteslesils		
dmu50542		~~~~~~	~~~~~~	~~~~~~	~~~~~~~~
celf35h12	SNRLEFQFES	ILEVKENDGT	VEKPSWKLCP	QHNGVFKVRR	DREKKIVVQL
Consensus	F			N	
man a					==0
	501	01/0DIFE	DIJOH MARKITA	acabbi caba	550 ECUTABECAT
Mur. tbcl			PVSETSFKLL		
TBC-1 dm150542	Isstishtsk	epsvcekear	pisessfkll	gasedissus	eshipeepap
cel£35h12	BUMBEKKAID ≈≈≈≈≈≈≈≈≈	GELLNIKKCE	GMLLAAGRNL	RHSDLOLLEM	DRNATGTDSA
Consensus	T	KF	L	S	
Secretary Secretary	551				600
Mur. tbc1	LSPQQAFRRR	\ANTLSHFPVE	CPAPPEPAQS	SPGVSQRKLM	RYHSVSTETP
TBC-1	lspqqafrrr	antļshfpie	cqeppqparg	sbansdrkTw	rynsvstetp
dmu50542	~~~~~~~~	DDDIMMERIA	NTETPRDTRV	EIMITIA I DUITU	QETQEDIDEQ
celf35h12	VFVIEANW	DPKVHMFEVL	NIEIPRDIRV	FHIVALDOIV	S
Consensus	A	F	F	•	•
	601				650
Mur. tbc1	HERKDFESKA	NHLGDTDGTP	VKTRRHSWRQ	QIFLRVATPQ	KACDSPSRYE
TBC-1	herkdfeska	nhlgdsggtp	vktrshswrq	qiflrvatpq	kacdsssrye
dmu50542	~~~~~~~~		~~~~~~~~~		~~~~~~~
celf35h12	MEAMSRVFHE	HERFYKTPQT	VVSEEFTLVL	EVRIRTEKLL	ETHGNMLLKS
Consensus-	E	V		R	
	451				700
M 4 2 1	651	ם מטונסט ביו	GPFGQYRKKR	GRRHASFESC	,
Mur. tbcl TBC-1	dygal cal nr	rsplenvced	gpfaphqrkr	kahlvssesc	akrlffnryc
dmu50542	ayacigeipp		~~MRKPAKRG	KRDAAELREL	WRTAIRQTIM
celf35h12	VPIDFAWOLE	GVYFLPTPSK	SCDQSDPNDR	KLTFISLESD	SDRKRSKQNL
Consensus		P	R	KS-ES-	R

PCT/IB99/01444

4/5 Figure 1 (Continued III)

	F	igure I (Co	ntinuea III	.)	
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	dnqrmdklek TNHHLEMLNR NEDPVLRMEK	tnsslrkqnl EKTQNQ EIGRHQANTL	dll.eqlqva HLE.QQLQFA RLERENDDLA	NARIQSLEAT ngriqsleat QSSIAQLETT HELVTSKIEL I-SLE-T	iekllssesk RSSQQAQITT RRKLDVAEDQ
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	lkqamltlel LQSQVQSLEL IETSANAIER	ersallqtve TIQTLGRYVG LTRQNMDILE	elrrrsaeps QLVEHNP ENKNLMRE	SR.ARLHPAG drepectqpe DLELPNEVRR YEQIKEMYRR	ptgd~~~~ MLQQLDDLDR DVLRLEENGS
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	QRRKPIFTER RAEKLLAEYK	KIGKSVSVNS	HLGFPLKVLE ENEREHFEVQ	ELTERDELGS KKAIIARISD	PQKQKKEKTP
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	FFEQLRQQQQ .WEKNRSPVH	QHRLNGGGQS SASTPTGPDL	SNVGESGSPT LTKLEEREDH	PPSRPNRLLD IKNLEIDLAQ	NASARTVMQV TKLSLVEAEC
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	~~~~~~ KLDELKLPEH	VDKFVANIKS	PLEVDSGVGT	PLSPPSTASN VGSSLKHHER	SSGGSIFSRM
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus		PLAQRQSYGV	AITTAPCPQH	MEEVAPATTM FMQTPSAVLK	AVMPQEDVEE
Mur. tbc1 TBC-1 dmu50542 celf35h12 Consensus	1351 PQPMHPLSMV NNMLHGINGV	GGDVNVRFKG	TTQLKSIRPV DDQYSNSSSL	HHMRAIPLGG	1400 VQHPSSTEPA GKAEDSTMVT

THE STREET STREET

5/5 Figure 1 (Continued IV)

	· · · · · · · · · · · · · · · · · · ·				
	1401				1450
Mur. tbcl	,	~~~~~~	~~~~~~~	~~~~~~	~~~~~~~
TBC-1	~~~~~~~	~~~~~~	~~~~~~~	~~~~~~	~~~~~~~
dmu50542		APPAATATTG			~~~~~~
celf35h12	VNLDQLPARR	TLLKCLVLVV	FGKSYTKVIF	QMLRFSGFLM	RRGVYPGTLY
Consensus					
	1451	1465		-	
Mur. tbc1	~~~~~~	~~~~			
TBC-1	~~~~~~~	~~~~		•	
dmu50542	~~~~~~	~~~~			
celf35h12	FRRCSKILLK	KYDRI			
Consensus					

Please type a	plus	sıgn	(+)	ınsıde	this box	\rightarrow	<u> +</u>
---------------	------	------	-----	--------	----------	---------------	-----------

PTO/SB/01 (10-00)
Approved for use through 10/31/2002. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION FOR UTILITY OR
DESIGN
PATENT APPLICATION
(37 CFR 1.63)

☐ Declaration Submitted with Initial Filing

OR

Declaration Submitted after Initial Filing (surcharge (37 CFR 1.16 (e)) required)

Attorney Docket Num	ber	46.US2.PCT		
First Named Inventor		Marta BLUMENFELD, et al.		
COMPLETE I		KNOWN		
Application Number	0	9 / 762,311		
Filing Date	February 1, 2001			
Group Art Unit	unknown			
Examiner Name	unassigned			

As a below named inventor, I hereby declare that:									
My residence, mailing address, and citizenship are as stated below next to my name.									
I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:									
NUCLEIC ACIDS ENCODING HUMAN TBC-1 PROTEIN AND POLYMORPHIC MARKERS THEREOF									
	(7	itle of the Invention)							
the specification of which	•	•							
☐ is attached hereto									
OR		as United S	tates Application I	Number or PCT International					
was filed on (MM/DD/YYYY)	02/01/2001								
Application Number 09/762	2,311 and was a	mended on (MM/DD/YY	YY) 02/01.	(if applicable).					
I hereby state that I have reviewed amended by any amendment spe	d and understand the co	ontents of the above idea	ntified specification	n, including the claims, as					
I acknowledge the duty to disclose in-part applications, material informed PCT international filing date of the	nation which became a	vailable between the filir	s defined in 37 CF ng date of the prio	R 1.56, including for continuation- r application and the national or					
I hereby claim foreign priority ben certificate, or 365(a) of any PCT i America, listed below and have certificate, or any PCT international	international application also identified below.	which designated at lead by checking the box, a	ast one country of any foreign applic	ther than the United States of attornion for patent or inventor's					
Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached? YES NO					
PCT/IB99/01444	WIPO	08/06/1999	0000						
Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:									
I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.									
Application Number(s)	Filing Dat	e (MM/DD/YYYY)	numbers supplem	al provisional application s are listed on a ental priority data sheet /02B attached hereto.					
	1								

[Page 1 of 2]
Burden Hour Statement: This form is estimated to take 21 minutes to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

DECLARATION — Utility or Design Patent Application

-								
Direct all o	orrespondence to:	Customer Nu or Bar Code L		000027	7206	OR X	Correspondence	address below
Name John Lucas, Ph.D., J.D.								
Address	Genset Corporat	ion						
Address	10665 Sorrento	Valley R	oad					
City	San Diego				State	CA	92121 ZIP	-1609
Country	USA		Telephon		597-260	00	858/597 Fax	7-2601
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.								
NAME OF	F SOLE OR FIRST IN	/ENTOR:			A petiti	on has been fil	ed for this uns	igned inventor
Given Name	Marta				Family I		IFELD	
Inventor's Signature							Date	
Residence:	City	aris		State		Country Franc	e Citizenship	French
Mailing Add	lress 24 rue Royal	.e		·-				
Mailing Add	Iress	''						
City	Paris	State			ZIP	75008	Country	France
NAME OF	SECOND INVENTO	₹:			A petiti	on has been fi	led for this uns	signed inventor
Given Name	e iddle [if any])	Lydie	9		Family I		BOUGUELE	RET
Inventor's Signature							Date	
Residence:	City Petit	Lancy		State		Switzerlar Country	d Citizenship	French
Mailing Address 24 rue Royale								
Mailing Address								
City	Paris	State			ZIP	75008	Country	France
	al inventors are being name		_suppleme	ental Additio		tor(s) sheet(s) PT		ed hereto.

Į
Transfer or the second
U
11.00
Ħ
Inhapping.
T.
T
Tun Hail

Please type a plus sign (+) inside this box —	PTO/SB/02A (11-00)
	Approved for use through 10/31/2002. OMB 0651-0032
	U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
nder the Paperwork Reduction Act of 1995, no persons are required to rest	cond to a collection of information unless it contains a valid OMB control number

DECLARATION

ADDITIONAL INVENTOR(S) Supplemental Sheet Page ___ of ___

Name of Additional Joint Inventor, if a	ny:		A petition has been filed for this unsigned inventor		
Given Name (first and middle [if any])		Family Name	or Su	ırname
llya			СН	JMAŁ	(OV
Inventor's Signature				1	Date
Vaux-le-Penil Residence: City	State		France	c	French Sitizenship
Mailing Address 24 rue Royale					
Mailing Address				****	
City Paris	State		ZIP 75008 C	ountry	, France
Name of Additional Joint Inventor, if a	ny:		A petition has been filed	for this	unsigned inventor
Given Name (first and middle [if any])		Family Name	or Su	ırname
Inventor's Signature					Date
Residence: City	State		Country		Citizenship
Mailing Address			·		
Mailing Address					
City	State		ZIP	Coun	itry
Name of Additional Joint Inventor, if a	ny:		A petition has been filed fo	or this	unsigned inventor
Given Name (first and middle [if any])	Family Name or Surname			
	<u> </u>				
Inventor's Signature					Date
Residence: City	State		Country		Citizenship
Mailing Address					
Mailing Address			T	1	100
City	State		ZIP	Co	untry

Burden Hour Statement: This form is estimated to take 21 minutes to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time—you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

Please type a plus sign (+) inside this box	\rightarrow	+	
---	---------------	---	--

PTO/SB/01 (10-00)

Approved for use through 10/31/2002. OMB 0651-0032

U.S. Patent and Trademark Office; U.S DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number

46.US2.PCT Attorney Docket Number **DECLARATION FOR UTILITY OR** Marta BLUMENFELD, et al. **First Named Inventor DESIGN COMPLETE IF KNOWN** PATENT APPLICATION 762,311 09 (37 CFR 1.63) Application Number February 1, 2001 Filing Date ☐ Declaration □ Declaration Submitted after Initial unknown OR Submitted Group Art Unit Filing (surcharge (37 CFR 1.16 (e)) with Initial unassigned Filing **Examiner Name** required)

As a below named inventor, I hereby declare that:								
My residence, mailing address, and	My residence, mailing address, and citizenship are as stated below next to my name.							
I believe I am the original, first and names are listed below) of the sub	sole inventor (if only or ject matter which is clai	ne name is listed below) med and for which a pat	or an original, first ent is sought on t	t and joint invent ne invention ent	tor (if plural itled:			
NUCLEIC ACIDS ENCODING HUMAN TBC-1 PROTEIN AND POLYMORPHIC MARKERS THEREOF								
<u> </u>	(7	itle of the Invention)						
the specification of which	(.	and of the invention						
is attached hereto								
OR		as United St	ates Application N	Number or PCT	International			
was filed on (MM/DD/YYYY)	02/01/2001		. фр					
A P. C. Nimb					applicable).			
Application Number 09/762	2.311 and was a	mended on (MM/DD/YY	YY) 02/01/	2001				
I hereby state that I have reviewed amended by any amendment spe-	d and understand the co cifically referred to abov	ontents of the above ider re.	ntified specification	n, including the	claims, as			
I acknowledge the duty to disclose in-part applications, material inforr PCT international filing date of the	nation which became a	vailable between the tilin	defined in 37 CF g date of the prior	R 1.56, includin r application and	g for continuation- I the national or			
I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or any PCT international application having a filing date before that of the application on which priority is claimed.								
Prior Foreign Application	Foreign Filing Date Priority Certified Copy Attached?				py Attached? NO			
Number(s)	Oddiay	(MM/DD/YYYY)	NOL Claimed	YES	NO			
PCT/IB99/01444	WIPO	08/06/1999			☑			
Additional foreign application numbers are listed on a supplemental priority data sheet PTO/SB/02B attached hereto:								
I hereby claim the benefit under 35 U.S.C. 119(e) of any United States provisional application(s) listed below.								
Application Number(s)	Filing Dat	e (MM/DD/YYYY)			.P 4			
				al provisional ap are listed on a	plication			
				ental priority dat	a sheet			
				02B attached h				
	i							

[Page 1 of 2]

Burden Hour Statement: This form is estimated to take 21 minutes to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time—you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

The state of the s

Please type a plus sign (+) inside this box

+ Approved for use through 10/31/2002 OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number.

DECLARATION — Utility or Design Patent Application

Direct all co		Customer Nur or Bar Code L		000027	7206	OR X	Correspondence a	address below
Name	Name John Lucas, Ph.D., J.D.							
Address	Genset Corporati	lon						
Address	10665 Sorrento	Valley Ro	oad					
City	San Diego				State	CA	92121- ZIP	-1609
Country	USA		Telephon	858/ e	597-260	0	858/597 Fax	-2601
I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.								
NAME OI	F SOLE OR FIRST IN\	/ENTOR:			A petiti	on has been fi	led for this uns	igned inventor
	Given Name Marta Family Name BLUMENFELD or Surname							
Inventor's Signature				·			Date	
Residence:	City Pa	ıris		State		Country Franc	e Citizenship	French
Mailing Add	dress 24 rue Royal	e		•				
Mailing Add	dress							
City	Paris	State			ZIP	75008	Country	France
) NAME OF	SECOND INVENTOR	₹:			A petiti	on has been fi	led for this uns	igned inventor
Given Name (first and m	e iddle [if any])	Lydie	2		Family I	ame	BOUGUELE	
Inventor's Signature			#			C.	25.	05.01
Residence:	City Petit	Lancy	•	State		Switzerlar Country		French
Mailing Add	dress 24 rue Royale	e						
Mailing Add	dress							
City	Paris	State			ZIP	75008	Country	France
D Addition	1			ntol Additi-	nal Inva-	tor(a) about(a) DT	O/SB/03A attacha	nd horoto

	72	
;	Ž	
.;		Minn
	J	ifere.
100,000	1	dian.
2000	Å	ŝ
	-	5
		Ŀ
	ž	
,	=	To the same
****	I	ilen.
******	1	ileans.
	- Fund	The same
	_	7
:	í	2

Please type a plus sign (+) inside this box —	PTO/SB/02A (11-00)
	Approved for use through 10/31/2002. OMB 0651-0032
U	S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to	a collection of information unless it contains a valid OMB control number

DECLARATION

ADDITIONAL INVENTOR(S) Supplemental Sheet Page ___ of ___

Name of Additional Joint Inventor, if any:				A petition has been filed for this unsigned inventor				
Given Name (first and middle [if any		Family Name or Surname						
llya			CHUMAKOV					
Inventor's Signature					Date			
Residence: City	State		Country France FX			French Citizenship		
Mailing Address 24 rue Royale								
Mailing Address								
_{City} Paris	State			75008 ZIP	Countr	y France		
Name of Additional Joint Inventor, if a	ny:			A petition has been filed	for this	s unsigned inventor		
Given Name (first and middle [if any	/])			Family Nan	ne or S	urname		
Inventor's Signature						Date		
Residence: City	State	•	Country Citizenship			Citizenship		
Mailing Address								
Mailing Address								
				l				
City	State	9		ZIP	Cou	ntry		
Name of Additional Joint Inventor, if a	iny:		ПА	petition has been filed	for this	unsigned inventor		
Given Name (first and middle [if any])		Family Name or Surname					
Inventor's Signature		Date			Date			
Residence: City	State			Country		Citizenship		
Mailing Address								
Mailing Address								
Cit.	04-4-			710				

Burden Hour Statement: This form is estimated to take 21 minutes to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time—you are required to complete this form should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231.

PTO/SB/01 (10-00)
Approved for use through 10/31/2002. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it contains a valid OMB control number

DECLARATION — Utility or Design Patent Application

Direct all c		customer Num r Bar Code La		000027	206	OR X	Correspondence a	ddress below	
Name John Lucas, Ph.D., J.D.									
Address Genset Corporation									
Address 10665 Sorrento Valley Road									
City	San Diego				State	CA	92121- ZIP	-1609	
Country	USA	T	elephone		597-260	0	858/597 Fax	-2601	
are believe made are p	I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.								
NAME O	F SOLE OR FIRST INV	ENTOR:			A petiti	on has been fi	led for this unsi	igned inventor	
-	Given Name Marta Family Name BLUMENFELD or Surname								
Inventor's Signature Date 26/4/01									
Residence: City Paris State Country France Citizenship French									
Mailing Add	Mailing Address								
Mailing Add	dress								
City	Paris	State			ZIP	75008	Country	France	
NAME OF	SECOND INVENTOR	:			A petiti	on has been f	iled for this uns	igned inventor	
Given Name	e iddle [if any])	Lydie			Family I		BOUGUELE	KET	
Inventor's Signature Date									
Residence:	City Petit	Lancy		8tate		Switzerlan Country	nd Citizenship	French	
Mailing Address 24 rue Royale									
Mailing Address									
City	Paris	State	-,		ZIP	75008	Country	France	
	al inventors are being name		suppleme	ntal Additio	nal Inven	tor(s) sheet(s) P1	ΓΟ/SB/02A attache	ed hereto.	

PTO/SB/81 (10-00) Approved for use through 10/31/2002. OMB 0651-0035

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it display a valid OMB control number.

POWER OF ATTORNEY OR AUTHORIZATION OF AGENT

Application Number	09/762,311						
Filing Date	February 1, 2001						
First Named Inventor	Marta BLUMENFELD, et al.						
Group Art Unit	Unknown						
Examiner Name	Unassigned						
Attorney Docket Number	46.US2.PCT						

l hereby appoir	nt:						ا ٦	
Practition		→ N	Place Customer Number Bar Code abel here					
× Practition	ər(s) nar	ned below:						
		Name			Registration I	Number		
		John M. Lucas	43,37					
		Peter Follette			46,21	46,213		
		Lukas R. Voellmy	1		43,358			
as my/our attorn	ney(s) or United S	r agent(s) to prosecut States Patent and Tra	te the application ademark Office co	identif	ied above, and ed therewith.	to transact all		
		espondence address f ed Customer Number		ntified	application to:			
X Firm or								
Individual Na	me		-					
Address		Genset Corporation	w Pood					
Address		10665 Sorrento Valle	y Road	G	CA	Zip 92121-160	 19	
City		San Diego		State	CA	Zip 92121-160		
Country		USA (959) 507 2600		<u></u>	(848) 597-2601			
Telephone		(858) 597-2600		Fax	(040) 081-2007			
I am the:	t/Invento	or.						
,								
Assignee Statemer	e of recc nt under	ord of the entire intere or 37 CFR 3.73(b) is ea	est. See 37 CFR (nclosed. (Form P	3.71. <i>TO/SB</i>	·/96).			
		SIGNATURE of A	Applicant or Assig	nee of	Record			
Name	Ilya Chu							
Signature		797/						
	0 6 01 101							
NOTE: Signatures of all the inventors or assignees of record of the entire interest or their representative(s) are required. Submit multiple								
forms if more than one	signature	is required, see below*.				•		
☐ *Total of		rms are submitted.						

Please type a plus sign (+) inside this

PTO/SB/81 (10-00)

Approved for use through 10/31/2002 OMB 0651-0035

U.S Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it display a valid OMB control number.

POWER OF ATTORNEY OR AUTHORIZATION OF AGENT

Application Number	09/762,311
Filing Date	February 1, 2001
First Named Inventor	Marta BLUMENMFELD, et al.
Group Art Unit	Unknown
Examiner Name	Unassigned
Attorney Docket Number	46.US2.PCT

l hereby appoi	nt:				<u> </u>		
Practition OR	ners at C	→ Nu	nce Customer mber Bar Code bel here				
× Practition	er(s) na	med below:					
		Name			Registration N	umber	
		John M. Lucas			43,373		
Peter Follette					46,213		
	Lukas R. Voellmy				43,358		
as my/our attorr business in the	ney(s) or United S	r agent(s) to prosec States Patent and T	ute the application rademark Office co	identifi nnecte	ed above, and to	o transact all	
		spondence address ed Customer Numb		ntified a	application to:		
X Firm or Individual Na	X Firm or John Lucas, Ph.D., J.D.						
Address		Genset Corporation					
Address		10665 Sorrento Vall	ley Road				
City		San Diego		State	CA	Zip 92121-1609	
Country		USA					
Telephone		(858) 597-2600		Fax	(848) 597-2601		
I am the:							
X Applican	t/Invent	or.					
Assignee	e of reco	ord of the entire inte	rest. See 37 CFR 3 enclosed. (Form P	5.71. TO/SB	/96).		
		SIGNATURE of	Applicant or Assign	nee of	Record		
	Marta Blumenfeld						
Name							
Signature		Vest 1					
Date							
NOTE: Signatures of all the inventors or assignees of record of the entire interest or their representative(s) are required. Submit multiple forms if more than one signature is required, see below*.							
☐ *Total of		ms are submitted.					

```
<110> Genset SA
```

 $<\!\!120\!\!>$ Nucleic acids encoding human TBC-1 protein and polymorphic markers thereof.

<130> D.18363

<150> US 60/095,653

<151> 1998-08-07

<160> 7

<170> Patent.pm

. <210> 1

<211> 17590

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> 1..2000

<223> 5' regulatory region

<220>

<221> exon

<222> 2001..2077

<223> exon 1

<220>

<221> exon

<222> 12292..12373

<223> exon 1b

<220>

<221> exon

<222> 12740..13249

<223> exon 2

<220>

<221> allele

<222> 9494

 $\langle 223 \rangle$ 99-430-352 : polymorphic base A or G

```
<220>
<221> primer_bind
<222> 9391..9408
<223> 99-430.rp
<220>
<221> primer_bind
<222> 9828..9845
<223> 99-430.pu complement
<220>
<221> primer_bind
<222> 9475..9493
<223> 99-430-352.mis
<220>
<221> primer_bind
<222> 9495..9513
<223> 99-430-352.mis complement
<220>
<221> primer_bind
<222> 9482..9506
<223> 99-430-352.probe
<220>
<221> misc feature
<222> 3953,4056,4167,4739,6217,6245,6860,9998..9999,10006,10012,10104
       10477,10822,10825,11095,11256,11273,11857..11858,11895..11896
       14057,15912..15913,16217..16218,16329..16330,17504
<223> n=a, g, c or t
 <400> 1
 aggacagtat ctagcacaat accccaaatc gactaactcc tccgtaaaga atagctacca
                                                                       60
 ctattgtgag agttttaagt caagctgtga ataaaactct tgggtccact taaaaatacc
                                                                       120
tcccctggat gtaagcatcc agggaaatca gggaatgcca taagacagcc ctaatctaaa
                                                                       180
 agcctacaag aagctcagtg ggcttcaagg aagacactgc tettggtacg atgaggaaac
                                                                       240
 ctggccctct atttgcctcc tgggccacag taatattgat aatagctgct gcttttagtt
                                                                       300
 gaggaccatg tacgtetgtg teactgcact ggccacttta ettacaettt cetgetttgt
                                                                       360
 cctcacaaag atcctgtaag gtgtgtattg gtcccattta gcaggtaaga caatgaagac
                                                                       420
 cagaggtcca gcaccttgcc taaaccacac ctgctgggat ttggattcaa gtccaaccgt
                                                                       480
```

WO 00/08209 PCT/IB99/01444

-						
acagctcaaa	cgctcagcca	cttccctaaa	gtccaccccc	agctacatta	agtaaaaaaa	540
tccagaaaga	tgccacctgg	gggtctggaa	ctgcctcctc	cgagcacccg	gctctcccct	600
ccctgcggac	tettetetgg	agaggatgtg	atgcttctta	cttttctcag	atccctctcc	660
ccaccctgcg	agtgacgttg	cgcctctgtg	cctggtggga	tagggatctg	ggagcttcgc	720
ctgttttttg	cacactgcca	tcccctagtc	ttagggagcg	agctctgtcc	cgcttttcac	780
atctccgcgt	ctttccttgc	actctacatc	accgctggga	atgtccccag	acctgatcgg	840
ggcatgcaca	ctggggtgtg	cgtgtgcgtg	tggtgtgtgt	tcctgcgcgt	gtgccgggct	900
cgcggggcag	gaaaaagcgc	ctaatccagg	ctctgcgtca	ctcccgcaat	tggttagaaa	960
tg gagtttcc	tggtgtttaa	tcccgggagg	gcacttcgcc	tt cgt tg ttt	cccagagtcc	1020
ctgattttcc	tgcctcgcat	gccagcgccc	catagggcat	ccgtgcctca	gttcacctct	1080
tgccatcctc	caaggacggg	gagaaggggt	aaggcggggg	agagcaaggt	ggcttggtcg	1140
ccccggccc	ccgccccca	tgttgtgtgc	agtttccacc	acgtctgttt	cggagggaga	1200
agagga g ggt	gcagatgagg	cgaggcgcct	tcgggagcgc	ggagagcggg	caggcagtgc	1260
cacctgctga	gagccactca	ggccgagcaa	gcggcgggca	gtgccacctg	ctataaatag	1320
gccgccaagg	acagggtgtg	cgactgtaca	tcccgccacg	agggcctgca	tcacgcgcgg	1380
ggccccgcgc	cccggctcc	ccagggaaac	gctgtgccca	gatcctgcgc	aggggtctgg	1440
atggggcggc	ggcccgagta	cttcccccct	attccccca	cagacactgg	ctgaggatgg	1500
cccgcgggct	tgggggcggg	gggtggcaag	gaggggaggg	aggccgcggc	ggacccgcag	1560
tgcagcagct	gttgctcgcg	tgtgactcgc	ccgtccgggc	cgtgctgccc	aggcacagtc	1620
acacggcgca	gtggggagga	ggaggacacc	gagtccccct	cccagctccc	cggggaccga	1680
gtggggagat	cccggctcct	gtcttcccct	cgcctccagc	gcgctcgccc	aggctgggag	1740
gaggaaacca	gagccgcgcg	cagacacctc	ctccttctcc	tectettett	cctcctcctc	1800
ctcctcctcc	tectectett	cggctgctgc	tcctggtgcc	gccaccgtcc	gccggtgcct	1860
gttgctgccg	ccgccgcggg	acctgctgtg	tcctcagctg	ggtggagaag	aggegggege	1920
cgagccgagg	ggagccccct	ccccgtcccc	ccgcggcggg	aagagcgcag	ccagccgggt	1980
gcgatggact	ccccaccac	ccaggccgtc	cccaggatgc	ccccaagcac	ctgcgcgtcc	2040
cggcccggcc	ccgggctçtg	agcgcgccgc	ggcacaggta	aggcgcttcc	tggggcttcg	2100
tcctggccac	cctgctggct	cctctcgggg	cgtcgcggcc	gcccctccc	gcagcacgcc	2160
cctgccccgc	ctggccgcgg	aggggaaggc	atctggccgc	ccacggacgc	gaggccaggg	2220
tctctcgggg	gaggaagttc	attgccatct	cgttgccccc	cttacccccc	cacccccgcc	2280
gcccttggac	gaaagcgaaa	ccttaatgtt	gctagcgacc	cgagagctcc	geeggettet	2340
ccccaaccc	ccgccagctc	actggtccgc	gcatctctcc	cctccccct	cccgccaatt	2400
atcctagcgt	gtttgcaagg	cgaccagatt	ggaaagagtg	tggtcagagt	gaccccaagc	2460
cacgctttaa	aagttcaggg	tactttgcag	tagtaacttt	ggcagctcca	ccagtgcgcg	2520
caacatttct	ttctatgggt	acatcctgta	ccagtcattt	tgaaaccctg	cttcattgtt	2580
tctagccgct	tcctgatggc	tctgtgatta	tgagaccccc	ctcaaacttc	accaggcatt	2640
aaggttttgt	ttttgctttt	ttttcagaga	ggtatcattt	cgtttgaaat	ccacctagat	2700
gtggcttttc	ctgttttgat	tttacttaac	atagcttatt	ctctggaagt	tgctttaaaa	2760
agaaattgaa	agtgatggtt	gttccttcca	ccaaacagtt	taattttcag	ggtgcctcat	2820
attaatggat	atgttttccc	ttcatagatt	tctcattgtt	tcccttatga	tgg gatgatt	2880
tcatttatta	ataaaatcag	actttgaaag	agcatttaaa	aatgacctgg	tttaaatagg	2940
tcacacccaa	gaaactcagc	tatctgtaca	agttcaaact	tctaaacttt	ttcaatgagc	3000

taggggtggt ggcacccacc tgtagtccca gctacttggg aggctgaggc aggaggatca 3060 cttgagccca ggagttcgag gccatagtga gctatgactg tgccacctca ctggagcctg 3120 ggtgacaaag tgagatccca tctcttaaaa aaaaagagtt taggggacat tttctgaagt 3180 gaacacaagt agagcattct aacactattg agtgcaagga gacctggaag ggactaagtg 3240 gttcaaagca ggaaataaaa tcatcaggtg ataattaaaa taatttcttt cctgtgqatt 3300 tgtccagcca tttgcaaacc aggagaatag gaaaaaaaat cactagtgta gttataaatt 3360 attacattac gttttcaaag gaaaattttg caaatgcgtc tccttgtcat agtctattgt 3420 tatctacccc actgagagtg ctggggettc cccttttcac cacgacagca tttctggttg 3480 ggtggcagtc atgcagtgtt gacctggtgt cccataaggc acagtttgtc aaaacactag 3540 tgggtattag gaggaaacgt gcaactetga agcaacagag ettgeceett ettecteatt 3600 atccagctgg tgataatccc tgtcccccac ttccctagaa gacagctttg accaggaagg 3660 ctgcaatgac aatgagatgt accectatge agagecagat gtgggegggt ggettttttg 3720 tggtccagat cttctaggat cttctaggat gtaaccctgg caagcagtgg ggagcctgaa 3780 tcaagcagca tggctgttac ctcttctgtg ttcacagcag catcttcagt tgtcttggtg 3840 cctggagcag gcaccacagc tgcctgctct gttggccacc agctttctag agtagatggt 3900 agggaggaga gcaaggggct caagaggatt ctgtctttga acatgctttt aantttgatc 3960 tgacagaatg gcageteect gaagteette etaetetete cacageattt etetgtaggt 4020 ecccagtttt tgetetttte agatteecag aggaentgaa aatgtateae ggeecatttg 4080 gggacttect gtatatgtgt gggtgeetea ggateatttg ttttgeeett ttecagteta 4140 eegtgetgee etteteaagt ttaatgnace aegttagttt caatatttta tatatttete 4200 ageagtttte atetettggt cattaaaett gagaagtaaa atetgeteat taaaatgaet 4260 gagtccatgg ccaggcatgg tggctcatgc ctgtaatccc agcactttgg gagtccaagg 4320 cgggtggatc acttgaggtc aggagttcga gaccagcctg gccagcatgg caaaaccctg 4380 tetetacaaa aatatagate tacaaaaaet agecaggeat ggtggcatgt geetgtagte 4440 ccagctattt gggaggctga gacaggagaa tcgcttgaag ccaggaggcg gaggttgcag 4500 tgaaacatga tegtgecaet gagtecatte ageageagag tagtgttggg gtttgtatee 4560 ctgtagtgat gacgaaggat ttaggttttc agtcagaact gttaccttac aatttccttc 4620 actgactttt cttcctttcc aacaccacat tccaataaaa aatatcttta gaccagattc 4680 ttcacgaaag acatgaaggt tttcatgctt caaggttttt gactttttt tttttttna 4740 aaggagtett getgtgteae eeaggetgga gtgeagtgge gtgateteag etcaetgeaa 4800 ceteogeete etgggtteaa gtgattetee tgeeteagee teecaagtag etgggaetae 4860 aggegtgete taccaeggee ggetaaattt tgtgttttta gtagaggega ggtttcaeca 4920 tettggecag getggtettg aacteeegae ettgtgatee accegeettg geeteecaaa 4980 gtgctgggat tacaggtgtg agccacggcg cccgaccagt ttttgacatt tctaagccaa 5040 aagttccatt tgatgaggtc ttagatgcag gggcaatgtg tcccttttca gatttcagat 5100 gtttagaaaa agatgtgtca tatttgggcc aactgaaaaa ctcttgatat gtaggttttt 5160 atgaagctgt gcagaatgta ggaaatacat tttagaacca acaaagaggc atttaatttt 5220 gagtgtgcct gtctcctttg agatgagcaa cagctatttt tctcttcaaa agacaatgcg 5280 tgtatttatc agcacatttt atataatcag caaatctaaa cctctgaatt aggtaagccc 5340 tataggtttg ttgccagaat tagtgaattt atacatgcaa agtgcttaga acagtgcctg 5400 gtacacagtg agcactcaat attatttatt gctattatta tgtttattta ttttatactt 5460 ttagagtata attttgatgt taggtttgga ttgctgaggc caagcaaaat ttagatagac 5520

caacccagct	aatccactag	aaagatattt	gagggttatt	cccatctaaa	gatctatggg	5580
atctttggat	atcatctgtg	agaaacaaca	gaagtttgta	gataagacag	atatagattc	5640
aaatgccacc	ttcacaattt	attagtgagg	gaaccctttg	gtaaaatgag	catgacaaaa	5700
cctacttctc	agctttgtga	acgtacagaa	gatcatgaat	gtaaaatgtc	catgaagtgg	5760
taggtggtca	acagatattg	ctctagcaaa	gtggttaaga	gcaagcaaac	tctggagcca	5820
aagggcctgg	gttcaaatcc	cgattctgcc	acttcttata	gtatggcctt	gggcaggtga	5880
cctcactttt	ctgtgccact	attcaatgat	aatattcctt	tattgtccaa	cgttttgtaa	5940
gttaaatcag	ttaataaaca	cactatgata	atgtgttggt	aactattctt	tttactttta	6000
gcagaataac	ttaaaggaac	actgcaggta	ggagggttat	acataatctc	tgagggccag	6060
ctaggacact	cgcccatttc	ccaccctttt	tcctgtgcaa	tgaagagtat	aagaagtgac	6120
agtgccattc	taaaggacta	gccttgagtt	ggctctaatt	tatatgactc	gtgcctgtaa	6180
tcccagcact	ttgggaggcc	gaggctggca	gatcacnttg	aggtcaggag	ttggagacca	6240
gcctnggcaa	acatgacgaa	atctcatctc	taccaaaaat	acaaagatca	gccgggcatg	6300
gtagcacaca	cctgtaatcc	cagcttctgg	ggaggctgag	gcagaagaat	tgtttgaacc	6360
cagggggcgg	aggatgcagt	gagctgagat	cacgccactg	cactacagcc	tgggtgacag	6420
agtgagattc	tgtctcaaat	aataataata	atttatatga	gaaagaagtc	attcaaaagc	6480
atcattactt	tacatgtcaa	attagaaagg	cacaccccag	tactaaagca	tccttgatga	6540
tgaaaacatt	tagaaccact	ggtttcagga	gctccatgca	atggtgaaac	agcctctact	6600
ccaagggttg	ttgctccctg	tgggattcct	gggtgaggaa	cacactgctc	ccgttggggt	6660
ggaatcctgt	ggaggaagtg	gatgaagagt	gtagccaagt	cagtgagcct	actgcatggg	6720
attagagtac	ttcaggttgc	agtataattc	tgttcaggtg	catgctcact	ccatctggcg	6780
taaagaacag	agaaattaaa	ccattgattc	acagagcaat	atgagtagct	gcctggggac	6840
ttcctgtcca	ttctggggtn	cccaacagcc	aatcaatatt	ggccggttcc	taatctgacc	6900
tagagctaga	ggccactagc	acccctccat	tcatcctttc	ttctctccct	acccactccc	6960
accagcattc	tgaggaggtg	agggctgaag	ctgcagaggc	tgttgtactg	tcagttactg	7020
taaagtcccc	atcctgctct	ctgtagtttg	ctgtgaagga	gtggagggg	ctgggaacaa	7080
agggggttcc	caataagcag	cttactaata	cctacccttg	cttctctcac	ttcctgatca	7140
ataattccca	tectttette	atgtacctcc	cccacatttt	tgttctttaa	gggaagaagg	7200
gagcagcttt	ccatccccac	acaatgttgg	ggacatttgg	tattatacat	tatgaaaata	7260
agatttacca	gattttagag	atggaagaaa	acttggggat	gatcttgttc	cattctctta	7320
taagaacaaa	caatatttgg	agaagctgag	taacttgctt	tttcaacttc	acacttgaaa	7380
gtgttttcat	gaggaagttg	gggttctctg	cagcacttgg	atgggagtca	gggacttgga	7440
ttgtcccaat	tctgtcacta	aatttggaca	agccacttaa	cttttccaga	atctagttgc	7500
ctcatttcaa	atattaggga	aatttctaaa	tggcttaaaa	ggagcttgtt	agctttaaaa	7560
ttttatgatt	ctaagtgtat	gctgccagag	atatgtagca	tagcaggaca	cattaacaag	7620
attattgaaa	ctgttctaat	aaaggacatc	tttgtgtctt	gggtagctac	tatgtttaaa	7680
gactgtgcta	ggtgggagtt	gtgcagaata	cacaggtttg	ctgtagaggg	atagggcgtg	7740
tacacagaca	actctactaa	caagaacgtt	actagaagct	tattggaatc	acagtatttc	7800
ttgctgaggg	tatgaaacat	aagagttctc	cttggaatat	gaggttctat	ttggggctta	7860
aagaatggtc	aaaggttgag	tgcaaataac	atggattgag	atggctttaa	aaaataatca	7920
aatggtttgt	tagtattaaa	ctggtgcaga	aataattgca	gtttttgcca	ttccttttaa	7980
tggcaaaaat	tgcagttact	tttaaaccaa	atccctaata	ttatttgcat	agtttatctc	8040

_			•			•
tgttatggaa	gtttttattg	acaagtaatg	tagatattca	cctgatctaa	gttaccctga	8100
atcttatatt	agcagaatct	gaattgctta	taaataatta	tggctatgtt	ggatgtagaa	8160
cttattattt	gatagtttat	gaacagtgct	aaggtctaat	ctacttttta	cagagaagct	8220
aagaacatgc	tacagctggt	tgaaaaacaa	aaacttcagg	cattgaaatg	ttttgtcaat	8280
gaaatggcag	gactcatttg	atgactgatt	attatcaact	gatttaaatg	actgaatttt	8340
tggtactgtg	tacatctata	ctctaagaag	gaaattgaaa	gtaattctgc	tatgcttgtt	8400
gccactatat	taataactgc	atcatctaaa	ataattgata	gagctcagat	ttatcctttg	8460
taataattct	agtacttctt	taaacatgtt	ttgggattag	cagctgtcaa	cagttagaac	8520
atgaaacaga	ttctgttaca	ggagtagaag	tcgatccaga	catttaatgt	cattttcacc	8580
tgtgagagag	agaataaaga	gaaagagaga	tcattattta	tgggattatg	tgaacttcaa	8640
gtccgttttc	attattagga	'gaagctgtgc	t ttaaaggac	agtcagggac	tttactttca	8700
tgaaatgcct	gagctgtaaa	taaagtattg	ctttattttt	tatttcttga	acatttgaaa	8760
taaaaaatta	gctatgagtt	atgttcaaat	tatattataa	aaatttgctc	ttagcattgt	8820
gcatatatat	tatacagaaa	aacacagagt	aaaaagaata	gacttcagtt	cctgttcaga	8880
aaaggtttaa	aatttgaata	ctgattttgg	aaaccccaaa	ccttaagaat	tcaagaagct	8940
tacggtcttc	ttgagggaca	cctattcaaa	ctcttaaata	tggtgattgg	gtagaaagtg	9000
cagaaaagcc	tgctgataca	tgccctaaaa	caccttggaa	aaaagaggtg	gtagttgctt	9060
gaggtaggac	ttaagtacta	gttggaaata	gaagaca ag g	atggagactg	ttggtagatg	9120
actctccatg	ggtccttcct	gtttctacac	accttgtaag	cagggcattg	agtgcctgtg	9180
ttccaaacta	ccttttccat	catgtttcta	cagcaaacag	tcatggaaga	tagaaataga	9240
gtcttcctct	ggagcaaagg	gcagacacgc	ttgcttcctg	tacttcccac	tataagatat	9300
tccggctccc	taaactcagc	tgcctttcct	gtaacccacc	atgatacaga	tgtcacctga	9360
cctgtgggaa	ttgggggtca	gggaaccaag	agaaatgctg	actgtctggc	tactgtgact	9420
gccctgagta	ataaattgtc	cttcgtctcc	aacccaggag	tctcatgttt	tctac cagc a	9480
ggataactgt	ggcrggctaa	cgtgttagtt	tgcaagtaag	gtaaaatctc	agaccctttg	9540
			gagaggaacc	-		9600
gcatgagaaa	cggggaacća	taactagttc	tctatcttca	gagcctttaa	aaggtg cac c	9660
aaggagggca	ttttagggga	gaatataaag	ttggagatat	agacacagcc	agattcct g a	9720
gagaccttat	atgccaggta	gaagacttca	gattgtatgg	gggaattatt	agagaatttt	9780
tagcaggggt	gtgatatgat	aaattttgtg	ttgattaagt	tactccagga	aatatgcgat	9840
g ggtggat t g	aaggatgggg	caccttttct	ctaggacgaa	aaagaaagag	tagttggtga	9900
agtcagttag	aggtagtaat	aggatgaaga	agggatctga	atgacccctt	ggccattcag	9960
tgagtagtga	tgctattcac	ctagatacag	cacatagnng	ggaaangaaa	tnctgggaag	10020
gagggagatg	agaccgagtt	agctttaaaa	taactaaatt	caggcctagg	agcctatagg	10080
ctatccagat	agaaatattt	aatngcctat	atggatctgg	aactcaggaa	ggaggcttcc	10140
gtgggagcag	aacacttggg	caccattagg	gtgtatgtgg	tagatgcatt	cttgtgcagc	10200
agtcaagggg	atgggattta	gactcaagtg	caaattgccc	cccatctcct	gtgataagtg	10260
actgaagctc	tccgggcttc	agtttcctag	ttcatcatag	tgggctctag	cggataaatg	10320
ttacaaaggt	taaatgagac	aacataggca	aagtgcgtgg	tactcaatag	aagtcagctg	10380
ctgtcatcag	cagcaggatc	accagaatgt	ggtgcttgac	accaaaagat	taggtgagat	10440
			aggatgnaag		_	10500
ctttgaagta	tgtggaaaga	aacaaccaga	aaggtaagat	aagaaccaga	agagattcaa	10560

•			,			•
gaaggaaggt	gtggccgggc	gcggtggctc	aagcctgtaa	tcccagcact	ttgggaggcc	10620
gaggcgggcg	gaacacgagg	tcaggagatc	gagaccatcc	tggctaacac	ggtgaaaccc	10680
cgtctgtact	aaagatacaa	aagaattagc	cgggcgcggt	ggcaggcgcc	tgtagtccca	10740
gctactcggg	aggctgacgc	gggagaatgg	cgcgaacccg	ggaggcggag	cttgcagtga	10800
gccgagatcg	cgccactgca	cntcnagcct	gggcgacaga	gcgaggagcc	gtctcaaaaa	10860
aaaaagaaaa	aaaaaaaaa	gtaaggaagg	tgtggccaag	attgagaaat	tcgtcagagc	10920
aaacaaggca	gtcaggggct	aaatagcctc	ctttaaattt	tacaaccttg	aggacctcgg	10980
caactttaac	agaatttcag	tggatcccta	gggcaaacca	ggccttacaa	accaggaatg	11040
gatggtcaat	aggaagtgga	gacagtaagt	gtagacctta	ccttggaggg	aaggnaagag	11100
aaagagccat	ggccaaggga	agtttgaaat	caaaggaaat	atctttttt	tttttttcg	11160
attggagaga	cctcagttat	tcttttaaaa	tacttattga	gcccctcagt	tattctttta	11220
aaatacgtat	tgagtcccta	ctttgagtca	ggcacnatgg	cagacacgag	ggngatagca	11280
gtgaatcaga	cagatgcaac	gcctgccttc	atggagtttc	accttagcat	ctgtccatat	11340
gctaggggag	tggggcaggg	gcagggagct	ggatacagga	gagactgaag	atccagggag	11400
caagtgagta	aagaataggg	cttgagatcc	cacagacaac	tcagctttga	acaaaagggt	11460
tttgtcatcc	aataggacaa	gaaggcgtta	ggatacatca	aacgtggttg	ttgaaaacag	11520
aaaagggctg	ggcactgtgg	ctcatgccta	taatcccagc	actttgggag	gccaaggtgg	11580
gcagatcact	tgaggccagg	agttcgagac	cagcctggcc	aacatggtga	aaccccatct	11640
ctactaaaaa	tacaaaaatt	agccaggtgt	ggtggtgcat	gcctgtaatc	ccagctactt	11700
ggaaggctga	ggcaggagaa	ttgcttgaac	ccagggggtg	gaggttgcag	tgagccacga	11760
tcgtgccact	gcactccagc	ccgggcaaca	gagcgagact	ctgtctcaaa	aaaaaaaaa	11820
ggaagaaaga	acatagacag	ggaaatgtag	ttaaggnnag	tttgggtttg	ggtttggtag	11880
aagcgttttc	tgttnnttgt	ttgtttgttt	tcagaaagag	tctcactctg	ttgtccagac	11940
tggagtgcag	tggcacaatc	ttggcttgct	gcagcctctg	cctcctggat	tcaagcaatt	12000
ctcctgcctc	agcctcctga	gtagctggga	ttacagacac	ctaccaccac	accaggctaa	12060
tttttgtatt	tttagtagag	acggggtttc	accatgttgg	ccaggctggt	ctcaaactcc	12120
tgacctcagg	tgatccacct	atcttggcct	ctcaaagtgc	tgggattaca	ggtgtgagcc	12180
actgcacctg	gcctaacatt	gatatctgtt	gatgagaaga	agccaggtgt	tggagtgata	12240
gcttatagca	catgaactga	ataaaacagt	gtttaagaca	atgtttgcaa	cataataggc	12300
actgaagaca	tgttaatgga	aggtggattt	gtgattcaga	acctctagac	tacctgggcg	12360
agtcttttaa	aatgtaagta	atatcttaag	tgatattact	tgtcccagat	cagttgttta	12420
aaactgaggt	ttaatgctgt	cagagtagca	ctgtatcgtc	ttctatcatg	ggggcctttg	12480
ttggctttag	gaggtttgtg	tttcatagta	gtttcccagt	gggctctttg	ttacctgtaa	12540
tgagtgtgac	agttatgcca	taaccaggtt	ttatatggaa	tacaattttg	agaaagttct	12600
ttctaggcag	agaagcttat	ttgaacctct	tattatattt	gggtttcagg	cttttgagtt	12660
cttctgaaat						12720
ctctggtttt	ctccccagg	tttctgcata	tgaagtgtgt	aaaatagatt	gcttgatcca	12780
aaacagaaaa	acagtgataa	ctgttttgct	gagttcccag	accettecea	agatggaacc	12840
aataacattc						12900
gcagctggtg	ggctccctgc	ctgtgcattc	cctgaccacc	atgcccatgc	tgccctgggt	12960
tgtggctgag						13020
agtccggctt	tgcgtttcac	cctctggact	gagatgtgaa	cctgagccag	ggagaagtca	13080

•			O			-
acagtgggat	cccctgatct	attccagcat	ctttgagtgc	aagcctcagc	gtgttcacaa	13140
actgattcac	aacagtcatg	acccaagtta	ctttgcttgt	ctgattaagg	aagacgctgt	13200
ccaccggcag	agtatctgct	atgtgttcaa	agccgatgat	caaacaaaag	taagtgagat	13260
ggagatccaa	aagactaagg	tgtggctggc	tggtttttat	tgtatggggg	tcaggatatt	13320
tattttaagt	atactgaaat	gaataaggaa	ttaatgctgc	agttataaat	tgattactta	13380
gctgaatttt	tgttttatgg	tgatagttta	tagttttaaa	gcacatttga	aaacagatac	13440
gagaaattat	cagtttttga	gttcaaaaat	tcaagagaaa	tcagtctaaa	actactaatt	13500
aagagcagaa	gtgttaagat	gtacattatt	tcagatgaat	gttctaaagc	catgcctctc	13560
aaactgaaat	gagcttgtga	gtcacctggg	gatcttgtta	aaatgtgaat	cttgattcag	13620
taggtctggg	gtggacccca	agactgcatt	tgtaacaagc	tgccaagaaa	tgctgatgct	13680
gcccttttgc	aggttgcact	ttgagtggca	aagttctaaa	tctccacatt	tgtaatccta	13740
ttaaga aaaa	tatagtcatt	cgtaaactgt	gtaaaaatgc	tactggccag	tttcccaagg	13800
cataatgttc	acttaggcaa	aggtcattga	taagaacgct	ggatatgcat	ctaagttttg	13860
atgcgatcag	gggttctttg	tgttttttc	tttcgcaaac	ctcaggtcag	atctgattag	13920
cttgttatta	tcacatgata	tggctgaaaa	aaaatgtgag	acatggtaaa	agttctgctc	13980
tttcctcgtt	catttgtgct	tgctttgtta	ttagcattcg	ttgtagctct	gggcaggact	14040
catttgaaga	tgcttgnccc	attttatgag	gattagctta	gataaaattg	aaaatataat	14100
gcaaatagca	actttctcag	ttgggctcag	ggctccacag	ctaaccccat	ggactgtgga	14160
gtcttgccgt	tgttttgggt	gccaagcaag	ccaagtcaca	tgtgattcaa	gctgtctgcc	14220
acatgtacag	ggcgaggatg	cgagtgtcaa	tccacctgtt	aactgtcagt	gaagccttga	14280
aagcttctca	tattttcaag	gttaaaatct	ggatagaaat	gctaaagttt	tctctctgca	14340
ctccattagg	ttattttatg	tactctctag	ggtgtaagga	ccttatttag	aaattaatat	14400
tcttggtatc	aagtagatgc	ccttttgctt	gttcatttgt	tggttcttct	agtcattcag	14460
aattgctgtt	gcaggtactg	ttggagatga	tattagcaga	ggcttgtagg	aaggcaggag	14520
catcagtggg	gaataggacc	aggtgatcta	tgtataggac	ataatggaag	gactgagaag	14580
ggagcctaac	acacacccaa	agggtagaga	aggctttgtg	aaataaaggc	taatatggag	14640
ctcaaaacca	ccatttcact	cacagaatca	aactctcata	ttataaatca	tttcatgtta	14700
ttgtccacac	atctcaagtg	ggcacggcag	catcaggctt	ggagattcag	agggactaac	14760
ttcctgtact	ctaatcctac	ttctgcaccc	ataaactggg	tggcctcagg	caattgagtc	14820
tgttttctta	tctgttaaat	ggggataatt	acagtattta	tccaatagag	ttgctggaaa	14880
gactaaatga	ggtagcactc	gacctgaaac	ttagtaagca	tttatagcca	taaaaacatt	14940
ttcattcaag	aaaattttac	tagaggcaga	ttatatgcta	atttcatttc	acgtcttagg	15000
taaaaagaaa	catgatacct	agatgagtgc	cttcagcttt	caaagatgag	attctggtca	15060
tatttgagga	acattttaaa	aactacacgt	ataacttaat	ggctcctatt	atttggacaa	15120
attccagaat	gaaaatgaga	ggactgaaca	gcctgtacct	cagtccagct	ctatatagta	15180
tttggactga	atttccttgg	ggagagtttg	tgcgtggaat	cgttgttcag	cattttacac	15240
atttgactct	ttcccaaaat	cttttacggc	catctgagaa	taggcttctg	gccagtcatt	15300
cggatgcctg	acaagagaaa	gagatttata	accaaattct	gtaattggga	cttccagtct	15360
ttccccaagt	agagaattgg	acttactcta	tatgctaaaa	acccatggtt	gaaatatgaa	15420
ttagttctta	agtgatttt	ggcttgcata	ccatttttgc	aaacacaaat	tgtcattact	15480
ctgctcattt	aataaaagaa	taatttgtag	tataggtata	tacctcaatc	agtgattttg	15540
ttgttggaaa	cagaacagta	aatcacactg	gccatgatgc	taacagcgtg	atagattttc	15600

-			-			
tgttcttggg	acaccaatgt	cactgtatct	catagcgaag	gattatctgc	tgtaggagca	15660
ttctcttgac	tacttataac	atttgctggg	tgaaataatt	ctccaggtta	aggcctcttc	15720
taaacagatg	aggtcagcac	taactgcatt	tgccagagaa	gacatatgca	tttactgcca	15780
gcatcataaa	cacaaaacta	cagtttgcga	ggaaaccctt	tgaccagcat	ctaattaatt	15840
cactgagtaa	tgtcttggga	gaagaggcat	gtaaaggaac	aattttataa	gcatgccatg	15900
agattgtttt	cnnattgtat	gttccataga	atatgaggaa	acttcaaaac	attttgtgga	15960
aaaattgaat	taaaaagtaa	aaaacacata	tatacataag	ctttatttct	caagataaac	16020
tttatcaagt	tcaagacact	tttgtaagca	atgttaacag	ccattgagtc	ggtctctaaa	16080
gaactgaggg	tc ctgggaat	ttaaccatgt	ttatacagtc	ttttatacat	tattaactgg	16140
agaaaaattg	gcgctcttta	aagattttt	aaaattgaga	agcaaaagga	cgtcagaagg	16200
agccaaatta	ggcctgnnaa	gtggatgcct	aatgatttcc	catggaaact	cttgcaaaat	16260
tgctcctgtt	tgatgagagg	aatgagcagg	aacattgtca	tggtggacaa	ggac tctg gt	16320
gaagctttnn	caggcgattt	tctgctaaag	ctttggctaa	ctttctcaaa	acactctcat	16380
gataaacaga	tgttatcatt	ctttggccct	ccagaaagtc	aacaaacaaa	atgcct tg gg	16440
catcccaaaa	aactattgca	accatttgcc	cttgaccagt	ccactttcgc	tttgactgga	16500
ccacttctgc	tctcagtagc	cattgcttaa	atttgtcttg	atctttagga	ttgcgctggt	16560
aaaactatgt	ttcatcacct	gttacaattc	tttgaagaaa	tgcttcagga	tcttgatccc	16620
acccgtttaa	aatttccatt	agaaactctg	ctcttgtctg	cagctgatct	gagggcaatg	16680
gttttggcac	ccatctagta	aaacgtttgc	tcagtgttaa	tttttcatcc	aggattgtgt	16740
aagctgaacc	agcagagatg	tctatgatat	tggctagttg	gtcctcttca	atgagggcat	16800
gaacaagatg	aatattttcc	tcaaacaatt	atctggatgg	tctgctgctg	caggcttcat	16860
cttcaatatt	gtctcgtccc	ttettttet	tttcccccc	gcttgagaca	cagtcttgtt	16920
ctgttgccca	ggttggagtg	cagtggcccg	atttcggctc	actgcaacct	ctgcctcccg	16980
ggctcaagcg	attctcctgc	ctcagcccac	caagtagctg	ggattacagg	tacacatgat	17040
cgtgcctggc	taatttttgt	atttttagta	gagacagggt	ttcaccgtgt	tggccaggct	17100
ggtctcgaac	tectgacete	aagtaatcca	cctgccttgg	cctcccaaag	tgctgggatt	17160
ataaacatga	gccaccaeac	ctggcctcat	cctttcttaa	aatgagttat	acatttgtaa	17220
gctgctgatt	tctttggaca	ttgtgcctat	aaacttttg	taaagcatca	gtgatttcac	17280
cattcttcca	cccaaacttc	accataagtt	tgatgtttct	tcttgctttg	a ttttagcag	17340
gattcatgtt	tctctgatag	ggggtctttt	caaactgatg	tcttatcctt	cttagagcct	17400
catcccagat	cctgttcaga	catgctacaa	gttaatacaa	gtttatttgg	tgccaaaaaa	17460
	tgcatagttt					17520
ttgtatactt	aaactgctcc	acatggaaaa	gcttccatga	tcaaatgcag	taaggcagca	17580
tctcaaacat						17590

<210> 2

<211> 99960

<212> DNA

<213> Homo sapiens

<220>

<221> exon

- I-

```
<222> 4661..4789
```

<223> exon A

<220>

<221> exon

<222> 6116..6202

<223> exon B

<220>

<221> exon

<222> 9919..10199

<223> exon C

<220>

<221> exon

<222> 14521..14660

<223> exon D

<220>

<221> exon

<222> 50257..50442

<223> exon E

<220>

<221> exon

<222> 56256..56417

<223> exon F

<220>

<221> exon

<222> 63326..63484

<223> exon G

<220>

<221> exon

<222> 76036..76280

<223> exon H

<220>

<221> exon

<222> 78364..78523

<223> exon I

```
<220>
 <221> exon
 <222> 85295..85464
 <223> exon J
 <220>
<221> exon
<222> 93417..93590
<223> exon K
<220>
<221> exon
<222> 97476..97960
<223> exon L
<220>
<221> misc_feature
<222> 97961..99960
<223> 3' regulatory region
<220>
<221> allele
<222> 1443
<223> 99-20508-456 : polymorphic base C or \mathtt{T}
<220>
                  1 -
<221> allele
<222> 5247
<223> 99-20469-213 : polymorphic base C or T
<220>
<221> allele
<222> 6223
<223> 5-254-227 : polymorphic base A or G
<220>
<221> allele
<222> 14723
<223> 5-257-353 : polymorphic base C or T
 <220>
```

```
<221> allele
<222> 19186
<223> 99-20511-32 : polymorphic base C or T
<220>
<221> allele
<222> 18997
<223> 99-20511-221 : polymorphic base A or G
<220>
<221> allele
<222> 19891
<223> 99-20510-115 : deletion of TCT
<220>
<221> allele
<222> 29617
<223> 99-20504-90 : polymorphic base A or G
<220>
<221> allele
<222> 42519
<223> 99-20493-238 : polymorphic base A or C
<220>
<221> allele 、
<222> 69324
<223> 99-20499-221 : polymorphic base A or G
<220>
<221> allele
<222> 69181
<223> 99-20499-364 : polymorphic base A or T
<220>
<221> allele
<222> 69146
<223> 99-20499-399 : polymorphic base A or G
<220>
<221> allele
<222> 76458
```

```
13
<223> 99-20473-138 : deletion of TAACA
<220>
<221> allele
<222> 78595
<223> 5-249-304 : polymorphic base A or G
<220>
<221> allele
<222> 82159
<223> 99-20485-269 : polymorphic base A or G
<220>
<221> allele
<222> 84522
<223> 99-20481-131 : polymorphic base G or C
<220>
<221> allele
<222> 84810
<223> 99-20481-419 : polymorphic base A or T
<220>
<221> allele
<222> 89967
<223> 99-20480-233 : polymorphic base A or G
<220>
<221> primer_bind
<222> 988..1006
<223> 99-20508.pu
<220>
<221> primer_bind
<222> 1509..1529
<223> 99-20508.rp complement
<220>
<221> primer_bind
<222> 5039..5056
```

<223> 99-20469.pu

<220>

<221> primer_bind

```
<220>
<221> primer_bind
<222> 5534..5554
<223> 99-20469.rp complement
<220>
<221> primer_bind
<222> 5997..6015
<223> 5-254.pu
<220>
<221> primer_bind
<222> 6332..6350
<223> 5-254.rp complement
<220>
<221> primer_bind
<222> 14371..14390
<223> 5-257.pu
<220>
<221> primer_bind
<222> 14798..14817
<223> 5-257.rp complement
<220>
<221> primer_bind
<222> 18751..18771
<223> 99-20511.rp
<220>
<221> primer_bind
<222> 19198..19217
<223> 99-20511.pu complement
<220>
<221> primer_bind
<222> 19605..19625
<223> 99-20510.rp
```

Ę. :-

```
<222> 19986..20005
<223> 99-20510.pu complement
<220>
<221> primer_bind
<222> 29529..29547
<223> 99-20504.pu
<220>
<221> primer_bind
<222> 30041..30061
<223> 99-20504.rp complement
<220>
<221> primer_bind
<222> 42268..42287
<223> 99-20493.rp
<220>
<221> primer_bind
<222> 42732..42752
<223> 99-20493.pu complement
<220>
<221> primer_bind
<222> 69026..69046
<223> 99-20499.rp
<220>
<221> primer bind
<222> 69525..69543
<223> 99-20499.pu complement
<220>
<221> primer_bind
<222> 76323..76343
<223> 99-20473.pu
<220>
<221> primer_bind
<222> 76771..76790
<223> 99-20473.rp complement
```

<220>

```
<220>
<221> primer_bind
<222> 78292..78309
<223> 5-249.pu
<220>
<221> primer_bind
<222> 78704..78721
<223> 5-249.rp complement
<220>
<221> primer_bind
<222> 81893..81912
<223> 99-20485.pu
<220>
<221> primer_bind
<222> 82353..82372
<223> 99-20485.rp complement
<220>
<221> primer_bind
<222> 84392..84412
<223> 99-20481.pu
<220>
<221> primer_bind
<222> 84909..84929
<223> 99-20481.rp complement
<220>
<221> primer_bind
<222> 89746..89765
<223> 99-20480.rp
<220>
<221> primer_bind
<222> 90179..90198
<223> 99-20480.pu complement
```

```
<221> primer_bind
```

<222> 9475..9493

<223> 99-430-352.mis

<220>

<221> primer_bind

<222> 9495..9513

<223> 99-430-352.mis complement

<220>

<221> primer_bind

<222> 1431..1455

<223> 99-20508-456.probe

<220>

<221> primer_bind

<222> 5235..5259

<223> 99-20469-213.probe

<220>

<221> primer bind

<222> 6211..6235

<223> 5-254-227.probe

<220>

<221> primer_bind

<222> 14711..14735

<223> 5-257-353.probe

<220>

<221> primer bind

<222> 19174..19198

<223> 99-20511-32.probe

<220>

<221> primer_bind

<222> 18985..19009

<223> 99-20511-221.probe

<220>

<221> primer_bind

<222> 29605..29629

```
<223> 99-20504-90.probe
```

<220>

<221> primer_bind

<222> 42507..42531

<223> 99-20493-238.probe

<220>

<221> primer_bind

<222> 69312..69336

<223> 99-20499-221.probe

<220>

<221> primer_bind

<222> 69169..69193

<223> 99-20499-364.probe

<220>

<221> primer_bind

<222> 69134..69158

<223> 99-20499-399.probe

<220>

<221> primer_bind

<222> 78583..78607

<223> 5-249-304.probe

<220>

<221> primer_bind

<222> 82147..82171

<223> 99-20485-269.probe

1 -

<220>

<221> primer_bind

<222> 84510..84534

<223> 99-20481-131.probe

<220>

<221> primer_bind

<222> 84798..84822

<223> 99-20481-419.probe

PCT/IB99/01444

WO 00/08209 19 <220> <221> primer_bind <222> 89955..89979 <223> 99-20480-233.probe <220> <221> misc feature <222> 3698,12593,13035,21712,27644,27655,31143,43084,43129,64585,66950 67301..67302,67926,75425,98821..98822 <223> n=a, g, c or t <400> 2 ctcaagcttg aatacttgaa tccaaacttt catgcttaga gtttacccca tctgttgaag 60 gatgtgcaat ataatgactg caatagaatt cactgtggag cctccaaatt agaaattatt 120 gtctgtgagg gccaggcacg gtggctcacg cctgtaatcc tagcactttg ggaggctgag 180 atgggaggat tgtttgaggc caggagtttg agaccagctt ggtcaatata gcgagacccc 240 catctctgtt tttttttt aaagaaatta ttgtctaaga accagtgtca tcttccaagg 300 agaaacttct agatacttgt tttaagataa ataagaaaca agtcatttct aaatgtgaat 360 tattttttaa atgcaatttt ttaaacattt tattttaatt atggcaatag acgtggaaaa 420 gactettttt tgatagtagg ggagageaga agaaacattg aattaagtae acagagatte 480 ttcagacctg ctttaaaaac acatgcatac aaatgcactt ctgtctctta ggatctacta 540 actgatgctg cttgctttag tcttttagct aatattttct ttctttcttt ctttctttt 600 tgttggagac agagtctcgc tctgtcgcca ggctagagtg cagcggcaca atcttggctc 660 actgcaacct ccgcctcccg ggttcaagcg attctcctgc ctcagcctcc tgagtagctq 720 ggactatagg cgtgcgccac cacgcccagc taatttttgt atatttagta gagacggggt 780 ttcactgtgt tggatgggat gttctccgtc tcttgtcctc gtgatccgcc tgccttggcc 840 tectaaagtg etgggattae aggegtgage caetgegeet ggeteatatt ttetttatat 900 atcaaaacaa ttcagcttgc ttcactttta tgaaagcttt attatgagtt tgaaagcaat 960 tctgcatttt cttaacattg taactggtgt tgagttgaag gcaggcccct gggagccctt 1020 tgtgggcaat tcccttcact ctggaggctg cctcgagcct ggacaggcac ttacacttgg 1080 tcagtgattg cacagaaccg gttgcaacag attctgtgca cctccctgtg gcgcgtagca 1140 tttagcagge acttggtcac tatttgctga gtgagtctgt taccttaggc gtgtatttcc 1200

cgtggacctg cctggggatc attgctcatt cactcatttt gaacaagcca atattacatg 1260 tccagggtac-gctctatagt gtgaaacaca aaggtaaatg atagttcccc ttctcaaagg 1320 aatttctaag gtagtagcca ttcttttgat gcatattctc attctcatag agagtccaat 1380 tatggataat tggacaaagc tgaatgtcgc ttttatgaga atccattctt tctcttttat 1440 gcyttgaaaa atgtgtagca ttcattagtg aattaggatt tcattattca aagaagacat 1500 aaggtcttcg aacagcagat gactgaataa aataatacct aacagcagta gaatgagggg 1560

aggacatatt caaggaacat tttatgccca ttagattggc agaaattttt aaaaagtgac 1620 aataccgtat aaaggtgaac tttcctatac tgatactggg aacatgaatt tgtaccattc 1680

agggaagaga aacttgataa tatctggtgt agtctgaagg ggcacagtcc ctgtgaccca 1740 gtgaggacat tecteattat tteeettgee aaacatttea catgagteta taaggagete

1800

-						•
tatataagag	aggtcactgc	agcctccttt	gtaagagcaa	gaaaaaaaag	caaataagtg	1860
tttaacaata	ggaacataga	taaattaggt	tatgcagtga	atatttgcac	tctgactaaa	1920
gtgagtgaat	caaaaaaaat	ttgtcaacag	gaataaatct	caaaaataat	attgaaagaa	1980
gaaagctaat	ttacagaagg	atgtgtacag	tatgacacca	ttcatttagt	ttcaactaca	2040
tatcttttat	ggacacatac	atataaaagc	agaaaacatg	aattgatagg	ataaacacca	2100
aatatttctg	catatggcca	ggtgtgggga	agtagtggtg	attaagcttc	aaagatgtct	2160
gcagtggttc	ccattaaaag	tagaaagtag	gctgggcaca	gtggctcacg	cctgtaatcc	2220
cagcactttg	ggaggccaag	gcaggtggat	catttaaggc	caggagttcg	agaacagcct	2280
ggtcaacatg	gcgaaacccc	atctctacaa	aaaaaataca	aaaattagcc	agatgtggtg	2340
gcgcacactt	gtagtcccag	ctactcggga	ggctgaggca	tgagaatcac	ttgagcccag	2400
gaggtagagg	ttgcagtaag	ccaagatcgt	accactgcac	tccagcctgg	gtgacagagt	2460
gagactccat	cccaaaaaac	aagcaaacaa	aaaaagctca	tagagtaggt	aatagtcatg	2520
atatctgatg	ttttttgatt	gtctggttta	cattttttat	ttttatttt	tgagacaagt	2580
ctcacgctgt	cacccaagct	ggagtgcggt	ggtgcgatgt	cagctcactg	caatctctgc	2640
ctcctgggtt	cgagcgattc	tcctgcctca	gcctcccaag	tagctgggat	tacaggcgtg	2700
caccaccaca	cctggctaat	ttttatattt	ttaatagaga	cagggtttca	ccatgttggc	2760
caggctggtc	tcgaactcct	gacctcaagt	gattcatctg	cctcagcctc	ccaaagttct	2820
gggattacag	gcatcagcca	ctgcacctgg	ccttggtata	tgtgttttaa	tttgtattca	2880
ttcatttaag	cctcatgaca	gctctgcgag	gaaagttcac	tatacgtctt	caggctgcag	2940
gtagaggacc	tgaaagggac	aggaggtaac	agtctggcca	agaccacaga	gccagggaat	3000
agcagaggaa	catttcacct	gggcattgca	ctccagagct	gggcttctca	ctgttctcaa	3060
cccctggcaa	atgctcactt	gaacaaagcc	aggtggtgat	acaaaggtat	ttgttatatt	3120
agtctctaca	cttttctgtg	tgcttgaaat	aactgcaaca	aagaatatat	cagtatttag	3180
agtaatgggg	gatttgcttg	tgtgtgtttg	tatttttgag	atggagtctc	gctctgtcgc	3240
ccaggctgga	gtgcagtagc	atgatcttgg	ctcactgcaa	cctccggctt	ctgagttcaa	3300
gcgattctcc	tgcctcagcc	tcctgagtaa	ctgggattac	aggtgtgcgc	cactacaccc	3360
ggctgttttt	tgtattttta	gtagagacag	ggtttccccg	tgttggccag	gctgatctca	3420
aactcccgac	ctcaggtggt	ccacccacct	tggcctccca	aagtgctgag	attacaggca	3480
tgagccactg	cgcctggccg	tttttttc	taacaaaatt	attttctaac	agaaagcaat	3540
caggtgagaa	tccacataag	aaacaattta	attcagagat	ttttgttgca	tattaaaaaa	3600
aaaatgtacc	ttcggctggg	tgtggtagct	cactcctgta	atcccagcac	tttgggaggc	3660
tgaggcaggt	agatcacttg	agctcaggag	tttgaganca	gcctggccaa	catggtgaaa	3720
ccccgtctct	acaaaaacta	caaaaaatta	gctgtgtgta	gtcccagcta	ctctgggggc	3780
cgagggagaa	ggattgcttg	aacctgggag	gtcaagactg	cagtgagcca	tgattgtggc	3840
cctgtactcc	agcctgggca	acaaagtgag	accctggcac	cctgtctcaa	aaaaaaaaa	3900
aaagtacctc	cttgtaaata	agtaacacta	agacttcatt	tagtggttgt	caagcaaact	3960
ccattgtatt	tttattttca	gtttttatgg	ctagtagtta	agggagagaa	gcttggttgc	4020
				agaggaaaac	_	4080
				agagaatagg		4140
				ggtaagttat		4200
				caatgtttta		4260
ctgtttgagt	ctcacaacac	agtgttatga	ttcgatcttg	ccattggtct	cactttactg	4320

The first from most find find in

aagaggaagt	ttgaggctca	gaaaagtaag	aaactggccg	aagaccacgg	ttagtgaaga	4380
	atccagttgc					4440
agcacatttc	gtcattttac	ttggggtaat	caaagcaact	ctctgaggca	aaattatttc	4500
ctggacttgc	agccatgtca	ctaaggagca	gatgaggtga	gatcacagac	aggatcagaa	4560
tgatggcctg	gtgccaaaaa	gatgtgtcct	agagatttt	cattccttta	agaagcagag	4620
aagggagcga	taaatgactt	ttcgtttttc	acttttttag	acatcgcaga	tggcagcaga	4680
gaatattgga	agtgaattac	cacccagtgc	cactcgattt	aggctagata	tgctgaaaaa	4740
caaagcaaag	agatctttaa	cagagtcttt	agaaagtatt	ttgtcccggg	taagtagcat	4800
aatttctcct	gatttaagtt	aaatcacttt	ttaggagagt	gtaagattga	gttctatgct	4860
tttattccat	caatgttcat	cataaaggta	aaagtataaa	acctttttt	atgttttctc	4920
aggcttataa	cagtattatc	tacattttaa	attgttttta	atttggccta	ggtttaaaaa	4980
aaatattcct	tactcttttg	tattatatcc	aatgggattt	ttttgccgct	ccaaagaata	5040
tttgttagcc	agtccctata	aagagcatgc	attagataca	ctgaagtgtg	gcttctgttc	5100
tccctactat	cactatgtat	aacttaaaaa	acagttactg	tcagctgctg	gtgttagcta	5160
tctaaaaggc	tatatagtag	gggtcagcaa	actatgccca	tgggccaaat	tctacccacc	5220
tcctatttt	gtaaataaag	ttttgtygaa	acaccgccac	atccattcat	tttccagtta	5280
tctaaggctt	cttttttgca	gacttcagca	gttgccacaa	acactatatg	cctcacaaag	5340
cataagacac	ttactatctg	gccctttaca	gaaaaagttt	gccaaatata	gctctataga	5400
aagaacaaag	tacacatgta	catcaatctg	ggagttcttt	aagaaattat	ccctccctcc	5460
catgagtgta	aatagcctga	tggcacgtct	gagaaatcaa	atctgatttt	ccctcagagt	5520
ttcacacctt	tctggagtgt	gcagtatctt	attatagttc	tttttgattt	tatggcacac	5580
ttcttttgaa	acatctgatt	ttattttatt	ttttaattaa	ggaaagttaa	attttatttt	5640
cttcgaagat	gtttctgaga	attttgcaat	atcttctgag	atcatgaaaa	acagttgatt	5700
tacaaaacca	gagttgggag.	gggctgcatt	tgagagctcc	caaagggata	gagtgctgtc	5760
cgagtgacat	gcggctggcc	gttatgatga	cttgtgaccc	aggggaggga	gttagttgct	5820
gagtgggctt	gagcacttga	ttttccttat	agacgaattg	tcttgtcttc	ctgcctatca	5880
ctcatgccaa	attacttagc	caccaggtgt	tttggaacgt	ttaggttagt	gtcttattta	5940
ttttttaaaa	aaatgatgga	aatgttgatt	attttaatgt	acaaatatcc	ttagtagcat	6000
ttctcagtag	ataacatttt	tttcctgagc	ttatttaaat	ggaccaatct	gcttctagct	6060
gatgcctttg	caaaagcctc	cagagtcata	actcgactgc	cttttcttta	tgtagggtaa	6120
taaagccaga	ggcctgcagg	aacactccat	cagtgtggat	ctggatagct	ccctgtctag	6180
tacattaagt	aacaccagca	aagtaagcac	atttctcttt	atrogacaco	ctgaagaaac	6240
caacaaatag	gtcttgctca	teteetgtet	acatacctcc	aatcataaaa	cgtttgctgc	6300
ttgcaaattt 1	cttggcacag	gtggaggact	ggtcatgcag	ttctatcata	acataaaagt	6360
tttacataaa	agagcagatg	gggctgggtg	cagtggctca	acgcctgtaa	tctcagcact	6420
ttgagaggcg	gacgcgggcg	gatcatgagg	tcaggagatc	gagacgctcc	tggctagcac	6480
agtgaaaccc	cgtctctact	aaaaatacaa	aaaattaaaa	attagccggg	cgtggtggcg	6540
ggcacccgta	gtcccagcta	ctcgggaggc	tgaggcagga	gaatggcatg	aacctgggag	6600
gcggagcttg	cagtgagcca	agatcacgcc	actgcactcc	agcctgtgtg	acagagagag	6660
actctgtgta	aaaaaaaaa	agcagtagat	tttcctatta	aaaaaataat	taatattggg	6720
aaaacatcag	aaagtggatt	tgtgaattta	gagaagtata	cagcttaaat	ttttctttt	6780
ttaagaaaat	tttattttgg	atttgggggt	acatgtgcat	gtttattacc	tgggtatatt	6840

gcatactggt ggggattggg cttctagtgt acccatcacc caaatagtga acattgtacc 6900 cagtaggtaa tttttcaacc ttcacaccc ctttcattct cccccacttg tggggaaatt 6960 aaatttctga aactttatcc tgtagctggc tctatgatta taatgaaaca ttactgtttt 7020 atttaaataa gcaagtatct atgtccttct tttaataact tgctttctag acatttaatc 7080 atatttaagc ctggtcagtt caactttata actcctgaaa agtgggtttg ggttttgtgc 7140 tagggaggcc agctttccct tctgctacca gaggactctc tttggcagta gtgagggagg 7200 gagtgtttgt ggaggccagc tccttaccac aggcagggtt tacagtcctc tqccatccct 7260 cctagacata tggctttcag aatttttcta acctacagta agaagcacat ttaacattgt 7320 ggcgtagttc acaaacacac atacctacac attcacacac aaaattaaaa gttcacaaaa 7380 caatatttac tgtgaacaac atacaataca tactgatatt ttgttctatt ttattttaa 7440 aatgctcatg gcaaactact cagttgtacc acctactaac atgatagagg gagcagtttg 7500 agaaacactt ccttagatgg atgagtgctt ctcaaatttc aggtgctccg cctcccgggt 7560 tcaggccatt ctcttgcctc agcctcctga gtagctgaga ctggttaaag tgcagattct 7620 ggttcagtag gcagggcggg gggagccctg aaatgctgca tttctgacaa gctccaaggc 7680 aatgctgctg ctcctggtct gcagaccgcc tctggggagt gaggtcctag acagcagtct 7740 tgtaaatgtg agtttctgag ttaaaatcca ggggaacata gtgtcgtcca gcctccatct 7800 aatacacact gatcccaccc tgcaattcat tgcaagtgtg ggaaggctat ttgcttattt 7860 gttgtgtaca gatgaaccac acaccgcccc tttcatgtag gaagttacct aggaggagag 7920 agatgacaga tacagaaaca gccccagcat caagcagagt gtggtaggag cccagaaqtt 7980 acaaataaga gacattggta acttcagtgt cagaagagca aggggaaggg aagttaggtt 8040 tggtcagtgg aaccagggaa aaggtgaggg gtgaggggc agtgatgctt ttggactaaa 8100 tcttggtgta ggaattgtgc ctatggaagt gaacagagaa gaaggcattc tagacagacc 8160 agtgtcaata gacataccat gaagacattc atgtcactca gtgggctttc cagtaagcct 8220 tattgcttgc tttttatttt tttccaaaag gcagatctag gaatatatac atattcattc 8280 ttcaggactc gatagttgtg aagattettt taaaaggatt taaaagtetg tetaagattg 8340 caatttctag agtcattcta agagagatgc aacttttcag aagctgcttg tatgtattgt 8400 atatgtttaa gtgtacttta catctttctt ttattcatct tgaattgaga aactactata 8460 ttctatttta tgtaattgga tcccttctaa aaaattgatc acctaggagt tgcaaagaaa 8520 ccaaatagcc ctgaaacttg acaaatgaaa atggcccttt cagttgtcca attaagctaa 8580 gggttagctc tttgatatga tttggaggga tattagtaag aatttagatc aacaggtttg 8640 catgatggag attgtgttct gtgatgtatt gtcttagaga gacttttaaa tccttaaaaq 8700 aatcttcaca actgttgagt cctgaagaat gaaaaacttc agttatgaaa gtaatcaata 8760 tttcatagta tgttgggaat tttttcctaa ttcttataca attaaatgta tgtaacttct 8820 ccctttggta aacacatttc ttttttttt tttttcaaat taaaaccctc aatacttgtt 8880 acctaaaagg cactcaactg tgtaaatgaa caggtagaat tcagagtctc cagtccactg 8940 ttagatgcat tcattcttgt ttactctatt cctgttgatt tattttttct cttccaacaa 9000 tttcaatagg agcaagctgc tacaattcct ctttttgaat attttgaata tattaaaaat 9060 atattggcca ctagccacgt cctgggtgca gtgttaaaca tcagtttqct tgaqtqqtaq 9120 tagttcattc ctttgaaaaa gcgtgcatcg tgaaggcata caactttaaa atattgtcat 9180 gattctcaac aaatgtttga gcactcactc catagattta ttgcatacct aataaacaa 9240 taacttatgt ttgtgtaaca ttttacaaca taaaaagtac ttttggttgt atcatcttgc 9300 tttgttcttg aaactcagat acatttttac tttaccctct tacagaagaa attgaggtgc 9360

-			23			•
agaaagaaat	tatttgccct	gaattgcagc	agtaagtgcc	tacagagtga	ttttccatat	9420
tctaagaata	ttgatacagt	tcttaatctc	aaattatgaa	gtcgaatctc	aacagtagat	9480
cagattegga	gagagcctta	aaatgtgggt	ttaacatgag	tgaacacatg	tggcaaagat	9540
aaagaacttg	gttaagcagt	ggagacaagt	tctccagcac	tcacacccct	tagaagctgc	9600
agtaaacagt	cctgttttct	agagagaggg	cactattcat	ggcgttgttc	agaacgttac	9660
agattgtggc	ttatgtcctt	cactcctgca	cttggccagt	ctccccattt	ctccagcaag	9720
ccagcagtgt	gtccttgagg	agcgggcatt	tatttaatgg	accttcattt	tcttctgctt	9780
ttggtggtgg	cttctagatg	gcattataat	cagaacacat	acttagatac	tgcaatgttt	9840
gcccgtgcag	gaactagaga	tttataaatc	ccacatattc	cccatggtgt	gtctgatctg	9900
ctgtgtgttt	gctcccagga	gccatctgtg	tgtgaaaagg	aggccttgcc	catctctgag	9960
agctccttta	agctcctcgg	ctcctcggag	gacctgtcca	gtgactcgga	gagtcatctc	10020
ccagaagagc	cagctccgct	gtcgccccag	caggccttca	ggaggcgagc	aaacaccctg	10080
agtcacttcc	ccatcgaatg	ccaggaacct	ccacaacctg	cccgggggtc	cccgggggtt	10140
tcgcaaagga	aacttatgag	gtatcactca	gtgagcacag	agacgcctca	tgaacgaaag	10200
taagatttgt	ttaaatttgt	tgcataaata	gctggggcat	atctgtgact	agccaggtat	10260
gtgcatccca	ggtatgttta	ttgagtgaga	gaaatgagtc	aggctttact	cttggtttgg	10320
agataaaact	ggaagcagtg	acatgttcgt	tcgagctgct	tgtgagtata	caagcaatgg	10380
gtacttgtat	tgtcaggaag	caagtgaaag	tgagcaaaaa	tggtacctaa	catgcatagt	10440
cattactcct	caaacaaagt	aagagacgtt	gttgactgtg	gaactttgct	gctgtgagga	10500
agagggcaag	cggatgagtc	tccccatctg	aagccctgga	gcagggttat	aatgggaggg	10560
agaggcgctg	atccttacag	gcagagcaag	agaggtatgc	tggcctcata	gggtgacagg	10620
ggtgcttcag	cttctggtcc	tagctctgcc	gtgaactaat	tgtgacctgg	acgaatttgc	10680
taaattctct	gaataacaaa	attggagtag	atgttttcta	aaatctctca	ctgtaagaat	10740
tctagattct	tctaacaaga	tttattcatt	gtaatagttg	ggttcctgtg	accagttaga	10800
atcgtctggt	tatggagaag	agtaatcaga	agttcccccc	attccttcca	agtgtccctt	10860
agtgattcat	ttaattctgt	gtgccagaga	ctataaatgg	acacagttat	ctttaaaaac	10920
	, -		taatatgaat	-		10980
gtcgctgcct	tcttctgacc	agcagccgca	gaagtcccag	gacctatgtg	ttcgtgtagt	11040
tcatacacgg						11100
agaagtgtga						11160
tcagccattg						11220
agcaggggac						11280
ttgggttgta						11340
ctgectcace*						11400
ttcctgtctg						11460
gataggactt						11520
gatgtcacac					_	11580
tggcccattc						11640
agacattgaa					_	11700
tcagttttt						11760
gaaaaagaaa						11820
tcccagagct	ttgggagact	gaggtgggag	gttgcttgag	gccaagagtt	caaggtcaca	11880

PCT/IB99/01444

gtcagctgtg atcgtgctac cgcactccag cctgggtgat agagtgagac ctattaaaaa 11940 aaaaaagtat tgttgggagc ataaacacgt gggaaatggt caagaacggc cgtcaatata 12000 ctctgttttt cactgaaaac tacctttgcc agagagcgag cagagatgag gaaaaggagt 12060 ggaagaagtc ctccactctg atagtgttac tggaacaacg agacaaaagc ggtgtgctcc 12120 ttccacctgt ttgctccgtg tccctgtcgg cgccccctct cctgctaacc ccccgtgct 12180 ttctctgatt gctgtttagt gtggatcctt cacctgtggg tgagtctaag caccgcccag 12240 gtcagtcttc agetcetget cetecacete gtettaacee etcegeetee tegecaaact 12300 tttttaagta cctaaaacat aattccagtg gagaacaaag tgggaatgct gtgccaaaga 12360 ggtgagcaca ctcacgtggc aagtttggtg ttgtctgttt tcctggggag ttcacactga 12420 tgaggatgtg ctgaatgggg ggaatgtcca tgcaggaagc agagccactg tgtgtgtgt 12480 tgtgtgtgtg tgtgtgtg tgtgtgcgcg cgcgcgcgtg tgtctttgtt tatattttgt 12540 cttattttca gctgtcattt gaaccaagtt aattttacta ttgatgactt ttnttaagat 12600 tattatgaaa acagatetta atggcagatt ggtttgtgtt tgtgtttgtt ttttttttt 12660 ttgagacagg gtctcactct gttccccagg ctggagtgca gtggcgtgat ctcggctcac 12720 tgcagcttct gccttgtggg ttcaagcagt tctcctgcct cagcctcccg agtagctggg 12780 actacaggca cacgccacca tgcccggcta atatttttat ttttttttgt agagatgggg 12840 tttcaccatg ttgaccaggt tgttcttgaa ctcctaacct caagtgatcc gcctgcctca 12900 gtctcccaaa gtgctgggat tacaggtgtg agccactgca ccctgctgca aattgttttt 12960 ttatacttat tttcacattt ccttgcccta gtggacactt acatgcatgc gtatatacac 13020 13080 cacacaggat aacatetgtg tttgateatg tacactgeaa tttgtgeeat ateagaaact 13140 tcctgattga tttaggggaa ttattttcc cagtttgaaa ggaagagtta tttggaaaat 13200 ggatggattt tctttttaa aaaattattg atcccattca tttaaaatca aattttattg 13260 gtgaaaatga aaattaaatc tcgttcgtga actactttta atttcttacc tagttttctt 13320 ttcttagcat tagaacaaaa atgtttcttt tattttgaag cttatatttt atactttgtg 13380 tttttatgtt tctttatcct aaactctttt ttcaaccaaa ctcttagcat ctcctactgt 13440 aatgccctgc ggaaaaaact tcattcttct tcctctgtgc caaattttct aaaatttctg 13500 gctcctgtag atgaaaataa cacctctgat tttatgaaca caaaaaggta gggcttaatt 13560 tagatatatc aagcctgggt gttactaagt gttgaatatc attagatata caagggtgtt 13620 ttaattacta ttttgccatt taaaaaatca tttcagctaa atctgttgta tcttcttct 13680 tatacttttt tcttactgaa tgccattttt aaaaatgtgc aaccaacctg ttctctagtt 13740 ttgacgagga ttagtttaag tgttgtctta agaaaagtct ttgccaagtc tctgagacca 13800 gtgtttctgg ttagtgagca tatgtctgtt tcaaatcagg atgtctgatc tgttcaggac 13860 gtctaatctg-taagttgagg ggattgctta cttacaggta cataacttgg gtataaattg 13920 gaagggcctt ctcaggttgt cctgtgaata ggagaaaaca tttatgattg tgtttatata 13980 ttgataactg tattttgtag tttaaaaaat acacacgtta aaacaattat catcatcaag 14040 tgactgcata gttattgcct tgctggttct gtgtaattaa attgcaagtt ttttcatttt 14100 ttgtgggaat cettggagac atgggeetgt getgageaga tatteceatg cacagaagag 14160 ggcagaatgg ggccccttgg catcaccccc tttccccctt taggcagttt ctctttatca 14220 aagtggcacc aagagaggcc caattggaac tatgatatgt ggaacatgtt tcttaatctc 14280 tgttacaatc gaaatcactt aagggcatgt aatctttctc ttttcatgaa aagaattctg 14340 taagaaagca gttetttagg aatgatgaee caetgtgage ttgatataae ttetgtgatt 14400

gattatttgt	ttatacaaag	atagttgata	atttagtgat	ttgtttaaaa	aaatgttaag	14460
ctaacaaaat	cccgtgaatt	cctccccact	agtcataaat	caatcatctt	ataattttag	14520
ggactttgaa	tccaaagcaa	accatcttgg	tgattctggt	gggactcctg	tgaagacccg	14580
gaggcattcc	tggaggcagc	agatattcct	ccgagtagcc	accccgcaga	aggcgtgcga	14640
ttcttccagc	agatatgaag	gtaaggccgg	tacctgaaat	gaaacctcaa	agagagcacg	14700
ctgacagagg	accctgggag	ccycatcata	ttggtaagaa	agcagagcgc	cgtcctcttc	14760
agtattggca	ggtctgaggc	aatcacaaag	gtaactaggg	agggaattta	gaggttaccc	14820
tccatttctt	agggaaggaa	tttaaagcta	atttagggta	acctctccat	aaacaggagc	14880
agagctctga	tgtttagagt	ggtcacagtg	ttaaccagcg	gtgaatccag	acaggtctgc	14940
ggcaacetca	cttcttgcct	cctaggacat	aaggcaaaag	gagagactga	ggcaagtttt	15000
agagcagcag	tgaaagttta	ttaaaaactt	cagagcagga	atgaaaggac	gtcaagtaca	15060
ctttgaaggt	ggttaggcgg	gcaacttgag	agatgaagtg	tgagatttgg	ccttttgacc	15120
tggggtttta	tatgctgcca	tacttccggg	gtcttgcgtt	ccttcttctc	tgattcttcc	15180
cttggggtgg	gctgtccgca	tgtgcattgg	cgtgctagca	cacgggggtt	gtgggggagc	15240
gtgcgcaggg	tgtttactgg	agttgtaggc	gtgctcactt	gaggcgttct	tccctgtcca	15300
gtctagcatt	cctagaggaa	cgtcatgcac	caggtaaatt	ccgccatgtt	gcctcttaat	15360
gcgcatgctt	gagcccactc	gcccagctcc	cgagatctta	ttgggaagct	gcagctcccc	15420
agttttaggt	gttttctatc	tactgggagc	ccgcccttcc	ttggtgcccg	ctgtgaccaa	15480
cgatcacttt	agagaaacag	ttgacaactg	cctgaccaac	acctgatggt	cgcctgacat	15540
tgctggtgca	tatctggaaa	gggccctctc	ctgccgtcct	catgtctgac	gagctacccg	15600
ctgtaaccaa	agcgtgggct	tcggagtctg	ctttcaaatc	ccagcttttc	cccttaggag	15660
ctgtgaacta	gaataaactg	tctaaagtta	ccacctataa	cctgggatta	attatgcctg	15720
ttgccacact	gatagagaca	aggcagcatg	atatcattac	tgatacattt	tttttaaagc	15780
attcaaaatt	catagtactg	gaaagaaaat	cagtgatgcg	aatgtttcca	gggtaatgtc	15840
acctcccatg	ctgtggaagt	ccttcgggtg	agcctggccc	cttgcttctt	ttgccccagc	15900
ctttctatgt	gggggcacca	tggagctgcc	actcaccagc	acctttttc	cctcaagtag	15960
tttgtaccta	taaagtattc	ctgccgtggg	tggcccctcg	gtggagetge	tgagcctagc	16020
cagggtttga	tttctcttcc	tgccagtgtg	agccagatgg	ccacatctct	cttcccctgc	16080
cccgtggaga	ggtctgctta	ccgcaaagaa	gggctcttcc	tcccaggtcc	tgtagcaccc	16140
tgttagaggg	tgtggagtgg	agcagtggga	accagagcca	ccagagggag	gccctggagg	16200
aggaacgaag	ctgattcatg	tctgaaaggg	gtgccagaac	ccaagtttcg	gtgtttaata	16260
aagagtgcct	cggtgttgcg	gtggccatac	ctcacagggc	atggtcgctt	ggaaatttct	16320
gctcggaaat	gctttgtgca	gtggccagga	tgcgttaggg	gccacagatg	actgcttgct	16380
ccatcataga	⁻ acagttccaa	gttttcaaac	gagcattcac	agactgagcc	gcatcctgcc	16440
tccctgtcct	ctgattcctg	gcttcttctc	tggtctctga	agccacacgg	aaatgtgttt	16500
gcatctgttt	cctgcccttc	agatgacaga	ggaccatgga	agctgctgcc	tectttaget	16560
ctcttctcca	ggggaattgc	cctcgtcact	gtttgggaac	ccctggtccg	agtcctgtcc	16620
tccgaagagc	ctctgcccct	cctggagtcc	tgagttgaac	ttggtgttca	cttggcctct	16680
ggctctggca	gtgtgttgct	ccttccgttg	acctgccact	gctctgttaa	tgcagattga	16740
tcttcataat	ctgtttctgc	tttaagtgat	taactcaaac	attcttggct	cttattctat	16800
cttgtccttt	gggatatgaa	ccattattta	aatttggact	ggtttcctgg	cttggcacag	16860
ttgaccatgc	ctgtaagctc	agtgctttgg	gaggccaagg	caggaggatc	cctggaggcc	16920

WO 00/08209

26

aggagttega ggeeaccett ggeaacatag tgagaceetg tetetacaaa aaaataaaaa 16980 ttagctgagc gtggtgttgt gaagctgtag tcctagctac ttgggagact gaggcaggag 17040 gattacttga gcccaggagt ttgagtttac agtgagctgt gatcacaccc ctgcactcca 17100 gcctgggcaa cagagtgaga ccttgtctcg cggggtgggg gagtcatgtc tatacttgag 17160 aagttttttt ccctcgcata gtttgtacct ataaagtatt catcagtttt gagcagtcct 17220 tttgcgattg ttttgagtct tgattctggt gaccagtaag ttgtatatat ttgcctgtca 17280 agtggacaaa catggccttt gtgcctttaa gtaatggcta aaagtaccaa acagaacagg 17340 gcctggcata gatgctgctc ctcctgttcc taggccgtaa tcacccctga ttcatcagac 17400 cccaaacaag tcagctcctc tccctcctgt gccccaccac ccaatctcct gcaggaagat 17460 gtctggagac ccctgtcagc gctaggcaga gatcaccatc catgtccacc tttcctctga 17520 tgcaggetee cactageece tetggettgt gecatgeeag ccatgaacte acceteatge 17580 cccacccgag ccctggcaca ggctattccc tctgcctgga atgctcttcg tcagtatccc 17640 catggetece teceteceet tecettgtat cetgaetete ceatageage tetetecetg 17700 taacacatgt tcacaggttc atcttcgtca cccatctccg gcagctcctg caggcttgat 17760 ggctgctaaa ggcaggcaag tcagtggctc agattcttgc aaacttagtg attagtgatt 17820 totcaactoo otootcatgo otootctgto totataggoa catattattt ottatotott 17880 ctcagaacca agccgcctga atttctgaat aacattgttt aagtgttctg tgtatgcaaa 17940 agaaaaacga gaataaaagg attattaagg aagaattaat ataataatag ccacatatta 18000 tgctcttttt atactctgct aagtgcttta catgaattat ttcctttaat tagacaatct 18060 taagaacatc gacattttta tgaagcccat tttacaggtg ggtgagtgga ggctgggagt 18120 ggcttaaatc actttcccca aaccagggag ttagtgggag ccagaggcag gacctgagct 18180 cgcgggtctg agctccaaag ctcattctct gaactgtgca cagcactggg ctgcagccag 18240 agatgcagga cgctgcggga ccctctggag gtggtcctgc ctgtgcttcc ctcttcccac 18300 aggaagetee etataggeat etgtgttggg egtggaetet eagtgtacet geatgtetee 18360 ctgttggcca gacaccaaca ctgaatggaa aacatgtttc tgggcatttt aatgtacgta 18420 cttgccttca gtcaatctcc tccgcccct tccatcctga ccgcctccct aatagttagc 18480 agtgggactg gagcttgáat ggcaactgat ttctgtctga gaggacaaat caggcatctt 18540 tgtcctctgc cactgtctgt tccccatcct taggatgcac gatgccagag ccctccactg 18600 tggtctgtga ccactttgac ccacactagc aggtctccat atgttccttc cagctgagag 18660 acatcacatc caaagacagt ttagagctct gaggtttctt tccccagagg tccctgcttt 18720 gtgcaaactg tetecageea agegtgcaca agactetgtt cetgatttge etgggegget 18780 gagccatggg cagctgagcc tgcagccgct ggactcactg cattcccact ctgactttgg 18840 catgaaagac acacaagtgt gettgtgaga aatagatett aacagtacet tttaacacet 18900 atttcaggtg ctcaaaatga ctgcctgttt tacatttata ttctggcagt gcaaacttca 18960 attggacagg aaatcttaca acctctcttc caggtgraaa agcgaggcag ggatgtttat 19020 acagttccat ccatgtcatc ccacttggaa gatactagta aaacacacca acagtaatac 19080 aaaaaccatg gtgtttgcaa tagtgataat gttactagtg aaggaaaaat agaaactttc 19140 tgtaatttgg agattctaat ttttataggt gggctaaaaa aaaaayctgg agagaagggt 19200 gttaagtgag taaggagtgt gtctctaact aaatatagtg taaaaagaga agaaaataca 19260 aagtcaggca cagtggatag aggtggatag tctaatctct aatagtataa tgggcaaaat 19320 tgtctcaaac aaaattagtc tgcctcttgt ttactcaggg atgtgtgact gttttctatg 19380 cacaaaatcc ccatgaaata attaagttgc aagaatctga actttatatt ttggaaacct 19440

atctgaggta	ggtaggaagt	taatttatat	ttagaaattt	gcttgcatat	gtctagtagc	19500
tccaggacaa	atattcccaa	atcccagact	atttttttt	ctttttaaat	tcaacagtga	19560
ccagttggtc	tcttgtaaga	attacagcct	taagttagca	aagtctaaga	gggctggttt	19620
taatcctgaa	cctcagaggg	tccctgcttc	tcaaatacta	agtaggtcac	gtgcacagca	19680
ggtactacat	tgaagggaaa	ttgtatgata	aataggaaat	cagcgatttt	tacttggaga	19740
cttggcaagg	caaatgtttt	tgtaataaaa	atagatcgtg	aaatagaatc	ctgaaagctg	19800
cctgtttaaa	tgtaaagcaa	atggctttag	tgatgcttta	agtgtggcag	tcacttctgg	19860
ctgccgcaga	aactatagaa	agtgcattct	ctcttggtgc	tgtgggttct	tag g gtgaat	19920
gccttgtgtg	acgctgagta	tgtggaagga	ccattcattc	ttggtaacta	tacactaggc	19980
agagggtggc	gttagcgaag	ctactgcagg	ttgggtgtgt	ttaagatttg	gatttatttt	20040
tcttttaatt	tttatttta	gttccagggt	acatgtgcag	gatgtgcagg	tttgttacat	20100
ggttaaacgt	gtgccatggt	ggtttgctgt	acctatcaac	ccatcaccta	ggtattaagt	20160
ccagcatgtg	gttattttc	gtaatgctct	ccctgctccc	tgccgcccc	caacaggctc	20220
cagtgtttgt	tgttcccttt	cctatgtcca	tgtgttctca	tgattcagct	cccatctatg	20280
agcaaaaaca	tgtggtgttt	ggttttctgt	tcctgcgtta	gtttgctgag	gataatggct	20340
ttcagcttca	tccatgtccc	tgcaaaggac	atggtctcat	tcatttttat	ggctgcatag	20400
tagtccatgg	tgtatatgta	ccacattttc	tttatctagt	ctatcattga	tgggcatttg	20460
ggttgattcc	atgtctttgc	tattgtgaat	agtgctgcag	tgaacatatg	catgcatgta	20520
tctttgtaac	agagtggttt	atattccttt	ggttatgtac	ccaggaatgg	gattgctggg	20580
tcaaatggta	tttctagttc	tagatctttg	aggaattgcc	acaccatctt	ctacaatgtt	20640
tgaactaatt	tacattctca	ccaacagtgt	aaaagcattc	ttacttctcc	gcaacctcac	20700
tagcatctgt	tgtttcttga	ctttttaata	atcaccgttc	tgactggtgt	gagacagtat	20760
ctccttgtgg	ttttgatttg	catttctcta	atgatcagtg	atgttgagct	ttttttcatg	20820
tttgttggct	gtatgaatat	cttcttttga	gaagtgtctg	ttcatgagag	agacatattt	20880
gctcctctga	gtaaagggta	aggatgctta	cgtctgtgtg	acagccttct	ctctttttca	20940
gaacctcact	gtggatcgcc	atcgttggcc	tgtactgaag	gtaaagcaga	tagaggcagt	21000
ctcatctgtc	agatgaagác	ctcatacacc	tgttgattaa	gaggetttet	tcagatcatg	21060
gtttagagcg	g tg tt tta ca	aacttgacgt	gcttggagtc	ttctggaaat	cttgttaaaa	21120
ggcagactct	gttgcagtag	gtccgggtgg	gttctgaagt	tctaacaagc	tccccagtga	21180
ggctgatgtt	tcaggtccac	tgtgagga g g	cagggcttag	aataaacaac	cgtgggaaat	21240
ccagtccaga	tctttgatgc	atcctaggtt	aggcctgtct	gtcaggctgc	cctgggtctc	21300
tagtgatgga	ctccaggagt	ctctcaagtc	tcaaataagt	ctgagtcatc	agggatattt	21360
tttgagaaga	gttgtgctgt	ctgaagaagc	aaagagtgag	tgtgatgggg	aaaatgcagt	21420
gattaaaaac	atggtaaggt	ttaaagaaag	atttgaccat	atgccaggtg	aacccaaatg	21480
tatggtgctt	tgcgtctttc	ctgccctttg	gttttccagg	gaggcaaggc	cttatctctt	21540
atggagcaca	ggagacacag	tgtgggcgtt	tgttttctca	gccgtgggct	ctaacctaat	21600
tgtcaagcct	acaaaaaaaa	aatgctgaaa	atcaacttct	gactagatat	ctggtagtac	21660
		tctgggtgta		_		21720
_		tcccaacttt				21780
tatgtgctag	ggaacaatat	taaatttagt	tttatacttt	tcctttaagc	atttcagatt	21840
atattgtgca	tttcaccaac	aatagaagct	ttcagacttt	atatgtcttg	taa aaaaaa g	21900
cctaatatag	ataagaataa	tttattgatt	tgaaacc cat	tgta ta agaa	at agtccag t	21960

•			20			-
gaaacttaag	ttcaaagttt	tttttgtctt	gtggatgtag	ctatgtcaat	atgcctagtt	22020
tatagtaaca	ttaagtctag	tggattagat	attagatatc	aattgagatg	taagcagtaa	22080
taaacagtaa	tgcctaaact	gaagtatata	atctgaatct	ttatgatgac	caatttatat	22140
tattgtgaaa	aacttaggaa	ctgatttgaa	acatgattta	catgttttac	atgaaacatg	22200
atttacatgt	gtcatatata	gttttcaata	atttacgtac	cagcaggaaa	ttttagtgga	22260
taagtggaat	aaactgcagg	tgaaactttg	ctggaaaata	caagcatagt	gacatetgtg	22320
caccaaaagc	acctggggag	atttttaaa	acatgggcca	gacatgcctt	cctggctgtc	22380
teceteactg	tcagtgagtg	tggggatggg	gtctgggcct	gaattctttc	tttttagact	22440
cctcaggatt	ctgattgctg	ccacgttgag	agggttgacc	tcaattcgga	cctcagaggg	22500
tgacttgaga	aactgtcacc	acttggtggc	agtgttgctc	cccgcatctt	gattgccctt	22560
gtttctttcc	aatcccggaa	aagtgtgctt	gtttttttt	tttccctgc	gtgtttttgt	22620
ttttgatctt	gctataatat	ttatattcct	tgctcatttg	caacttattt	gaatggagag	22680
ctactttctg	aaatctagat	gtttttcttt	ttctacaggg	ttttagggca	tgggcaaaac	22740
acggaagaaa	aaagttgtct	tcagttggca	gagacgtgga	tttttaagat	tgttcttaat	22800
ttactttctg	tataactttg	cttttctgtg	gtgaacaaag	accaggttca	agataaaata	22860
ttgcaagcca	agaatctgat	tgttcatgga	tttctatggt	taaagatact	tgatcacctc	22920
cccatccgcc	ccctacccca	cccaccctgc	gccgcccca	caccccattg	tgcttcttgg	22980
cttgtcattt	caaaagtcaa	ggaagtcaca	gtgaatggca	agattttacc	tegaettget	23040
atttttgtgc	ctgttaacaa	ttgtgagtta	acactgactg	agcttttcct	agtgaacctc	23100
cggcgtttaa	acagccagtc	cataacactg	tgtgagggct	ggagctaagg	ttattggtga	23160
cacaagatag	cacctgagcc	agtgctgctt	ggtaggaggg	ctgaggggaa	gagggctgag	23220
ggcttggatg	ctgagatgct	agagtcacat	cgcctggatt	tgaatccctg	ccctcctgtt	23280
ctgataccag	ctgacccatg	acgatgctac	agcacctgac	agcagattcc	tccttagggc	23340
tggtctaact	ctagagtgtg	tgcctgtgtg	cctgcaggag	aatgtccaaa	gtgggtgatc	23400
ttgatctgtt	aacctttgaa	ttttaaccta	taccagggag	ccattgaaga	gtttaaagca	23460
agtgaatgac	gagtagtttg	aaaatatttc	caggtggata	gaatttgtgg	acatacatga	23520
acatgagcag	cctcaaaátc	agggctggga	ctagagtgag	gccagcacgt	gtccagggtg	23580
caaaatgtaa	ggaggcattc	actttcaggg	cctggcaggt	gtggaccctg	aacttccagg	23640
accttgagag	tgagtgt ct c	ctaaggatta	caccctgggg	gcctatttgc	ctcatcctgg	23700
tccctggtcc	tctgtgtacc	ctattgcctg	cttcagtaaa	caggcagccc	tgcaagggaa	23760
ggaagggttg	gatcagctct	gaggaggag	tttttttaga	aggatagatt	tgttttgttt	23820
aaaaaacagc	tttattgaga	tataattcac	atcctataca	gtttgttcat	ttaaaatgta	23880
caattcaatg	ttgtgaggtt	attttttggt	atatccacag	agttgtgtga	acatgaccac	23940
aatctaattt	tttttattt	ttttttttg	agacggagta	ttgctctgtc	gcccaggctg	24000
gagtgcagtg	gtgcgatctc	ggctcattgc	aacctctgcc	tcctgggttc	aagtgattct	24060
catgcctcag	cgacctgagt	agctgggatt	acaggcatgc	cccaccaagc	ctggctaatt	24120
tttatatgtt	tactagagac	ggggtttcac	catgttggcc	agactggtct	ccaactcatg	24180
gcctcaagtg	atccttctgc	ctcagcctcc	caaagtgttg	ggattacagg	cgtgagcccg	24240
acccaccgca	gtctaatttt	gaaacatttt	ttgtccccct	agaaaaaac	ctgtagttgt	24300
cacttgccaa	tctactgccg	tecaceteta	accatagaca	gcccctaatc	tactttctgt	24360
ctctatagat	ttgcctattc	tgaacacttc	atctaagtgc	aatcatataa	tatgtggtct	24420
tttgtgtctg	gcttctttga	tttaacatgt	tttcaaaatt	cattatgtca	taatacatac	24480

•			2)			•
cagtaatcca	ttctttttta	atgacttatt	aatattccgt	tgtatagaga	catcacatat	24540
ggtttatcct	ttaccagtcg	agaggcattt	ggattgtttg	cacttttggc	tgttacggat	24600
aataccgctg	tgaacattga	tgtatgtgtt	tttgtgtgtt	gaatgtgagc	tggtgtggaa	24660
actcctcctc	caggggggcc	ttacctgtga	ttctacccac	ggggatggtt	aagccagcag	24720
ggatgggaag	ggtttggtcc	tgctggccct	aggettteet	gcaggctgcc	atgtgccttt	24780
cttctgccta	ggctgaaacg	gaggctgccc	tggtttctgg	cactgccctc	gtgagtgtgt	24840
gggaaggctg	ggggaagcca	agtctccatg	gtgcctccat	cagggaccct	gcagctggga	24900
ggcagccaga	gggccacagg	ttggtagcat	tcacacagag	ctacatttct	ttttttt	24960
tttttgagac	aatcttgctc	tgtcgcccag	gctggagtgc	agtggtgcga	tctccgctca	25020
ctgccacctc	cacctcccag	gttcaagg a a	ttctcctgcc	tcagcctccc	'aagtagctgg	25080
gactacaggc	gtgcgctgcc	atgcccggct	aattttttgt	gtttttagta	gagacggggt	25140
ttcaccacgt	tgaccaggat	ggtcttcatc	tcccgacctc	gcgattcacc	tgcctcggcc	25200
tcccaaaaag	tgctgggatt	acaggcgtga	gccaccatgc	ccagcctaca	tttcttttt	25260
ttttttcttt	gagatggagt	cttgctctgt	cacccagget	ggagtgcagg	ggcaccatct	25320
ctgctcactg	caacctctgc	ctcctgagtt	caagtgattc	tcctgcctca	gcctccggag	25380
tagctgggat	tacaggcaac	tgccaccaca	cctggctaat	ttttttattt	ttatttttta	25440
atagagacgg	agtttttcca	tgttgaccag	gctggtctcg	aactcctgac	ctcaagtggc	25500
ctcaagaggc	caatccgcct	tggcctcccc	aagtgctggg	attataggtg	tgagccactg	25560
cacccaccca	gcccgtagct	acatttctgt	cagctgtttg	caaactgtgc	cccagaatcc	25620
cctggaggac	ttgtagaacc	accagttact	gggttacgcc	cccaaatgtc	tgatgctgga	25680
gatgaattat	cttgggtgga	gccctcaagc	cgcagcagct	gataagcat g	gggacctcct	25740
attctgataa	aaattccaaa	aaagtcctga	gtgattaata	aacagcacat	tgaaaattag	25800
aaatgagttc	tatggcaggg	gatgaaacag	gcaacaaagc	ctattttctt	tgcaatgaag	25860
cgcatcagat	attaataata	gccattgtaa	ttatctttat	catgtattaa	gcattttgtg	25920
tttttcactt	ttacacaatt	agatgatccc	cataggtatt	accgcctttt	tttttttt	25980
ttttttgaga	cagagtcttg	ctctatcccc	caggctggag	tgcagtggca	cgatcttggc	26040
tcactgcaac	ctctacctcc	caggttcaag	ctattctcat	gcttcaccct	cctt ag tagc	26100
			gctaattttt			26160
cagggtttcg	ccatgttggg	caggctggtc	tcgaactcct	gacctcaggt	gatc cgccca	26220
cctcggcctc	ccaaagtgct	gggattatag	gcgtgggtca	ccacaactgg	acttactgcc	26280
			gagtgacaca			26340
tggttcacat						26400
			aaattatcaa			26460
acttaaatta						26520
			ttgctaataa			26580
ttagcaactt						26640
gttatttata 1						26700
gagtacagta a						26760
tcccctgcct						26820
aatgtttaca 🤉						26880
cttgaattaa a						26940
cactgegeca (ggcctccatg	tatttgaatg	aaagagcaga	catctcctgg	aggt gg caaa	27000

			- 0			-
gctatgcate	g cccccctgg	aggggagctg	ggggctctgg	ggttacagtg	atggcacatt	27060
	teegetttgt				•	27120
					aaaatatttg	27180
cagcatgagt	actaacaaga	ttttttttt	cttttacccc	gatggagtct	cgctctgtcg	27240
ccgggctgga	gtgcagtggt	gcaatcttgg	ctcactgcaa	cctccgcctc	ctgggttcaa	27300
	tgcctcagcc					27360
agctaatttt	tgtatttta	gtaaagatgg	ggtttcacca	tgttaaccag	gatggtctct	27420
atctcttgac	: tttgtgatct	gcccgcctcg	gcctcccaaa	gtgctgggat	tacaggtgtg	27480
agccacgctc	ccggccaaga	ttttaaacat	tatttaccaa	agtaggaacg	tggtaattat	27540
ggtcttatat	aattctgaaa	atgatttcta	gtaccaaact	atgaatttta	tacttgaaag	27600
aatgatgggt	ttttcacaga	aagttgaagt	tattatggtt	tgtnttcctg	ttcanggtgt	27660
ttttgctgga	gaatgttcga	tgaacagcag	ttctggtgat	aagttatgga	tgtacacagc	27720
tggtgtggtt	tttaggattt	tattttgcag	cagcatcttc	ctcaaacagt	tgccagggga	27780
aggctttcct	tcttcttact	ggtaccagcc	tttctcttgc	agacaaggca	gtatgggagg	27840
gttgggagac	aaaacagaag	ctgttggttt	cttcagcctg	gcaaggattc	agattgcagg	27900
ttatagattg	gaggccgtca	gtggggatac	ctttccggac	aaagtggtgt	ttctgcctgg	27960
cactgcttgc	cagagaagtt	tcagttcttc	attctccgtc	agagaaaccc	atatggacca	28020
cattctgata	gttttcttct	gtttccctaa	caccgaaggc	tcagcccctg	gtgcaggtcc	28080
cagtgtacag	caggctgcat	acagttagac	cagatgttct	tgtagtacga	aaagtcaccg	28140
agtttccatt	cacttgtggg	tggcaggtat	ggccctcctt	acctcccatg	gcccaggttt	28200
ctctgtcctg	ccgttttcac	attttccagg	ctttcacctc	caggtaccaa	aattcacatc	28260
atttagagat	tgtgtctgcc	tgccaatacg	cggatgtacc	agtgagggat	tgttctcgcc	28320
tgacgagagg	tctggatgat	gagagagcag	agctggccct	ggggctcagt	ggtgacaccc	28380
tcgagcttgg	ctgcttctgt	tcttctgctt	cctctgcttg	gattccttcg	cctttggctt	28440
cccctccagt	tccaagcaga	acaaaacagg	agatatcaag	gaggaaaggg	tgacccctct	28500
atatctggag	agcaaaactg	tcgcggaaat	ccctagtgta	cttccatttg	tgtctcatta	28560
tctgaaaccg	agttacctgg	ctggtcacgt	gcagccacca	gaggcaggaa	ggtagtgatt	28620
ctgcctgtgt	ggaatgttct	agcattccct	ggtagctttt	gtttcttcag	gcagccatga	28680
ctttgcatag	atcatttcct	tttgcccagg	acactcctgc	tcgttttctc	ccctcctcac	28740
caaacccaca	gtgcattaac	agc gacagac	ttctcctcat	cctctcaggc	cacttggatg	28800
tcaccatttc	ttctctttac	ccctcaggcg	tagtcagcct	ctctgtgcct	gatgttttat	28860
ggctttgtgt	atgccccgat	ggagagcgtc	ttactgtgtc	ttcgggttat	ttatctcaac	28920
ctcgcatctg	tgctatcctg	tacagtaacc	aacagccaca	tatcactatt	taaaattaaa	28980
tacaaactaa	ttatacttaa	atgtaataaa	aatgtagccc	ctcacattag	cctcatttca	29040
agagccacat	gtggctaccg	tattgtaagc	agagctcaag	aacattcagc	aatattgtga	29100
gagtggctac	catattgaga	gcagagctct	agaacattcc	cttatcccag	aaagttctct	29160
tggacatgct	gctcaaggtg	gtgaactctg	agatctccag	tccccccagc	tccgtcactc	29220
agaaccacaa	atgtggcacc	atcagccttc	agggtggtgc	ttgtgttgta	ctgtctcctg	29280
actaaagaag	taaacttcag	gcagtcaaga	ttttctacaa	cccacactgc	tcctaaaact	29340
agtgttactg	gatatgtaaa	agctat tgag	cccagtgctt	tcaaggaatc	ctaaaagcaa	29400
	tcatgtatcc					29460
ggaaaggggt	ccactcaagg	tcatac aggc	aaggtcatac	agtcagtggt	agaatggact	29520

			31			
ggaatcttga	ttgttcatct	cactcccctt	tccattaagc	cataactgat	tgagtatcac	29580
caacctgttt	gttttccctg	attatgttcc	cttctcrcct	gtttaatcag	gtttcagctc	29640
cttgtgagag	gaagttgttt	tcagcttctt	agccccctac	ctgtaggcgg	tgctctggga	29700
acgctcagga	agcagatgca	tgtgagcctg	tctcagccaa	tgattgctta	gttgcaagaa	29760
acaaaaaaat	gactcaaact	agcttaatca	aaaggaggct	tttaaacagc	aagataaggg	29820
taaaggctgg	gaactaggaa	gttgtcagga	accaaggctg	tctctctgga	tctctcttg	29880
aggccatgta	attttttt	cctctcggtc	tctttattct	gcacaccagc	tcacccagct	29940
tgcttgttac	tatatgcagt	catgccaacc	cccagatctg	tatgacctgt	tagcctcagg	30000
ggcacccaca	tagctggctg	ccaatctgtg	ttttcttcca	ggtttcgaga	gggagaaatg	30060
attggcccag	ctcagggtca	cttacccagg	gagaggtggg	ggaagtatgg	aggcaccgtg	30120
gtatcaggga	acccctgggc	caagettgte	caacccgcgg	cctgctttct	ttcatttttt	30180
ctgttttttg	ttttgttttg	ttttgtttt	ttacagctca	tcagctattg	ttagtgtatt	30240
ttatgtgtgg	cccaagacaa	ttcttcttct	gctgtggctc	agggaagcca	aaagattggc	30300
cacccctgtc	ctaggccatt	atctgggctg	tgggaggtgt	ggagcagggt	cagagctgga	30360
gggggagggc	atageeteca	gccaccataa	gttggtgtgt	tcttggtaat	tatattgctt	30420
gtcaaccgaa	ggcagaatca	ggacaatgaa	agtaatgaga	atccctagct	ttgtaacagt	30480
tagtggttat	ctaaaagtag	gtgaaattgt	acatgagtga	gtggcatgaa	tttcttatta	30540
ctaaagtgct	cagatagctg	gctaactttc	tgtcaaagat	ccctctgcta	ggatcaacat	30600
ttgattaata	tatttatcct	gtaataagaa	tttgggattc	ttaaagcaaa	atagttgtca	30660
tgtggctgac	tacacaacca	aagatggtcc	aggtgtcgct	ggaagaggag	agactgaaga	30720
gctgttgcca	ggttcccacg	tggaccttcg	gcatgacccg	gccatgggga	ggcctcacac	30780
gctcctgcat	cgcccacatc	ttgccaagcc	atggaa a aca	cttgggattc	atatctaaat	30840
cctagtttaa	gcttggtgag	gacagtggcc	tggtgc ag ag	tttgggtcat	agatggtgct	30900
tggtttcttt	tgtataaagg	ggtatatgat	tttggaatat	ttaccaaatg	tgggcatttt	30960
ttctataaaa	attattgtat	ctactgagat	tatagtatgt	aaaaaaaca	tacacatgga	31020
gaaagaatac	aaagàgagca	ttgatattct	acagaagtgg	caagaagatg	tggtgatagg	31080
tgatatttt	gccttttgťt	tcaattttgt	attgtagtga	cttttttggt	agaaaaaact	31140
aantttctaa	ttaagggaga	aacatttgaa	gtacatttag	tctttctaga	aaacctcatc	31200
ttctcataga	agtttaagat	ggagacatac	ttccattgtg	aatcatgatg	ctaaccagta	31260
ttcagatttg	ttggaaatgg	actcagtttt	aaaattgctt	ctctcttgtg	ggctggaact	31320
gcaaatgatt	gtttggggat	ttttcccctt	ttcttctatg	gagttattca	acttggcatg	31380
accagtgatt	tgagctgaga	acatggaacc	cttgatttgc	agaaatcaag	ccccaaagg	31440
tacagataca	gtggtcattg	tctgaagggt	ttcttttgtt	cttggccctc	ctgtccctgc	31500
tcttactgtg	gcagctgcag	ctgcaggtgc	ctctgaagcc	ttg ccatcca	tggtcacttc	31560
ctgcctgctc	cccacccacc	cctgggaaag	agcccccaag	tgtccaaaag	cactgtgttg	31620
cctaatgctt	gttgagagtc	tacatttctc	tagat ctag c	agaagt aa aa	tttcagtttg	31680
ttatatttat	agtttcagga	atagtt tg gg	a atggattta	ataaaaaatt	taaaagccca	31740
tcatttttat	atctctttt	cgatatttga	tggtttaaaa	gacatcaaag	ttatcttctc	31800
ccattactca	tcctatacaa	ttaaaacctg	ttttttgaag	ttgtaatagg	taagttagcc	31860
ttaggtcacc	ccatatttat	gtaaac tcc a	gcccactgcc	acagctactt	tgattgtgat	31920
ctgtcattgt	gttacccact	gtagggcaga	aatggttcct	gcctcatgcc	gttgctgctt	31980
tactcttcct	gaagtggtgt	ggttctgtct	ctgtagtcct	tggcacactg	taggttctca	32040

-			32			-
gatggcaggg	tgaaaagttc	ttctgtttgc	ttaaatctct	cataataccc	tgagtctgtg	32100
gagctcaata	aattctactt	ggtattattg	atagattatt	tggagccttt	tatgttagaa	32160
aagggattct	taatccaatg	ctccgtttta	cagatgagaa	gactgaggct	caaagaccat	32220
acccccagga	gccatgattt	gcactgtatt	taggaatagt	gtctagggtc	agcacctggt	32280
gttggccgac	tgcagagcag	cctggttagg	agccctgggg	ttgggcgggt	ctgggctgct	32340
ggtgccacag	cagtctccct	cccctgggac	tt tg ggcctg	ctacccaccc	ctgttccttc	32400
ctttgtgaga	tagggctagc	agtaactgtc	ttgtttcatc	agaggcagta	ttgcataatg	32460
aatgagagct	ggggcctaaa	ttaggcacaa	gtgcaagccc	tcagaaaact	atgtacacct	32520
agagagagag	agagacacac	gtctgtatga	cagagaggca	gggtttggga	atgttctgat	32580
ttcatgtttt	gaattggtgt	gacctttggg	aggatatcct	tggaatcgca	gagcttcgtt	32640
tacatcatga	ctttcctgcc	cacccacatt	ttctgagaag	ccagagtttt	aaatgtggac	32700
cccgtgagct	tttctctgtt	gcctcatttt	ggcctgtggc	cttttgtttt	cttggtatgt	32760
catgaggcaa	aataaaatga	aactcagtgc	tggttaataa	ctcccatcat	aatgtatatt	32820
tctgtgaatg	gctttttagc	catttgagag	gaaaaagggt	catgtaaatt	tcagaaaggc	32880
ctgattggct	ggagagtcag	tgtagtgtca	cagttaagag	tatagattta	aaaaaaattt	32940
tttattgtgg	taaaaaacat	aaacataata	ctaccatcta	aaccatattt	aagtataaag	33000
ttcagtagtg	ttaagtatat	tcacattgtt	gtgcaatgga	tctgcagaat	ttttcatctt	33060
gttaaactga	aactctatgc	ccaataaaca	actcctattc	ccctctccc	agcccctggc	33120
aaccaccatt	ctactttctg	tttctctgag	tttgactact	gtagataact	catttaagta	33180
gagtcatatg	gtatttgtct	tcttatttct	ggtttatttc	gcttagcata	atgtcctcaa	33240
ggttcattca	cgttgtagca	tatgacagga	tttctctctt	ttttttccgc	ctttttttg	33300
agttatattc	tgttgtatgt	atatttaaca	ttttcttcat	tcatctgttg	acattcatct	33360
gcttccacct						33420
atgtctcttc	aagateetge	tttcagttct	ttcggatatg	tacccagaag	tgggtttgct	33480
ggatcgatca						33540
cattttacat	<i>f</i> .				_	33600
cttgtaatct						33660
tgtgttgggc	ctggatgttt	agaacagtat	ccctcctttg	gagtggtaaa	tatgtaagtt	33720
tttattataa						33780
tttccttcat					=	33840
catatttggg						33900
tttgattttc						33960
gcccttccat						34020
gtaatgccgg						34080
gaagcactca						34140
aggggtctga						34200
cagccattct						34260
caaccaaaag						34320
ccaaggatct						34380
gctttttggc						34440
agatagcagc						34500
tctgaattga (gggacagcaa	aaacctctaa (ccaaccatac	tgaaaagcag	gcatt ggg gg	34560

•			33			-
ctttagggga	aggttctttt	caaaactcat	gatggggaga	gaccaaagac	tgggaatcat	34620
tgtaaagaag	ttagtcatag	atgetteact	ctttacaatc	atcccaacac	aaggttaaac	34680
aacatgcagt	tttcacgatg	tcccagaaag	cgacgagtgc	agtgaggtga	aacgtggcca	34740
			atcgatttcc			34800
			gcactcgatt			34860
			aaatatgcaa			34920
			teceetgttt			34980
			tgatgccaca			35040
atttgccata	ttcagttatg	gtcagtagag	actttcttag	tctctctctc	tttttttt	35100
tttttgagac	aaattcttgc	tctgtcaccc	aggctggagt	gcagtggccc	aatcttggct	35160
			attctcctgc			35220
ggattgcagg	cacgcgccac	catacttggc	taatttttgt	atttttagta	gagacagggt	35280
tttgccatgt	tgtccagact	agtcttgaac	tcctgacctc	gtcatccgcc	tgccttggtc	35340
			caccgcgcct			35400
			gtttttagtc			35460
			ttcaaggagc			35520
agtcataaaa	tggattaaag	tgtttgaaat	ctattgggat	tgtaaattta	tgtcagtgta	35580
ctgactttca	agagatettg	atgatcatgt	cgtctgtttt	cattttctac	tacatgagaa	35640
cattgaagcc	tgaaacttaa	cacaaacccg	agttccccac	ttgcctaaga	gtcatggata	35700
			ttctttcagg			35760
			aggtcacatt			35820
			cactatttta			35880
			cacaaaagta			35940
			tcacgtccag			36000
			gcatgagagg			36060
			tctcttgtca			36120
			ccatcacggt			36180
			cctcctttct			36240
			accagetett			36300
			gttgaatagt			36360
			ctgctaataa			36420
			ttcagcatgg			36480
			tgtccttcac			36540
			aaaaccatca			36600
			ccccatgat			36660
			caattcaaga			36720
gccaaaccac						36780
gggtcatgat						36840
			aagggtggtt			36900
cgacacactg						36960
aaaatccaaa						37020
ggtaggatta (taggtaactt	ttttttcct	tagataaata 1	tatagataga (tatattagtg	37080

			3 -4			
tttacagtt	t ctctgccaco	aaccaaaata	a tttttttcag	gaggaaaaaa	aaccccagcc	37140
	a cctaaaaac					37200
tgaaattgc	c atttgtgcca	ctgttgtatt	attttaccag	taactccaga	ttccaggctc	37260
ctgtatctga	gttctctctc	cttccacagt	ggagctcata	ccttcctgtt	tectggetge	37320
cactcagatt	taggctccgt	ttttcagaco	tcagtggctg	taatagctgt	tccttctacc	37380
tcttaggato	gttctttctg	taatagcctt	tgtcatcaca	tcatcagagg	atgatagctc	37440
ttaatgagga	tctaaaattt	gcaggtaaga	tatccctgcc	tctgacatga	gatagatgta	37500
ttgcatgcta	tttaacatac	aactatactg	g agtgtgcagt	tgtatgtaaa	agcattgttc	37560
taggtattgg	gttgaaagtg	gatcaaatgo	: tagacaaagg	agcgtacaag	tcttgtaagg	37620
aagacagctg	, ccaagagaga	agaaaggatg	gggaaatgct	gcgtctacta	agttcaaggt	37680
tctgaattgg	aaagctgcag	ctattgagga	gaagagtctt	ttaaaattcc	taaagggttt	37740
ttgttatctt	ttattgatgc	aaatgctatt	ttgtggcata	aaccttaata	attt tgg ggt	37800
tgaaactctt	atcaggataa	aatgatcctt	ttctatccca	agcttaataa	atattgttta	37860
agtacaaatt	aaatatatga	aatctgccca	tctatattat	aaatgtcata	tggcagaaat	37920
tataccttga	cttttggttc	tttcacaaaa	ccttaatttt	tttttttt	ttttgccttc	37980
aatgaatttt	gtctgatttt	acattaaaag	cctgtaattt	ctcaagtctt	gagtctgggg	38040
agccgtcgtc	atccttttt	cccctctccc	ttgtcttctg	gatgttcaag	cgattttaat	38100
tagatgttgg	gcttttatgt	caagtgctgg	cattgcactc	catgataatc	cagggactcg	38160
gaagcacatg	ttatgcgtca	ccctgggttg	gtgcagtgga	actggggtgg	gttggaagta	38220
gtattctaaa	tctgcttcct	gcgatggggt	aggtcaggtt	gtcctgtgtt	gacaaggaag	38280
aagtctgggt	gaggaagcgg	gatgaaagca	gaccagacgc	tagagtccac	tttcaagtcc	38340
gatcccagga	cctggcttaa	agttaaagaa	cagcaaagat	gaaaggtgcc	gcacagcagc	38400
acaggtcggt	ggccacgtta	atgacataga	aagcaagtgc	tgtgaattca	aaagaaagga	38460
cagctctgag	ccagagtact	tggtgacttt	gctcaaacaa	atccctttct	ggcaccccca	38520
ggccttccct	cccgcttcaa	aaaaattctg	aattgtgcca	atccattgag	gctcagctca	38580
aggccatccc	atgcctttcc	atcgtaataa	agccttgttt	cctgggcttt	aaacatattc	38640
ctttttctt	aggtacagat	tgaacttttt	taaaagggaa	gttgtcagag	gctctgtaaa	38700
acgttaaatc	aaacctgctt	tgttttaggg	atggggtagc	ttggaatcag	atttgctcct	38760
gctatggact	gaacatttgt	gtccccccaa	aattcctatg	ttgaagccct	aatgcacagt	38820
gttatggtgt	ttgaagggag	gcccttggga	ggtgattaag	tttagatgag	attgtgtgag	38880
tgaagccctc	atgaatggga	ttactgtcat	cccaaa aa ga	ggtagagacc	ccagagette	38940
ctctctctc	accctgtgag	gatacagcaa	gaaggaagct	ctctgcaagt	cagga aga ga	39000
gagggctctc	actagaatac	acttgtactg	ccaccctgat	cttggacttc	ccctccagaa	39060
ctgtgagaaa	caaatgtgtg	ttgtttaagc	cacccagtcc	ctatgatttt	attagagcag	39120
cccgagctcc	attctccact	ccctggcttc	ctgcatggac	tttgcaacca	gagcttcacg	39180
gggtatagtt	taatagctgt	ttctctgtaa	cgtagccact	tttctctttc	caggtctagt	39240
tttgaccc tc	ataacacttt	gttaggggag	atttgagggt	gaggaagttg	gcttgctttt	39300
cttttcacca	tgtctcagta	gaaacagaag	cagaaaggcc	ctgagatact	gagcccacct	39360
ttctcagcag	ggtgtgacag	cccggagtac	cctgggctga	ggaggccagg	gctgg agg gg	39420
	ggtggagggg					39480
	tgggttgaga				•	39540
tgttaggtga	gtcatcagtc	tgtctctgac	tatgttcaga	tctggaagtt	ttctg g aagg	39600

WO 00/08209

aaatttgtta ttgctgtaat agtgtaggtt gttgatctgg attagcaggg agcggccct 39660 taatacattc ttaagaaaat ggtatttagt tcagtctttg gctttgaact ttgcctttga 39720 caaagatgaa agtgcgactt gactggtgtt tgaaaaacat ggtgatatgg ccaggtgtgg 39780 tggctcatgc ctgtatccca gcacgttggg aggccgaggc gggcagatca cctgagatca 39840 ggagttcgac acctgacttg gtcaacgtgg tgaaacctca tctctactta aaatacaaaa 39900 aaattagcca ggtgtggtgg tgtgcaccca taattgcagc tacttgggag gctgaggcaa 39960 gggaatcact tgaaccctgg aaggcggagg ttgcaatgag ccaagattgt gccattgcag 40020 tccagcctgg gcaacaagag cgagactcca tctcaaaaaa aaaaagcaag ttatattaca 40080 ttttaaaact ctatttaatg gtcaggtcat ccatccataa tgggtagagt cattgcttaa 40140 ttaatttaaa acaatgtatt taaaaggtac ctttgttccc tagtgtcaca taacgtgaaa 40200 tatccaatta aggtaactgt aatgtaaagt aagtggctaa aaaagtgctg aacgccaaag 40260 gccagagatt caaccttttg tgtgcattag aatttcccaa ttgttcaaat ccaggttgct 40320 ggatctaccc cagagttttt gatccagtag gtttggggtg ggaccaagaa tttgcatttc 40380 taacaagete eeaggtggtg ttgaggetga agetegtgtg gggaccacat tttgagaact 40440 tetecegtag actgaactca tggtetaggt tetgteaget gtgacceetg tgetgetgga 40500 gggagtggtc agatgtcctg acctctgtgc ccacagtgag gtccaagctg agtaggtttg 40560 accagcaget gtaatcacag agtgaacaat gtaaacgace aatgttgggt ggtctgacat 40620 cttttaaaaa aaatccacgt ggatgagatc acagggttaa gtgtgggcag cagtcagggt 40680 aactccatgt ggttactgcc catgcactct ctgctgtttt tcacctcttc ttcagagtgt 40740 40800 acccacetet tageatetge aggeaetece tgteeetteg etgggeeeeg tggggaacta 40860 cttgcagtca tcaaattcat catgctgctt tcttttaatt cccacacttg ccaaggtggg 40920 actgccccgc atctccttcc cagtcgtgtg tcagaactca gcactggacc tttccccttt 40980 ceccaetece acceetecte acceegacga acgteteact tgggateate tettetgagg 41040 ttggacctgc acagccgccc tctgcactct cgccacctta tgggctgccc ttgaccctt 41100 ggcacacaga cetggaagtt ggcetgetea getgteteet taggggtgga gettggtttt 41160 ctttcatcac tgttctgcga tgaattgaat gcatgattgg tcacaggaag gtaggggagg 41220 gataaacacc ttatgatatg tttcttataa ggttttatat gtagaaagtt atatgaaagt 41280 gtcagatatc tatatatgaa gtatatgtga agttttatga tagttttgca taatttaaga 41340 ataaactett taaaggaget gagteecaat eeettgggte gagagttgeg tggeteeegg 41400 ggcctgcttg tttccttcca ctctgcgtgt tcgttgctgg cccctcatag gctgtcccag 41460 acctetttga ettetetet ttetgeccag tettecetga gaegetecag getecetgge 41520 ctcctgcttc tcggagcttc tcttgtgttt gttttctgtg ctcagggcgc catggtgcta 41580 taggccacag aggaggcgtc tggggtccct cggggcaggt gcagcaggag gaagccgtct 41640 ccgagggcat gaccttggaa ctgagcattg acagaggaga gtcagccaga caaagaaagg 41700 ccaaaacccc acccctctcc caccctattt ctacgtgacc atgggccctg gacacagcaa 41760 gacggtgacc cegggeetee tattgttgeg aggageeeet gggaaaatgt tggcatttte 41820 ttcatagaac aggtttctct tctccagtat tcttcagtaa atcaactttc ttttttatcc 41880 ccaaccccag tetgattgeg aagaagteta agcaacagaa agattttgec aaatagatta 41940 tettttttag aacaaaatag atcatgatat taataggaat teageaetta etettgteta 42000 agtactgttt ttaagtgctc tcaaggattt ttcatttaat ccccacaaca aagctgtggg 42060 gggtggatgc tattattatc ggtgatttat gaatgaggaa actgacacag aggggtggtc 42120

gaggagcttg	cccatttcct	ggtagttagt	accagggctg	gcatcatcag	ttgcctgctc	42180
cttttcctct	ttgcttttgt	gtccattacc	ccaaggcatt	aggatgagcc	agccaagttc	42240
tagtcctgga	ttcaccacct	aattagctct	gtgtcccatg	tcttgccgtg	gagggataaa	42300
accaattcct	agcttatccg	ttggtggtga	agatgaaatc	agtggggtac	ttgtaaagca	42360
cactgcccag	cacatagtaa	gtgcccagaa	aatgtgacgt	cggacctctt	taagcttcag	42420
tttccacatc	tgggaagaga	gggggagttg	agctaagtca	ttttccagtg	tccctttcag	42480
ctccatgttc	ctgtgagcac	tgacagtttc	cccacaatmc	tgaagaaaga	aggaaaataa	42540
gggcggggtg	gcgaaggtcg	ccactgtgac	gtggctgctg	gtgggaagtc	cctggggagg	42600
caaggcccag	cttcccagac	acagccctca	ggtgctcatc	ctggtggcac	tgaccagggg	42660
ccatggtggg	cttttccacc	ccaccatgtc	tcataaaatt	acaagaacca	'cagttgaaaa	42720
tcagtgttac	agaaatggta	ataggatagg	gcaaactgtt	acaaagatca	gcacttaaga	42780
ttctggctga	ggcggaatat	ttgtttctct	ttagttttgt	tgtctttaat	caagaactga	42840
gagccctgac	tttcagctcc	tcaaaaaata	cagcttcctt	ccccttgcag	atgcaaaaac	42900
aaacgccact	tctttccaag	cataattttc	tcccatgcgt	tatctcctgt	ctacagettt	42960
ttcttgatcc	ttctccagct	cctgtagacc	tcccatttag	agccaccagc	cgcccatcac	43020
tggggctgcg	cagagctctt	ggtgctctgt	gccctgggct	cgcccaccca	ggcctgttct	43080
ctgngcctct	tcctggttct	cttccctgga	cttcccactg	ccgtgtggnc	ttcagtgctc	43140
ctctgagctg	ttgtcatgac	ctctaaccag	actgagtcag	gactttttc	ttcctcatct	43200
ctaagtcatc	cttacacagc	cttggaagtt	taccctaaat	ggctattttg	ggagggagtg	43260
gggataaaga	tctgcaggcc	tcttgctcct	ggtccttgtt	tctgcttatc	ttggcttctg	43320
tttttaagtg	tgtgtgcacc	tctttcctca	tcacaccctt	cccctccgta	tggctcccat	43380
ctcaggcaga	gttaggtgct	ctgttctgtg	tccatagctc	tttttcgagc	ccttcttctc	43440
actgtttggt	agtggccttt	catgtgtgtc	tgatccacta	ggctgtgcac	tccctgcctg	43500
ccaggatatg	gttaaagtgc	taaagaatgt	atatatgaga	tcacttttgc	ttaaaaaacc	43560
cccaatcttc	tggaattccc	aatttctaac	caattaatat	gtggattgac	tagaccttaa	43620
gcaaccaaga	gtcagccagc	cttgtcttct	atattcaggc	gcatactatc	tggtcgttag	43680
acaaaatggg	tcattatcag	tgatgagtta	ataattacct	gcacatcttg	tttatgctgg	43740
ttctttacct	aaagtggctc	ccatcaatta	aacctgtatg	gattttacct	gttcttccag	43800
aaccacccca	ctttccacaa	aaactgacaa	caatgatggt	aagaagaatg	gtagttgaca	43860
ttttattaaa	tgtttactgt	gtgcaggctt	gttttttcc	acacatttac	ctacttaatg	43920
ctcacaataa	tcctatgaac	tagtcagttt	tatgcagatt	tcgcagatta	ggaaactaag	43980
gtggcaagtg	atcagataac	ctgtttgagg	ttgagtagct	agatcatggt	agagccaggt	44040
tcaatcccag	atacctggct	ccagggccca	tgctcttgac	cttataaacg	gctgaaattc	44100
atctttttt	gctgaacttc	cagaacactt	tctttgtatt	tcccttattt	tggtagtctt	44160
gtacttctct	gctaccctga	ttcatacttg	gatttctagc	agcatgcctg	gcatgaggca	44220
acaacttaac	agtatttctt	tataccaaat	gaatgttgtc	tttttttt	tttttttct	44280
tgagacagag	tctcgctctg	tcgcccaggc	tagagtgcag	tggcactatc	ttggctcact	44340
gcaagctccg	cctcccgggt	tcatgccatt	ctcctgcctc	agcctcccaa	gtagctggga	44400
ctataggcgc	ctgccatggc	gcccggctaa	ttttttgtat	tagtagagac	ggggtttcac	44460
cgtgttagcc	aggatggtct	cgatctcctg	acctcatgat	ctgcccgcct	cggcctccca	44520
aagtgctggg	attacaggtg	tgagccacca	tgcccggcca	tgaatgttgt	ctttaaaaaa	44580
ttctgttttc	ctctagctag	actgtcatat	aatgcaactg	taggaaataa	tcaggttctc	44640

tttggagtat	tttccataaa	agatccacag	aagtcatggc	agggttgaga	gtggacttgg	44700
gcaaatgaat	ctgttcattc	attgaatatt	ccatgcatat	ctgctgtttc	ccaggcatgg	44760
gatatggcag	ggaacacaga	aatctctgcc	tcctgggctc	tgctttctgt	tgtagtagag	44820
gtaaagctgc	tcatactttg	taaacaatat	gacaacatta	agtctacatg	gtcattttac	44880
tttgttttt	tctaagaaat	tttgagctgt	tcgtaacaac	agacgctgca	gatgttaatc	44940
ccgttgttgt	taacttttct	ccagagattt	aatgttcaat	tttctccttt	ccagaatcga	45000
tttatgttgt	tcaaacagag	gtttgagaat	aactggaatt	tttttaactt	cttttttt	45060
tttcgcatgg	agttcagaat	tttcaagagg	gatgaagaga	gttataaaat	gctctatggt	45120
gggtaacaca	cagaaaaagc	cagaaaattg	gagaataagg	atctgtctac	tcgtttcctt	45180
ctagagetee	tctttcttac	agggcactta	acatgtgatt	taatgtcgtg	tctttaaaag	45240
gaggagaact	gcagttcaga	acttaatgtc	agtgctttgt	gaaagtgcaa	gaaagaagcc	45300
ctgtattctg	cacttgagag	agccagatac	tgggcagata	ggaggtggtg	tgcacgttgc	45360
tttttgtctt	tctcgatcat	ggcattgatt	ctgttcataa	caatgatgca	atgtcatcct	45420
cttccccaca	catttgtgtg	cagatagaaa	gaatgcaaca	gcacagagtt	gttggggaat	45480
aatttggcat	ctaaaatatc	gacataccag	catagatcat	atttatgact	ctgttgggag	45540
tgtcacagca	atgatttaat	aggaggcagt	tgtctccaag	gcctcctgaa	ttatgactgg	45600
ttttaaaatt	cttagaaccc	attggaggct	attgtttctg	aaaggctaca	taatttaagt	45660
gctccacatc	cgtcattata	ggagatgtca	gaatagtaaa	atctaatcct	ggactaagtt	45720
gttatcgcag	ccctttggtt	tggtggcttt	gccgacttta	taaatatgcc	tgtcagtgcc	45780
tgtggtctct	acagttgggc	agtcggcggt	gaatatcatt	tctcacattt	tacactgggg	45840
gactggaacc	cagaaggcat	atgttttccc	aagaggcacc	aacacagttg	gcccatgagg	45900
tagagcagcc	cctccttcgg	ctcagcctcc	gctgcactga	gccaagccaa	gcttcctaca	45960
ctggcctctg	tgcagctgtc	tctcagcaag	aatgcaagtc	ggggagagaa	gccggatccc	46020
tgggattgtt	ctagagagta	gaaacctcag	agtagccctc	cttagaccac	ctaacgcatt	46080
gcatcgctgc	atacatgtaa	gggactcaat	gctggtagga	ttggcttagg	aatgatgcaa	46140
g tgaaaacag	tgccccggtt	tatcattaga	acaaggttct	tagctgacag	ttgcctcaga	46200
ctttgatttt	gttctccttg	acctgccact	ccactcgagt	ccacatctct	caagactgca	46260
cacgcctgaa	ggaggactga	ttacaaacca	aagccttgtg	cccagtctgg	atctttttgc	46320
attgttgaga	aagcagctta	ctttctttgg	actgattcag	caggccaaat	ttagaacaaa	46380
gatttttaac	tatctccctt	tataaattac	tgagctattt	tgtagccagg	ctactcttaa	46440
tatgaacaaa	aaatattata	caaatttgtt	gttaatcgta	aactataaaa	aaatcagtaa	46500
ttgttaccac	gtgaaatgaa	tttggataaa	agagatacgt	ttttgcccct	tcccagggtt	46560
taggagagac	gaaatggtga	gattttagct	ctgaatcaga	ggttcttatt	agaggtggtt	46620
ttgttcctcc	tgacccctag	gggatattta	gcaatgccta	gaggcattga	tggtgggcag	46680
atgctactat	gccctctgct	aaacattcta	cagtgtataa	aactgttcct	cctgacaaag	46740
aatcatccag	ccccaaaatg	tcagtagtgc	tgaggttgag	aaaccctcct	ttaaactctt	46800
gggtttattt	gctgaccttt	acagtggatc	agcttttatt	tagttcatgt	agaggtgaaa	46860
ttaatactag	tgctcaaata	tgtctttgta	ttctggactt	ggcctggatc	ccccgaccaa	46920
atttgggaca	agctcctgcc	atgtgttgag	gacctgaatt	caggcagcta	acaacagtat	46980
ttgaactgtg	ttttcagtgg	tgggagtgaa	ggagat gag c	cgacgtgcta	gcaagcgcat	47040
					ttcagagtgt	47100
gcttggagaa	acagcagtgg	aggcattact	ggggagcttg	atggaaatgc	tcccctcaga	47160

cttgctgaat caaaatcttt aatttagcaa gatccccagt gaggcttgtg catgtagaag 47220 ttagagaagc acggggtaaa ctcttcttt ttttactttg gaggaaaata caccttttt 47280 cttattatgg ctctgaccct tactagctgt gtgaccttgg ccaagttata aaacctcact 47340 gcaccttatt tgttttagct ggaaaatgga gatcataata tcacctgtcc tatgagattg 47400 ttgtaagaat caaacaagct tatttatgcc aagaacccat atggtaaaag ctcaacaaac 47460 tgtcactagt gataataaga aaaagatcac aaaagtagaa aacattaggg agacagctta 47520 ggtcttaaat ctcacagttg tcgtccccaa acaatacttg tatttttgca gatccagttt 47580 ctctgaatac taaaataaaa ccggagtttc ataaacttct atagacagtg gtccttgtca 47640 gtagcccaag tggcagagag tacatggatc tggggacaaa cagcctctac tgttaggaat 47700 gttccatcct cctggcctga gttacacctg ctcattgtga ttccgaattt gaaaggaaca 47760 cagtaggaat tttcaagacc ctgggaagag gaaggctgtg gtaaacagga aggatgagat 47820 tagaagaagg agtttaggtg aggtgagccc ttgttttact agtagggttt aagaatatcc 47880 aagtcagctg gacatggtgg ctcacacctg taattctagc actttgggag gccgaggtgg 47940 gcagatcacc tgaggtcagg agttcgagac cagtctggcc aacatggtga aaccccgtct 48000 caactgaaaa tacagaaatt agcagggcat ggtggcgcat gcctgtaatt ccaactactc 48060 actcgggagt ttgaggcagg agaatcgctt gaacttggga ggtagaggtt gcagtgagcc 48120 aagattgggc caccacactc ccacctgggc aacagaatga gattccgtct ccaaaaaaaa 48180 aaaaaaaaga aaaaaaaaa aagaatatcc aggtcaaccc cacctaaccc tcagcgggc 48240 tecettetgt tgeetgggtg ggteetgggt tetettgaeg cacaegagat tgtgagagtg 48300 tatggaaaca ctgccctcgc tatcaggaca gcgcctgcca tgccagccag aacacatcat 48360 aggaattgca aaactctttt gcaaaccagt gagagatatg cttccaatgt gaggtaaagc 48420 agaactttaa tcacagctgc agtgttccac agaattccaa gagccaagat ggtaaaagaa 48480 taaaaaaaaa gaaaggaaag ggctcaaatt aaagacttca agctgcagaa taagattaaa 48540 taaaaggatt caattgaact gcatcatatt cagtaatgac taatcctaag tatacagggt 48600 ttgggggtga aaggatttgt aagtgttttg caggaaaata ttttttccat ctttcatttt 48660 aattagaata gatttgcatt attttttctt agtttttatt tttaaaatat ttattgccac 48720 aaatttagaa aatacaggga aaacataaat aacagtacat gtaaaccaat attttgtccc 48780 ttcttttgtt caacagctat ttctcaggca cctgctgggt gtcagcagct gtgctcagtg 48840 tggtgaccaa aaccettgte aacaaggeag caaggtteta acctggttag ggettacagt 48900 tgagtagctg aaattttgat ttcttttctg tgcccctagt aaagatatga tagcaaacaa 48960 taagagetat tttttttatt gtgttettae tetgtgttgg geeetgttet eagtggttta 49020 tagectatta acteagtete tttaceacea etetgagggg aggetetgte atacecaett 49080 gacagatcgg gaagtggaag catcaggagg ttaagcaact tgttaaagat cacaaaatca 49140 ataatgacag agttttgatt agaatcccag cagcctgtct ccagaacctg ccctattaag 49200 tgcagtgcaa ctgtactgcc tttcataata tgtatcaaat tgagatgata ctttataatt 49260 tcaattettg ettttetatt gaacagtaca cagtaacate etectataat geatataaae 49320 ccccaaaaga tgtagaattt taatttattc atttgtctga taggctcata atgaaataag 49380 actctataaa gctgtgtaat ttagatatag gaaacatttg gattatagtg gtatgtagtg 49440 ggaacaaatg gtcttctgaa tcaggaagac atgagttaga gtatgccggt gtacctcctt 49500 actcactgta tgaccttggg caagtttctg aactttagtt tcctttccag gctaatatct 49560 gccttctgga cttgtcatca ggattaaatg agtctaccta tataaaatgc ccagcgcagt 49620 gcccagcacg tggtagaagg tctgctagtg gttactgtta ctgctggcta ttaaatacat 49680

tttaatcttc	cttcagaata	cctggccaga	tagcacagtg	gttaagaatg	cacatgaaag	49740
ccagactgtt	gggttccagt	cctggctcga	ctccttccta	gctatgtgac	attaggcaac	49800
ttacataaac	tccttgttcc	tcagtttgca	tttctttaaa	actgcatagt	tatcataccc	49860
atgtcttaga	gttttgtgag	tgtaaattat	tgtatataaa	gctctgagaa	cagtttggta	49920
cacagtaggc	actgtatgaa	cattttctgt	aattatcaat	aatataatta	ttaaataaca	49980
ttttcagaag	gagataaaaa	tattacacct	taaaaagcag	gtatctttaa	attcttcctc	50040
agctactgaa	gttttgctta	ctatttgaca	tatcatttgt	ttcacgtttg	tggctcagac	50100
gtggcttatg	ccaatgcata	ttaacacagg	aattttaaat	ttggtgatat	tattatattt	50160
tatctgaatg	aacagaattt	gctgatttga	cactgtgttt	gaatgtgcat	tttttgttga	50220
aaaatgacaa	ttctggaatg	ccgtctccct	ttccagatta	ttcagagctg	ggagagette	50280
ccccacgatc	tcctttagaa	ccagtttgtg	aagatgggcc	ctttggcccc	ccaccagagg	50340
aaaagaaaag	gacatctcgt	gageteegag	agctgtggca	aaaggctatt	cttcaacaga	50400
tactgctgct	tagaatggag	aaggaaaatc	agaagctcca	aggttggttt	gccatcttga	50460
tattgaacag	gcctggtctt	atcttggctc	tgaagttaat	cacatcagac	ataagcatgc	50520
tgtcttaaaa	atacagcagc	acgatagtct	aatgtataca	tctatctata	tctgtttact	50580
ttttcagagt	aatattaaca	ctgtttactt	tctggtgatc	taatgatagt	ttcaccaaca	50640
atattcatta	ttcctctatg	gtcactgtta	gtacagtgtt	tagaacttct	gagatccaag	50700
ctttaaatct	aagctctaac	acgctgaaag	gtgcttttca	ttttgttttg	ttttcccctc	50760
tgtctctctc	tctctctcta	ctttatcctc	agccatggtc	tgtgcctgtg	tgttaggtat	50820
gaacttttct	tgtgtaagtc	attaacatac	gtaacttcac	tctgtgtgct	ttttcagtga	50880
tttgcaagta	atctgaaaaa	aaagaattag	ctgagttcta	cctgtactga	tatcaatagt	50940
gtcaaaatat	gacatgaact	ttgaaagttt	agattttgtt	catttcctgt	ttccatgctg	51000
acactggaac	caattaatgt	tatcttcaaa	gtagcttaag	atgcaaagtt	tacatactct	51060
ttggaaagag	catgagtctt	agggtatcta	gagaactgcc	cggtgataaa	gtagtgaaga	51120
ttttgagcag	gaagtctgca	taatctcttt	caaagggaag	atgtagcaga	tggttcagtc	51180
accctgccat	tgccċcagaa	caattttgga	attacagtac	atttcattca	gcatcattct	51240
tgattgcaaa	ttttgatctt	ttaaaatgac	cttgatgctt	gtatagagct	aaaaagtcat	51300
taagacacca	actctgagga	ataagctcct	gagaatgtgt	tgcatctgtg	agtttcagtt	51360
gcatagctag	tgtcatagcg	agtggataga	cgttctctgt	gcatgtccct	acaatgcttg	51420
tg agttatga	caacactgtg	tacgagcaac	atagtttctg	cagttgaaaa	gtacgaattc	51480
at agaatgta	aagagatagt	gtctatatct	tttgactgaa	aacagaaaat	gagatataaa	51540
ggaa taagac	ctttcgacat	gaaagtaacc	ccacagttgg	aataggctag	taagctttcc	51600
<pre>aacatgcagt</pre>	tttgaagctg	agaaagacgg	gtcctctcat	cagggtgctg	tggaagatga	51660
tagcacactg	gggggcgttt	agagcaggtg	agtgctgttt	tcttccaacc	cagtttttct	51720
gccactttct	tatgtttttg	tgaaggtaat	tttaaaagca	gatgtctaaa	agatgtt tg g	51780
tagtgatggc	attactgcat	gtgtcatcag	ttaaatgaca	gctcgggagc	acagcagtta	51840
tgttcgtgtg	tatcttggga	tttttgttga	agaggaaaaa	ggcagttatg	ttcatcatgt	51900
aggtcaaact	ttaatgccaa	tactggccaa	tattcttgca	aatgacagcc	atgtaaaatc	51960
agggcatagc	tataaaatgg	gaacggtgct	cacagctggc	ttctttgtgg	tgagga ca gc	52020
ta taattggt	gaggcaaaac	cagtgtgcca	caaaagcaga	atacattctg	ctgtgcaagc	52080
aatgaccaga	cagactagaa	tgaaaaggca	agagtttcct	aaggttacct	ggaacccctt	52140
gccaggtgtt	gcattaagtt	tactggccct	tgccaacatt	cttctaatgc	ttcctcattt	52200

•			40			-
catctggctt	cttggcagtg	ttcagttttt	gtggtctttt	atttttactg	tttgacttca	52260
tttctcttct	tagctctgta	aagttccaca	tgtgtttatc	tttgtggtga	aaacacaata	52320
aacttgctta	atataatgtt	ggaagtatta	atccattgta	ttagtgtgta	caggacctgg	52380
attgctgata	aaaaaataac	tagcaataac	agcctgattg	cttaaaaata	tttagtaagt	52440
tttgtcgggg	tggattgggg	cagggcagaa	cttttacatt	aaatatagat	gcaagatttg	52500
ataagaatca	gccagagtgt	acagtaagta	ttcacttaat	gttgccaata	ggttcatgga	52560
aactgcgaat	ttaagcaaaa	tgatgtataa	tgaaacaaat	tttactaagg	gtttattgat	52620
aaaaacaaga	gttaagttcc	tatggcatat	ttctgggcac	aaaaacatca	ccaaacttct	52680
aaataaagac	ccaagacact	tctaatatta	aatattgatg	taaacgtgag	atatgcaaac	52740
atttaagcaa	gattaataca	aatatgataa	ttattggctt	ggcacagtgg	ctcactcctg	52800
taatcccagc	actttgggag	gctgagacag	ggagatcacc	tgaggtcagg	ggt tc gagac	52860
cagcctggcc	aatgtggtga	aaccctgtct	ctactataat	tacaaaaaaa	aattagccag	52920
gtgtggttgt	gcacacctgt	aatcccagct	acttggaaga	gtgaggcagg	agaatcgctt	52980
gaacccagga	gacgaaggtt	gcagtgagcc	aagatggtgc	cactgcactc	cagcctgggc	53040
aacacagtga	gactccatct	taaaaaaaa	aaaaaaaga	aagaagtaat	tatttttcca	53100
cttattccac	ttcagggtct	cagggggcca	gaacctatcc	ctacagcttg	ggatgcaagg	53160
caggaaccag	ccctggaccg	aatgccattc	catcttgggg	tgactcacac	acacactcag	53220
actgggacca	tgtagacata	ctgattaacc	taatgtgcac	atctttgaga	tgtgggagga	53280
aactggagca	cttggagaaa	acccacacag	acatgaagag	aacacaaact	ccacacagat	53340
aatggccccg	ggctaagaat	ccatttttt	cttgtcaaca	ttataagaaa	gcgacattga	53400
gcataaagac	attatttgag	gacctgctgt	actatgtact	tagagagata	ggcattctat	53460
cttgagttcc	tttttttt	cccttcttga	aggaaggtta	aattgcatct	gagatggctc	53520
ttgaaattga	tcaggggttc	aagctgactt	gcatactctt	tgggaaagaa	tttagaagga	53580
tgtgtatgag	gaagttctta	tggttaagcc	tgtttcctga	cttgaataga	tgaatcaaat	53640
attttttact	attctggaag	catcgcattc	tggaaagaac	catactatgt	catctcagtc	53700
tacctcactc	cattgtaggc	acttggaagc	tgaagttgtg	atttctccaa	aattagatag	53760
ctaattttca	ttggtgttág	aacaaaaagc	gctgcctctc	tttgaagaca	ccagtcctcc	53820
accgtcctcc	tctgcaaggc	cgttttcccc	cccttttt	tttttttga	gacagagttt	53880
tgctcttgtt	gcttaggcta	cagtacagtg	gcacaatctc	ggctcactgc	aacctccgcc	53940
tcctgggttc	aagcgattct	cctgtcttag	cctccagagt	agattacagg	cacccaccac	54000
cacacccggc	taatttttat	tattagtagt	agtagtagta	gtagagatgg	ggtttcacca	54060
tgttggccag	aatggtctcg	aactcctgac	ctcaggtaat	cctcccacct	tggcctccca	54120
aagttctggg	attacaggca	tgagccactg	tgtccagcca	atttttctgt	att tttaaa t	54180
gaagatgtga	gcagcctaat	gtaagatcac	aacatgtgat	tcaatacagc	cgtggcttgg	54240
tgttgacatg	ttattaccag	ttgagctaat	ccatgtaact	cagcatttta	tgctttacta	54300
agattaaaat	gatgtgataa	cattaaattt	tgaattacag	ttgatgtttt	ttatttaaaa	54360
aacatttttc	ttagttaaat	aatacatgat	ggtttaaaaa	tcaaatattc	agtgcaattc	54420
ttctaaaatc	tctgcaagtg	tgggggtcat	ttaattgctg	agcctcccag	cctattagct	54480
ttccattctg	agctttcaag	agatggtggc	agctggcaag	gcagttttgt	ctgggaaagc	54540
cattgttaac	agagcagaat	tggggatgga	gcagccatag	cccacccacc	agagtaggca	54600
caaatcagac	ctgaacgtta	tcacaaagtc	caagttggct	cagacatttg	tgttaaatca	54660
taataaatat	tttagagaac	ttggttgcaa	atttacattt	gatctcagtc	agtcctcttc	54720

•			41			-
ccctatctct	acaagcttac	aaaccgcatg	ggtgtgtggg	ggtcttattt	aatattgcga	54780
acagctggtt	cctgtatctg	aagttcttgc	cctggagcct	gggtgtttgt	tgtagttctg	54840
caccatctgc	cttggttgat	aaggcatttt	ggaggccact	gattttaggc	agcagtgttg	54900
ttaggatacg	gaaacagcag	gatgtttgtg	gattgagcct	tttcagctga	atcttctggc	54960
cagttctttc	tggctgtgtg	aagttgtgtc	gactacagag	caggatgete	atgttgcctg	55020
ctgggctctg	ttagggtggc	cagacgtgct	tgtagcagcc	ttactgccag	aggaacgtac	55080
gttggcatcc	agagtccagt	gctgccgcca	gttgcagtgc	agcaaggcta	gccccaaacc	55140
tgatttgctg	caaggattag	ctcaactcta	gtgacattta	ttgtgttttc	tcatagccca	55200
aatcacagcc	aaaaaaaaa	aaaaaaaat	ctagggttga	catttttaaa	aatttttta	55260
aaaaacattt	ttcttggtta	aataatacat	gatggtttaa	aaatcaaata	ttcattgcag	55320
ttctaaaatc	tctgcaagtg	tgggggtcat	ttaattgctg	agecteceag	cctcttagct	55380
aaaaaatcta	gggttgacat	ttttaaaaat	gtattcaaca	gagtacgagg	gaaa agatt a	55440
aagatggtgg	atggaaaacc	ataaaagctg	agaggaaggc	agcactgggc	ttagag tca c	55500
ttggcttccc	tctagctagt	aaataaccag	caccaaatca	cctgatcctc	ctgaacttca	55560
gtttctgtgg	ccatgaaata	agaggttggg	tccaggaatc	aatgtaaatt	gtcaatttaa	55620
catttccctt	tattgatatt	actccccct	gggcttgata	atttagttat	aattcttcat	55680
gcagctttag	gttgagtaag	tttggtggga	aacagtagct	ctcttcatat	atttgagaga	55740
tgtcatttga	aaggggtaga	tttattcagt	ttaactccaa	gaagcagaaa	tgggacccat	55800
ggtagaagct	accaaatgga	ggtttggctc	taaataagaa	aacgatcttt	ggagtgcctc	55860
tectagttta	gatgaaaaaa	attgcatcaa	gttgtaacca	tgctagtcat	tgggaatttt	55920
attaacaaca	cgtagctcct	gtcctgggga	ggctcatagt	ttgatagggg	taagatggaa	55980
agaattgggc	agatgtggat	tatgtcttag	cagtagagcc	aacagagtat	gttgggggtg	56040
aaggggtaag	agaaatcaca	tacctcctag	gtttttagca	ttttccaaaa	tgaggaaaat	56100
gggtagaggc	atggacagtg	acttatattt	agacgcgtta	agccagttgt	aactgcttga	56160
cgtctcagcg	ggataacaag	taggcagcca	tgttgtgtaa	tggaaattcc	atagctgtag	56220
	tgcgctctgg				-	56280
gcgcctgaag	ctcgattatg	aagaaattac	tecetgtett	aaagaagtaa	ctacagtgtg	56340
ggaaaagatg	cttagcactc	caggaagatc	aaaaattaag	tttgacatgg	aaaaaatgca	56400
ctcggctgtt	gggcaaggta	agcttcattg	ggaagcatct	agtcaacctc	acccctcatt	56460
ggtgattggg	gagaagtgtg	gaattaaaaa	aaagtcaagt	ctaattttag	tggccatctc	56520
	atcacatctt					56580
accacctttc	atgattcctc	ttaattaaaa	ttcccagaag	gccgggaaat	aggaagaaga	56640
cagaaaaacc	caagggtttt	gttgcctata	aactagataa	tgatttgatg	atatactttg	56700
	taaactagaa					56760
acatagtgtg	gggagcacag	acttcagaca	ggtggtcttg	ggcttaaatc	tcaggcctgc	56820
	gcagtgtgat					56880
	atgtgggtgc					56940
	ggccttagag					57000
	tactggtcca					57060
	caatgtgaca					57120
	tttaaatttt					57180
atcagcctgt	agtagattgg	aaaattttta	aaaagaaaaa	aaattgatgc	ttcacagata	57240

Ī

4

H

San Tank

1 CETTO 1

gtttgagaac cgctattttg aagcttacct tcagtcatta ttagtgttct agtcaaacaa 57300 tgatttettt aaaaatatat gttaatgtet tetggeaaga gtaaaageet gagtetaate 57360 tgattctatg ctactgagtt ctggttgage teatcatgaa taaccaggtg ttetgaataa 57420 gggtttcaag tatgtataga atgggttttt cctgagttta tcagttgtgc agtgggaaaa 57480 egttgtatat geacttttte ttttttgaga tgtagtttea etettgttge ceaggetgga 57540 gtgcaatggc gcgatctcag ctcactgcaa cccctgcctc ccaggtttaa gctattctcc 57600 tgcctcagcc tcctgactag ctgggattac aggcgcccgg caccatgcct ggctaatttt 57660 ttgtgggttt tttttttt ttttaagaca gagtcttgct ctgtcgccca ggctggaatg 57720 cagtggcgtg atctcagctc actgcaagct ctgcctcccg ggttcacacc attctcctgc 57780 ctcagcetec tgagtagetg ggactacagg tgecegeeae catgeceggg taaatttttt 57840 ttgtattttt agtagagatg gggtttcact atgttagcca ggatggtctc gatctcctga 57900 cctcatgatc cacccacctt ggcctcccaa agtgctggga ttacaggtgt gagccaccgt 57960 gcccggccaa ttttttgtgt ttttactaga gacgggtttt cactgtgttg gcaaggctgg 58020 tettgaaete tggaeeteag gtgatetgee tgeeteggee teecaaagtg etgggattae 58080 agatgtgagc cactgcaccc ggcctgcata tgcatttttc atctctagga gcataaatgg 58140 aacaaagcag tgttttttac tatagttttt taggcatttt taaccttttc tgaattttga 58200 catcaatttt agtaatcatg ggaagttatt gtttgttacg cattttccct ttctatggat 58260 aaggaaactt gggcttagag cagttgaata gtggcttagg gccacagagc tgggttcaca 58320 ccaccgtact gcactgcctc ctgttgaaca ggatctccag gtgcttatct cagaacacgt 58380 atgcagtggt gaagaccgaa gttctggatg gacaccagct ttcagtgtga ctttagcagg 58440 taccctettt etgggetett geceeettae tgatagaagg agagaettge actgagtaga 58500 ggatettgga getgtettgg agttetaata tteettgeac etgtaetttt tettgaggtt 58560 tacctttaca ccaaatgacc ccaaattgct gttttgaaaa gggagaaagc agagaaaaga 58620 atgagtetgt tettececca tteacagttg cetagatgat cacetteagg tgtetttget 58680 totgogaaag gcaaattgca tgggtotgtg acagotatto caaatatttg agottottag 58740 aagcctggca cctggatatt tgtttttcac tgggcatatt ttgtgggggc taatagaaat 58800 actotaggaa tetggaceet gggtagtgaa agttgggcac agatgattga gcattetgta 58860 tactggagtg agctaaggct gacctggaat ttccttatgt gttgcctgac tttgccacat 58920 caetttttac tgcagaaget etaaceataa agggggettt gtcagtcagg tggttttaac 58980 acattaagat ttaacaactc caaacaaatg agggcggtct attttgtggt tcagaataaa 59040 aatgtgaatc aaaaaatttg agcctaaatt tgaatcatat ctttgacctt tgaagtagag 59100 gccaactcac ctcagagacc ttgtaagaga ggacagttgt gtggattaag aggcccttcc 59160 tcatagtgac ataaaagacc ctgaagtgat ggaaataaag gaatttataa aattttccca 59220 gttaaaatta gatgaggggc caggattagg gtatcaattt aggagaagat aacataatcc 59280 tatgacatta tagataatto agtttagtao acatcaaaat gatttotota aagatatota 59340 gatagaacct tataagctgg aatgtctttt ttaggaatgg gattgcagag gggctgcctg 59400 ggctgctgac agtaggggcc agatgcaaac tctgcttgcc tttgacccgg caatgccatt 59460 tataaaaact tactctagaa actaatcagc caaaaatgta ctgcagtaag aatgcttatt 59520 gtgacattgt ttaatagtca aaacaaaaca aaaaacccaa catgtgacta tcccatgtca 59580 tattccttga aaatgacacc ataagtagat ctgtatttac tgacgtaaaa gatgtctaag 59640 ttgttaaatg aaaagagtac agcatggtct cctgtactgt tgatatttcc atgtgcgtat 59700 acatggaaaa acacccacga tgcagatgtc caggttatag acaggatgac catagggccc 59760

-			15			•
aacctggcat	agccctggtt	tatgactcct	gtcctggcaa	aattattaat	agccttcctc	59820
ctcttcactg	tcaaaagctt	cctgctttgg	atggtaaata	tatgcttatt	ctagttatgg	59880
gtggttttc	actttcttct	ttatacctct	tgcatttcag	aggtttttt	gcaccacttt	59940
taaacagtga	gtgtata tt a	ttttttaagt	gagtaagaag	ctatttacat	gggggatgga	60000
ggaatggcct	cctgccctcc	cagaccctgc	ctgcaagccg	taggtgggct	ccactgccag	60060
gtttctcttg	ggttaggagt	gaaggcagca	ccatggtggg	gaagggcatt	ccaggccatt	60120
cttagcaaaa	acattgggtc	caacctgcat	gatcctgtgc	tttaaatcac	agaatctaag	60180
cttactcctg	aataccacaa	tatctggtac	tgtccagtga	cacagccaat	attcttttct	60240
ttcaaaaaat	aaaggtctga	taagacaatg	ggaatgattt	agtaatagga	aattggacat	60300
ttcataactt	gggaaaattt	cccagtttga	gaaaaagtat	tttgtgaaaa	aaagccccac	60360
tataaatcac	ttatcatgct	gactgttttc	tagcccacat	ttacttctca	tcagcatttg	60420
aagtatttgt	ggggagggtg	tgcgtgtgtg	tgtatgtacc	caggatatat	ctatgagctg	60480
gaatagcaga	gggagacaag	aaatagaata	atagtagaaa	gcagagatca	gggtatattt	60540
gcttcctgtt	gctaccataa	caagttacta	caaaaatagt	gactaaagca	acagaaattc	60600
ttctctcagt	gttctggagg	ccatagetee	aaaaccatgc	cgttagcttg	tctgtacttg	60660
gcctcttcca	gcttctggtg	tctgtcagct	tcctagactt	gtggtcacgg	cactccagcc	60720
tetgeeteet	tggtcacatt	gattcccctt	ctcatctcct	cctctgtatg	tctattataa	60780
gaatgcttgt	cactggatat	agggcccatc	tggataatcc	aggatgctct	cctcctccca	60840
aagtccttac	ttaattatat	ctgcaaagaa	ggtaacattc	acaagctcca	gggattagga	60900
agtgaacaca	tctcttttga	ggggacacca	ttcaactcac	tctacagggt	cattatatta	60960
		aaggtatagg				61020
ttcttctaca	ggccaactta	aaaaaaatga	tacgtgaaag	ggaaagaaga	aagtacttac	61080
tacacagtaa	gtatttccaa	gaggtggccc	agtgagactt	ttgaatctgt	taataaaatg	61140
attactattt	ggttcaaatc	cacagatggt	tattttatca	ttaattgcaa	gataggaaca	61200
caaaatattt	tttctctagt	ccccatttga	gtagcagcct	tgtttgacat	ttctgacatg	61260
gaggacacca	agagaaaatg	gcagtcagca	tecetggget	gtcactcacc	ggcctaatga	61320
		ctcactgcct				61380
ttcccttaga	gggttatgag	tatggtatgg	gccaatacac	ataacgtgca	tggaatggcg	61440
		aatcagtgct				61500
cttttcccaa	aggtggccag	agacagaaac	cagagaaacc	atccttctgg	acaggctgtc	61560
tgagtggcag	ggcagggtac	aaagcggcca	cttttttcc	cggatggaaa	gaaagatcaa	61620
tgcctaactt	ggaggcttcc	tttctcccaa	aagacaagaa	agacttggca	tcttattctt	61680
cagtcttctt	gatatacaca	tttccacctt	tttggccttg	taatagctga	gtaatgagct	61740
aaagaatttt	ggttcaaaac	tgtcaccttt	taaaattagg	tttgccctaa	ataacatcct	61800
tgactttaag	agaattttct	taagttttag	acatttttaa	tcactgtgag	tattcaaatt	61860
aatcacatgo	: aaagcattag	ttagaggctc	ttggacattt	tctgtttta	gagctttgtt	61920
					gctcttcttg	61980
					tgattagett	62040
					ggggcaggtt	62100
tettgeegte	aattttgcct	ctcacaataa	cccttccago	ctt cttg cca	getgetetet	62160
					: ctctcttgta	62220
cttttctctt	ccttttccct	catccttaag	g gctcggctca	ı aatgaaggat	: tctgtggaac	62280

The first first transfer from the first transfer from the first from the first first from the first first from the first from

cttgattgct cagttagaaa tgagcaaact gtcgcaagga cagaaaacca aacatcgcat 62340 gttctcactc ataggtggga attgaacaat gagaacactt ggacacagga aggggaacat 62400 caaacactgg ggcctatcgt ggggtggggg gagcggggag ggatagcatt aggagatata 62460 cctaatgtta aatgacgagt taatgggtgc agcacaccaa catggcacac ttatacatat 62520 gtaacaaacc tgcacgttgt gcacatgtac cctaaaattt aaagtataat aataataata 62580 ataaaaagaa atgagccagc ttctctttca tctgagctct acttcctttt gattctctct 62640 gctttctgag atcacatctt acatgacaat ttttcatact tggctttatt tccctagaat 62700 gttgttaatt ggcaccaggt tggagctcag gtcgtatact ttattccttg cagagtctga 62760 cagggtcaga acatgataac acatttgaga agtgagaaga agggaggaag gggccaggga 62820 agtgagggga gaataggggg tggaagtagg ggaagaagca aatagggcaa ggttttagtt 62880 gectecette tgttettatg etgttaatta ataatggaac cagtggecag geatgatgge 62940 teatecetgt aateceagga etttggagge tgaggeagga gtategettg ageceaggag 63000 tttgagacaa gcctggacag catagtgaga ccctgtctct acaaaaataa aaaaaaatt 63060 agccaggcat ggtggtgggc acctataatt tcagctactt gggaggccga ggtgggagga 63120 tcattggagc ccacaaggtt gaggctgcag tgagatgtga ttgtgcctct gcactgcage 63180 63240 aatggaacat gaaacacaat tcatttttat tattaagttg tattctgtgc ataaattatt 63300 tccatgtctt ctctcccttt taaaggtgtg ccacgtcatc accgaggtga aatctggaaa 63360 tttctagctg agcaattcca ccttaaacac cagtttccca gcaaacagca gccaaaggat 63420 gtgccataca aagaactett aaagcagetg aetteeeage ageatgegat tettattgae 63480 cttggtaagt ctgtgccatc gattggagat gacaatggaa gtttcactca catgaaaaat 63540 ctgaagagac tgtccaagtt atgtattgac ctgcctttag gtttagcaat caaaatttac 63600 tactgagact tttaatttaa aaagccctag ggtaatcaca aatgtcatct tcaagcatat 63660 aaaaatctct gtattttcac tggggagctt gttaactttg cttggcatgg agggagggtg 63720 ttcattaagg ctgcagtcat aattgtggtt cagtccagta actcaaatat tgataggagg 63780 tttttacagt caacegaagg aacatectgg aaaaegtata gatgttcaga acegaggett 63840 ggtttaatta caggagccac teestegttt ttactgetca caaacagaat teatcagaaa 63900 aattgtagaa agcagtttgt gtgtgtgcct tgaatgattt tattttggaa actgggtggc 63960 accttgtctc ttgaatagtt tttaaaataa gaagatggga acaatataca gtcagccctc 64020 catatctatg ggttctgaat ttggggactc aaccaacctc agatggaaag tatttgggaa 64080 gaaaaatcaa tgaaaactaa acaataatat agattttaaa atatagtaac tatctatgta 64140 gaatttacat tgtattaggt gttataggta atctagagat gatttaaggt gtgtgggagg 64200 atgtggccgg gcacagtggc tcacgcctgt aatctcagca ctttgggagg ccaaggctgg 64260 tggatcatga agtcaggaga tcgagaccat cctggctgac acggtgaaac cctgtctcta 64320 ctaaaaatac aaaaaaatta gccaggcatg gtggtgggcg cctatagtcc cagctactca 64380 ggaggccgag gcaggagaat ggcgtgaacc caggaggcgg agcttgcagt gagccaagat 64440 64500 aaagtgtatg ggaggatgtg tgtaggttat gtgcaaacat agcaccatct tatagaaggg 64560 ccttgagcac cgtggatttt ggtgntctgt ggggactcct gcaacctatc ccccgaggat 64620 gccaagggat gactgtattg gatagatttg cagttgccac tgtgaaggac ttgttgaact 64680 ggggtgtgat tatgatgcac agagggccct cctgacttgt cagtggccat gcacagggcc 64740 aggtggcaat gcactcccgt ttgcctgccg cctatcaccc aagctgctgt ctctactggt 64800

()原排機構(對)對於公司持續(計)

-			43			-
ggtgagctgg	ctcgatgtgg	taggagatgg	gccctgctgc	ttttagagca	tgtggccctg	64860
cttccagaat	acctgttctg	gttgcagctg	ctgctgctga	aggctccaca	gaacacacag	64920
tgctttgggg	ccctgcggtg	gcccggttct	ctgattgttc	ctgcagccac	gacagaggat	64980
gcagtgtgag	ccgcatcagg	cagtatgaag	tcctttcctc	tcaagccacg	tagctagcct	65040
taaaggttaa	tttcataacc	cttaaggtta	ttttttt	ttaattttt	tttttgagac	65100
ggtgtctcgc	tctgtcgccc	aggctagagt	gcagtggtgt	gatctcagct	cactgcaagc	65160
tccgtctcct	gggttcacag	cattctcctg	cctcagtctc	ccaagtagct	gggactacag	65220
gtgcccgcca	ccatgcctag	ctaattttt	gtatttttag	tagagacggg	gttţcaccgt	65280
gttagccagg	atggtctcaa	tctcctgacc	tcgtgatccg	cccgccttgg	cctctcaaag	65340
tgctgggatt	acaggcgtga	gccaccacac	ccggccccac	ttaaggttat	tctttagctt	65400
gaacatcatc	tctgagaaac	tttccctgac	tgtggtctcc	tctcccacct	caagactgga	65460
tgaggtgtct	tgctaagccc	cctgtagcac	cccacactct	ccccatggtg	cgtatcacat	65520
ttctcatcat	caccgttatc	tgcttattat	catcactgct	gctgcctaac	ttcaccttgg	65580
gccaaatgtt	gtgcaaaggg	acttaaactc	ctttcttaa	tccttacaac	atgatcaggt	65640
agatgttgtt	ctgtttctct	ttagagttga	gaaaatagaa	acagacaggt	tacgtaactt	65700
gctgaaagtg	acacagccga	tttgccgcta	atcagtgtga	cttcggaagc	tgcacttttt	65760
tttcaacttt	tattttagat	tccaggattg	cgtatgcagg	tttcttacaa	aggtgtattg	65820
tgtgatgctg	aggattggag	tgtgattgaa	cttgtcaccc	aggaaccaag	catggtaccc	65880
aataggtagt	ttttcaaccc	ttgccttcct	ccctccctct	ccacccccca	ggagtccctg	65940
gtgtctgttc	tcatctttat	gtccatgtgt	acccagtgtt	cagctctcat	ttctaagtga	66000
gaacatgtga	tgcttggttt	ctgtttctga	attagtttac	ttagggtaat	gacctgcagc	66060
tgcatccatg	ttgctgcaaa	ggacatgatt	ttgtcccttt	ctatggctgc	agagtattgc	66120
atggtgtcca	tatatcacat	tttctttatc	cggttcactg	ttactgggca	cctgggttgg	66180
ttccatgtct	ttgcaattgt	gaatagtgct	gtgatgaacg	tgtgagtaca	tgtgtctttt	66240
tggtaggatg	atttattttg	ttttgagtat	atactcagta	atgggattgc	agggtcgaat	66300
ggtaattcag	ctcttagcag	aacctgtatt	tcttactcca	cctcccccgc	ctgtccttag	66360
tatacagcag	tggctcttta	ttgccttttt	cccttatagg	atacagccct	ctgcggactg	66420
ggctggggct	gtttggccat	tataccctcg	gcttctagga	cagtggctgt	gacacagcag	66480
atgctcaaag	aatatcttta	agattcagag	tgtgagacac	tgcactagca	ccgccatctc	66540
atgggccctc	acaacagccc	tgggaaggtg	gcctgcaccc	tctctaagaa	atgaagaaac	66600
tgaggtcaca	tgttgaccat	ggtcacaaag	tcacctgagg	ggaggtgaca	ggaactgaac	66660
ccactgtcac	tctgtgtttc	cctgggaccc	tctgagcgca	ggaggcccgt	gttgctgtgc	66720
agtggcaggc	caaggcaatg	ccttggtgga	gctggggccc	atttggccca	ctgacctgag	66780
gaaagcagtt	ttgtgaattg	gcagtagctg	catttgctga	catggtgagt	tacaggaaat	66840
gccatcatgt	tcctatcatg	tgaaacaaag	tgagaaatag	gttcag g gtg	ggaggctgaa	66900
agggaggaat	gcagacagcc	ccgctcccca	cacttgetee	aaggctgggn	aggaggaacg	66960
ggaaggtgtc	teceetectg	gattcagtca	ccttcttctc	ttcattcccc	tgcagtatcc	67020
			tgcttcttgt			67080
			ga ac aagggc			67140
			acggcctccc			672 0 0
			ccggtccact			67260
agtcctctcc	atccctgcct	ctccccaccc	tctccctctg	nngccccctt	ctctgacagt	67320

getgacecec etetetette eccaetettt eccatecteg eetggeetee ggtttggatg 67380 ctgtccacac acttcccgag ggcctgagag gacctccgtg tgaggcaatg catttcccag 67440 gteacetetg tgtetetete eaggettttt ceagggaete eeeggggtea gteteetete 67500 eccaetggaa eggggaaaet gggattggee tagaeeegge agtggagtee eaggtgeeet 67560 geetgeeegg etgaeteege ceagggagge eteccaeaga ageteeteea gaeteeaeet 67620 gttacctccc ccactcctct cacccaagge tgtgctgtgg ccaagtcagt tgtttagtct 67680 acactttctg tttagtctac accatggcta cctcaaggcc cagtgaaggt gtgtagtata 67740 aagcaaaatc aaatccatat ttcagttttc cttaaaaagt gaccttcata ttctggccag 67800 aagaacagaa tggttggttg gatatatttt gagttttcat gggtttttgt tttcctgcct 67860 cttgttatac tttctgaaat tggcttttag tctaaacagg ttttttttt tttttttt 67920 tttttnggca atgtgttttc ctccaaagag taagaataat aggcctcatg gctgggtcgt 67980 gttctacagt ttgtgaatat tttctcaacc tttgtcaaat ttcatcttta cacatcctgt 68040 gtgaaattgg gcacgtgccg ttatttccaa cttagagggg atgaatgagc ccttaagagc 68100 ttgagttete tgeecacatt gegagttaet cagtgecaga aggagttetg gaacceaggt 68160 ctcctcagtc tccataccac atcccttcta gggcaccatg ttgcttctgt gtttcttggc 68220 tetgeceact ceatgecage acaactetee ceacecetge tittggtggaa teatgtteet 68280 ttggggtaga tcacaccagc cagaggcaac tgctctcagc ttagcagatg gtactcatca 68340 cattattett gaageettgg gteaggagee tgeeecacee atetgeatee atttgteeag 68400 ccctcagaca attgccactg ttttcatgtc tattctttga ctctctatcc tgggtagaca 68460 acatggactg cccagcatcc tgtcttctgt ctggggctcc cactgtcgtc ctgaccacgc 68520 tgggggetge cagtgacact gggaaactee egagggacee cetteagget teacateate 68580 tgctccctcc ctagcatccc agcctagaac actttccagc catcagctgc attccccagt 68640 gaggegtgea geeteteeca tgataggagg getteageeg aaagaacaet teaacaggee 68700 cagaaaccca ggagcaccat tagatcagaa agcagaagca agaatgcatc taatctcccc 68760 cacatcaatt gctatagttt tattaatctg catattatag gtcagtaagg ggatggcaca 68820 68880 gtttataatc cctgcaagag tctgatgatc ttttggtgac cagaagtgcc attttttgat gggcttctag agatcctcca-tcagggatac cagacatgtt tggcatgcct gtgctgccgc 68940 gagacgetaa gegtgtgtee agactacaeg tgtgggteat gggteeagea geagagetgt 69000 catattgatt gtttgcttct actaaatgta taaagcctgc ctggtgtcca gaagaaaaga 69060 aactataato caatttttta gaatocataa aaggtaagaa gtaggagaac atttagaato 69120 cacaaaagat gagaagtagg agaacrgttg gattttttag aatccataaa agatgagaag 69180 waggagaace tecaaaagga aggaateage tgagagtatt gaagatgace aagtacaaac 69240 aggeagaggg gagegettee cetteteete teecaggegg tgggetgeet egeteggeea 69300 ggacacacag agcagcatcg tgcrctttga ggggcaggtg gagctgctca tcactagcag 69360 gggtgctggc ggggaccaca gtgttctctt ccatctttga gttgaagtcc tgtgtgagaa 69420 atgagaaacc ttcatggcaa aagacagaaa gggacctaga atgtaacatt cagcagtctt 69480 gttatctcac gcacctgtct gtccagttgg ggacgttgct gtatggaggt cagttgaaca 69540 atcacagttg aggagectaa tgaattettg caccaccage cacacacatt attetgaaga 69600 69660 gtgagccatt gtctctgatc ttatcaggat cacatcgtgg gatcatattt atttggtcat tctgaatata ccctttaagt ccaaagtgaa ataactaaat gtcgttgata aaaggaaaga 69720 69780 ataaagtggg gtatgatttc ctttcacaga ggtctggaat cttcctgcct ttttcaagtc agteggtggt getggcaaat gtttaataac eageteetet eacceeteag aggaageeet 69840

•			77			-
tggtgttcag	tgtttgcaga	tttccattgt	gcaactagtc	ctcccacacc	ccattttaaa	69900
ctacccactt	gatgtcactg	gtcatggagt	tgggctcaca	gagccagtgg	gagtcaactg	69960
gagcagccac	tggactcatt	caagtgtttc	ccaaaacaat	ctgctcctag	aaggactctc	70020
ccttaatctc	ctaaccctgc	cattcaggat	gattccctgc	actctgggaa	gcacacgttc	70080
tagtgggaag	actgatactg	ggcaactgat	aaccaagtga	cttaaacttc	tgagggttac	70140
aaagggtgtt	tgtatcctca	gtgtctcatt	tcagattctg	ctcagagcta	aatgcaacaa	70200
tgtgagaaga	tgttagtatc	ccagatette	atccaggaag	gaatcttaga	gatcattagg	70260
ttgtagggtt	tctcttctgc	agaggagata	gagggtcggt	gtcagattgc	tgqtttgcca	70320
gtaccactcc	ctggagaaaa	gagcaaaaga	aagaaacttg	ttagtcaact	gtgcagagcc	70380
accgtgagac	tgaatagctt	tgtgggtggc	cccgtgtttg	ctgcaagaga	cctctggcct	70440
cttgtagcag	ctgccacatg	gtaaacagag	ccgagatatc	aggagtctcg	ctgaaaatgc	70 500
agtcagatgg	gctctgaata	gaggaaggca	ggacactctt	gagatgggat	ggggtttctc	70560
acagcaccgt	acagggacca	cctgcaagat	ctcttgaggg	gcttgtgaaa	aacacatccc	70620
tgaggtcacc	attcttgacc	tgctgcttat	tgagtttctg	atgcctggga	tgtgcaggtt	70680
taacaagccc	ccagatgatc	ctaataggat	tcctgcctga	aaattgctgg	gtgaaggete	70740
ttccccctcc	aagtgataaa	gaaggaaaag	attgatcctg	gaagaacatc	cgttagatga	70800
			gaattactag			70860
aattgtggat	catcttgtag	gcctttctgg	catatttctc	cacttagatc	cacaagacac	70920
atcgaatgtc	tttttataaa	ggggttttt	aatgcccatg	tttgaccctc	tccacttaac	70980
			atctgcttta			71040
			atgcagttta			71100
			ttacccagct			71160
			gctagcatgc			71220
			tcaatctaga			71280
			catagttaag			71340
			cccacggctc			71400
			atcagcacta			71460
			attatacgaa			71520
			: acgtgacctg			71580
			gttagcctag			71640
					tctgtttcca	71700
					g attagtattt	71760
					a aaaattatta	71820
					ctacattgaa	71880
					tcattcattc	71940
					g atctatagta	72000
					c tccaggggtc	72060
					a ggaaaacagt	72120
					a gatgggtgca	72180
					c tgttccccag	72240
					t tgcatttggg	72300
agtctcatag	g aacacactg	g aagatgtgt	t ttagcttgg	c caggttcat	g caggacagat	72360

-						•
tttctgcata	aagaaaatca	atgacagttt	ctgaaactgc	atcctggaag	ccttgaccag	72420
tttgggtaat	aacaagagat	ttgaaagtgt	ggggtgtaca	ggtgttttgc	tgaatctagg	72480
tggtggtggt	gattattatt	atttggaatt	cagctttcag	ttctacctgc	ttg tgagttc	72540
caaactttgt	gaaaattagt	tgcttggacg	aaacttttct	ttgcctctgg	aaggctgtca	72600
gaaagcgaga	tttcccagct	tatgtgcagt	gttatagtta	atagagtaat	ggctctgcaa	72660
agttgttcct	ttactttaaa	tgtaatttat	tttgcatttg	tgctacagaa	cggtca taag	72720
tgtgcccttt	tgtcctcttg	tttggaaact	gggtttttat	aatgtgtgtg	gtctatccga	72780
agattattgc	ccattattga	acaccattca	tagcaaccat	ttgcattagg	cattgtacgt	72840
gtactctcca	ctctgcaaac	tatgtgttct	gtcccttttt	aaaaagagga	agctaaggtt	72900
cagagaagct	aggtagtcca	ttctgagctt	cacgtgccag	aggccatttt	gtacttactt	72960
caaatgccat	tgaaataaat	gcacatcaga	gaattgttct	tagcataagg	ggcgctacat	73020
gtaacttttt	attagtgaaa	tggatgatgt	tcaagggctg	tggttgatta	gaaaggcgtc	73080
cagaccctgg	ctccagggac	tatggagcag	aactcgaggc	cagtgcctgt	cgagcgggtc	73140
cccacactcc	atctgtgtga	cctgactgtg	gatggcctgg	ctctgccgtt	agattgccac	73200
ggtgccctcc	tctggttgaa	cctttctcga	gaagtgcttg	ttggaggctt	gagtgcagag	73260
cctgtgagaa	gctctatgtg	gttcctattg	cctgtcagct	tgctgataaa	ggtcattggt	73320
ttggcaaaat	ttggcccaag	gtttgccttc	tcataacata	ccactcggta	gcaaggctgg	73380
gaggaaggtg	gctatagcta	tttctggaag	ctgcttaggg	ggctgcctcc	ccctaaattg	73440
gtacataatt	tgcagggcct	attgcaagat	gaaaatgcag	aaccctttct	t gaaagatta	73500
ttaggaattt	caagacagag	acaacagagc	atgaagcctt	gtgcaaggtc	cttctaagca	73560
cagagccagt	gtgaccgcac	agaacacaca	cccgtgaagc	cagctctgcc	cccaccatct	73620
gaccactctt	gagtggccaa	ttagcatagg	tcactcccca	ccctgctagg	cccaccctct	73680
taggaatgtt	gtgaggctta	aataagaaat	agccactcta	caagcggtgt	caattagcat	73740
gggctctggt	ttctgtgtga	ggtagtttgc	taacatgaga	gggtatctga	ttagctaaaa	73800
cgataacact	gacagattaa	attcagaata	actaaacctt	ccctgtgttc	ctttatgcca	73860
catgactcct	gcatattctg	ctaccagcac	ctgtttgata	ccagacggag	gggtccattt	73920
gggatgggac	aggagcatéa	gcagaaatgc	agaagtgggg	aagtgctcca	tcttcttgga	73980
agctgagctg	gcaagggtaa	tggaatgaaa	gagattgtga	atatttttga	gactatgagg	74040
aaaccagtac	actggtgttg	cccagtacag	aagccacatg	tggctgttaa	gcacttgaga	74100
tgtggctact	ccaaattgag	gcgtgctgtc	agtataaaga	acacactgga	tttcaaagac	74160
ttggcatgaa	aaaagaatgc	ctaatgtctc	agtatttta	tattgattat	gttgaagtga	74220
tagtattttg	tgtatgttgg	gtgtaacaaa	atatctaatt	aaaattaact	tcacctgtta	74280
ttttctaatg	tgggtgctag	aaactgttac	atcctgcatg	ggggtcacat	tccagttcag	74340
ttgcatgtgc	tgctacccat	tgttctacac	acacacacac	acacacacag	ctgcacacaa	74400
cctagagggg	tcagagaccc	caggageeee	tgcttctggt	gcccaggcta	agcgctggag	74460
tggaagataa	agctgggagg	gtgggtaagg	aggtgagtgc	acggagctcc	aggctaacag	74520
agtggataat	ttgttctttg	agcactgggg	agctatggat	tgcttactag	cagcaaggtg	74580
acttgtgcag	ggtatatctg	ggggaggttt	actgggggaa	gag atagag g	aggcaa gaag	74640
				gtagctttcc		74700
agttcagata	acacatctgg	actgatgaaa	ttcttttca	ggaagctgag	gaagagccca	74760
				agtgaacaat		74820
cctagatcca	ccttttgcct	tegetgatee	aagcaggttc	ataattcttg	cctggg ccca	74880

•						
agcttggccc	tggctgccag	ctgcctggct	ccagatgttt	cttaatcgtt	tcaagtactt	74940
ctctgctccc	tggaaacagg	cactcccatc	agtcacattc	cagaggagga	ggaagaggaa	75000
cttgacaagt	atcagctaca	aaagcctcct	gaacaaaaga	aatcctttaa	gcctatttga	75060
ataacagttt	tttgtgaaaa	taatcaggat	gttgagagct	tttttttt	tcttttaaac	75120
tctttttgga	aggtaacttt	t gtgaaaaga	aaacacctgc	tgctcctcag	gctgtttcaa	75180
aacactgcct	atagtttgaa	agtacggaga	tatgcatgtg	gtatgaagca	tttgcaggca	75240
taatatgtgt	agtctgggaa	aagcagatcc	agagagtgct	tgtagtaagg	cgaggccttt	75300
tagctgcatt	tagatgatgc	tgggattggg	gtgggtgcag	ggtgcagcag	tggggaggaa	75360
gaactgtgtg	tgttcctctt	gagaataggg	gttatgtcta	gaggattaac	agttttcttt	75420
tttcntttt	tttttttt	ttggagttgg	agtttttctc	ttgtctccca	ggctggagtg	75480
cagtggcatg	atctcagctc	actgccatct	ctgcctccca	ggttcaagca	attctcctgc	75540
ctcagcctcc	cgagtagctg	ggattacagg	cacctgccac	cacgcctgac	taatttttc	75600
tattcttagt	agcgatgggg	tttcgccatg	ttgggcaggc	tggtctcgaa	ctcctgacct	75660
caggcgatcc	tcccgcctag	gcctctgaaa	gtgctgggat	tacaggcatg	agccaccaca	75720
cctggccaac	agttttcttt	tttcgattga	agttcagcta	tttgcaggac	cgaaggtagt	75780
tctgattact	ttcacctgta	cttccaccaa	aaaataaata	aaacaaccat	gagtaattgc	75840
tgatttttaa	ttgaaagcat	tattccagga	ataactggtg	gacttcgttt	gcagaggaag	75900
tggcaaagac	tgattgatat	tatgatccag	cttctaaaga	ttttgctgct	taatctgaag	75960
cacattggat	ttctggttca	ataggctttc	tttttttgtt	tttattatta	caactaatat	76020
gtattcttt	cacagggcga	acctttccta	cacacccata	cttctctgcc	cagcttggag	76080
caggacagct	atcgctttac	aacattttga	aggcctactc	acttctagac	caggaagtgg	76140
gatattgcca	aggtctcagc	tttgtagcag	gcattttgct	tcttcatatg	agtga g gaag	76200
aggcgtttaa	aatgctcaag	tttctgatgt	ttgacatggg	gctgcggaaa	cagtatcggc	76260
cagacatgat	tattttacag	gtatagagtg	ttccttatgt	ctttaataca	acaaaatgct	76320
aagaatgttt	cttatccctc	tccagatgtg	cctcaggagc	tttttcaccg	tcaggtaaca	76380
ttgtaatagc	tgtcactgct	gataaaggac	tctgtgctag	gcattattcc	aagcgcttca	76440
tctgcacttc	cctctaatáa	caggaagaca	ctgttcatcg	téteaatttg	tagattggaa	76500
aactgagtct	ccaagagatt	ataaattggg	cccagtcaca	cagctagcaa	gtgtcagagc	76560
tggactggaa	acccaggcct	ctctgactct	agggccttcc	ctcttgcccc	catcagccat	76620
cagatgatct	cagacctacc	tcccagcctc	tgcatctgct	cttcctctgc	ctcaccccca	76680
cccttgtcat	ctcaggttca	gctcaaatat	cacatectgg	gagaagctca	ttctgactac	76740
cctgatgttg	tgttccccac	ttccaccttg	ggcacactgc	gtcacgttat	cctggctgat	76800
ttcttttgca	caacacagcc	actgccagaa	atgatcttgt	ttccataatc	atctccctgt	76860
ctattttctg	atttttcata	gcctgtgaac	tttaggagag	ggaagggatc	ttacgggtct	76920
tggagccgag	ttcctagttt	ctgaaacagt	gcgtgggttg	aagtaggcac	cccataagta	76980
tttg ttgaat	gaacaattct	gtcagagaaa	accaaacaca	gtagcgtatt	gcaaatacca	77040
cgtgctgctc	ttgctgcctg	tcagagggaa	aactctggat	cctgcttcag	gaatattcct	77100
aaatgttgca	gcacatgttg	atatgttcat	ttactaccag	taagatacta	tgccttcaga	77160
gctctagaga	gtatcctggg	agggaataca	ttagagccaa	ggacttgctt	tgagagcacc	77220
aaattatgtg	attcaaaatc	ttttcacctt	gacctgtgaa	catggaccac	gtgaatgcaa	77280
atatcataga	aggaactcat	tcactgaaag	attttgacca	cataacactt	tccacatgta	77340
ctgtgaggtt	cttcctacat	tccctttatt	aactttaaag	acagtggtca	ccaggcagtg	77400

-			50			•
gaatttttga	gttttctata	atttatgtaa	cacacaactc	ttttggggtg	gtgcctttgg	77460
ttgattagac	agtcttcgat	atgggagagc	cacagctggt	gcttatggga	ttatattatc	77520
tgageetetg	aaaacggttt	tgttttcttt	ctctcagttt	agataggaca	tatccaactt	77580
ggtggatctt	agcggattct	gaccctctgt	aggttgttgt	ttctttaggc	tcaggccgtg	77640
gcactgctca	gatctgggct	ggctctcggg	cctctgtgag	cctgtaactc	ttggtggcac	77700
tactaggaac	tggcatgaga	tttctgccag	aatcatgtca	ttctgtgaag	ttggagttcc	77760
actttagttg	gaaaaagttt	ttatttcatc	ttaagatgca	cacttgtctt	cttgttttaa	77820
cttgccaggt	atctggatat	tccatatatt	atacaccaaa	agaaattatg	cttctcctgc	77880
ctattgagta	atttcagggg	tccagaggga	acttgctgag	tgaacatgta	caatggattc	77940
ctatggaatc	ataagatgcc	cctaattcag	tcttagtaaa	gagactggct	tcttatttct	78000
aattcctcca	ggcttgagtt	gtgcaaagag	tatgtatttg	taagagaatt	tatgaaatgt	78060
ttgcacaaga	cagatttta	gatcttctta	gtggaggaat	acaagggaac	aataaaaagg	78120
aagtggcagt	agaagaccca	gcgttagcgt	cctgggccta	cacccagcca	gtgcctggca	78180
ccagcaggca	cttgggaagc	acttgttgga	tgaattagta	gctgagctca	gtggatcgca	78240
agccaaatcg	aatgtttaaa	gttctagtaa	gtcttctctt	acacccaccc	tgtgagcagt	78300
aggcataact	ttattgctgt	ggcagatccc	taattctcag	cccttgtggc	tgtcttcctg	78360
cagatccaga	tgtaccagct	ctcgaggttg	cttcatgatt	accacagaga	cctctacaat	78420
cacctggagg	agcacgagat	cggccccagc	ctctacgctg	cccctggtt	cctcaccatg	78480
tttgcctcac	agttcccgct	gggattcgta	gccagagtct	t tg gtg agc a	ttagtaaatc	78540
tgtttgccag	aaccagcctt	ctcttattag	aggggaaaca	tttcctgtct	ctccrtggtg	78600
attcttattt	ttatacctgt	agctcttacc	agaacagggt	attgtttgat	agtctaagat	78660
tagtcagggg	tgggttttgt	gactttggag	tcctccttaa	cttctgataa	tcacggggct	78720
tecettagat	gccttcatct	tgtgggatgt	ggatccgatc	cgtgtagatc	cgatcgctca	78780
ccatgagggt	ctccctagag	cagacatttg	gaggacttgg	ctgaggagcc	acaggtgtat	78840
gtttctcatg	aattgccttc	ctcagccact	ctgggttgtg	agtattgact	gatgctgact	78900
gtgggcctct	gggccctttc	tagattccct	tggcatctct	tcctcccctt	tetettettg	78960
ccctgccctt	ggctctacác	tttctcccaa	gtcactgtct	tggagaccag	tgtcaggacc	79020
ttgagtaaca	cctccgtgtg	gatggctcgc	tctccccgct	cagccttgac	acttcatgaa	79080
ggcctcttgc	ccctgagccc	acatgtcaca	gccactgcca	ctcccgtgcc	cccgctgtta	79140
accttgggtg	gttcacatgt	aaaacctgcc	tttatattct	tgatttactt	tttgagaaca	79200
ttgtcaaagt	taggtgagtg	ttcatacaca	aagccttcaa	cctgccttca	tatgcaggga	79260
tagggctgtc	cacgtgcgca	tcaggaaccg	agtggaatgt	tgtgagcatg	gtcagttcgg	79320
gcacagtttg	ttttccctac	tgcagaataa	aagtgatatt	tttgacaatt	caggttcttt	79380
tttttattgt	aaaggaggag	gctactaaaa	aaatgatagt	tattatatat	caaatgtttt	79440
taagcatcac	ttgacagctt	aaaaacatgt	gatctttaaa	aaatttgttt	ttatgattag	795 0 0
agagcatctt	aagggaaatg	ttcaaagaca	ttgatactac	ttcagacatg	ctttgggtaa	79560
acatcttaaa	tatccaaatt	ctagaaatcc	taaaatttgc	tttttaatat	aagtgagcat	79620
ttacccttct	tctctcttt	cctttccccc	caaatactag	atttttatta	ttcactttta	79680
tctacaagaa	cctttaaaga	gtttcccatt	ttgctttact	ataagaattc	atattccttc	79740
ttttctgtcc	ctgaaaaaat	aaaatcacta	aattaaaata	gatacaaaaa	gctatctcct	7 980 0
				ccatggttaa		79860
agattaacta	aaagcctcaa	aattgtgtgc	ttagtttatt	aacaaaagag	ttacagaaac	79920

WO 00/08209 PCT/IB99/01444 51 .

-			<i>J</i> 1			•
taaaatctca	agctctaggc	tttaagcttt	cttgccaata	acttctatgt	ttttgacttc	79980
tctaacactg	gaaattaaaa	gaaaattatt	aatctacctt	ccttacattt	tctccacatt	80040
ttagctatga	ttttcataca	gggtcatgaa	gaggagtgag	gatggaaatg	gggaggaggg	80100
agcgctgttt	gttaatggtt	tgtaaacagc	tcaggcatta	aattacttgg	ttagtgaaga	80160
aaattctacc	aaggcaacca	ggctgaccac	agactggagg	gctgaggggt	catcactgag	80220
tcatctctgc	cctggggccc	caggcactgg	agctgctgct	tgcagaaagt	tctggggctc	80280
tggaagagaa	atttttcctt	cggctcataa	atgggtaaaa	agacgttaac	aaacaagcag	80340
actccacttt	ggaaatgata	gcccttctat	tgcagagtaa	tttgaagctc	tctgaagctc	80400
actctaatga	cttcataaat	caaagctgca	gcttgtaaag	gtaagatatt	tttctgtaga	80460
c tttgtaggc	agtgg tg aga	ctcggagttt	cataaacatt	atgcatagag	atgccagtgt	80520
ctacatcatt	cctggatccc	acagactcct	gctgtgctaa	gtgggtcgtt	gtccagctgg	80580
ccaagggctc	ctgggattag	aggcgggaag	tgggatctca	aggccgcact	ggcttgtgat	80640
gtcactttgc	aagaactgct	ttttttcca	cagtccatcc	catctttcag	tacttaaaaa	80700
cagaaaagat	aggtttttac	aaaccatttc	tatttttagc	actgatgact	tagagaatgg	80760
tgatggagat	agcttagttt	tatatttcaa	agcctgccat	tcagtcacta	tagtcttttt	80820
tggcctaggg	cccatttcat	tataagcctt	taagtctgga	taaactctaa	aaacatgtag	80880
aactttgttg	actaagtaaa	atatttcagt	ttgcaccacc	ttagctcata	tattagttaa	80940
taggttacct	ggcattaagt	atgtatctgc	tccttggagg	ggcggctgcc	agtgatgtgt	81000
gcgccttaac	cttcacatga	tactcacacc	ttgctgaatg	gcagttcttc	tacctggtgt	81060
cacataatgt	caatagcccc	ttcctgtatt	tttctagctt	gagtacagca	gggccctggg	81120
agagacgctt	gctctgttca	ctttctcatc	acatctacct	ttgggggaaa	aaaaatctaa	81180
aaaacaggac	ctggcttgct	cctgatggag	gaggaggctg	cagtgttcag	cctctgatgt	81240
ttttctatag	gacatgctgc	caaatagatg	agggaggagg	aggagtataa	aaactaaggg	81300
tttggcaaaa	aacacagaag	ccacctgcaa	tatagtgaag	gcttcagaga	gactttagga	81360
tgaaaataga	ctgaaaacaa	gattgtttct	gtggccagga	aaatctccag	ctattcaggt	81420
gacatgatgc	cgcgtgatga	tgagtgtgtc	cagtctgtct	gtgctgttgt	tetgcacage	81480
actgtcatca	gccttcagcg	tcccctttac	ccgttactca	tagaatgtag	cggagccacg	81540
actggaggac	cgcagccttc	cagaagaaag	ttgagaaggc	tcagccttga	caaagacaaa	81600
ggtggctgga	aaaagagatg	cagtgcaatc	tcacatagga	agattgcact	ttgagatcat	81660
gaagtttagt	ttcaaataga	gttccaatac	acagtaacgc	aataagaggg	ttgctgaaat	81720
gtcctcaaag	aaaagcagtt	cctttgtgtt	gttcccagcg	aatacagtgc	aaagtaatag	81780
agtcagctga	atttaagatt	cctatttcct	gccggataaa	acgtcttgcc	tgtttctagg	81840
tggcttgaaa	aaggagagga	gagaaggaag	aggcaggaga	aaagtcccac	tgaaaggacg	81900
tgggctacag	tgtagtgagc	taggctactg	cctcactgcg	ctgggcggct	ccaacagttc	81960
actgtctcct	agggaggttt	tcaaatgcag	gacatttgct	cacttttcca	aggagagtta	82020
ttgtttttt	gttttgtttt	gttttgttt	gttttaaaaa	attccagaat	gtaaatgtat	82080
aagataccgg	aagacaggca	aaataaaaat	aattggtttg	gggcagtggg	tttataggta	82140
acatttttct	ttactattrt	tttaaaatta	gatgtgattt	aaaaaaattt	ccaaagccaa	82200
aaaacggtac	aaatgcttta	aaggatgaag	atgttgtccc	caagtgtcat	cagacaaatt	82260
taggaggccc	ttcctcccaa	gcaaagcttc	ctgcagtcct	tccttcaact	ctgaattcaa	82320
gcacattcct	gcattgtgca	cccaaatgat	ctcccgattt	aagaccccct	gtgtctcaca	82380
gaagcttctg	gggctgaact	ttctccggcc	ttggagggtt	ggacgctttg	aatgggagga	82440

WO 00/08209 PCT/IB99/01444 52

•						
gtggtggtga	gtggagcatc	tctggcagca	ggcatttggg	agtctctggc	aggaatcaat	82500
cagcgtagtc	tccaaaggtg	gcctttctct	gacactaact	agcccttgca	ggggtcatac	82560
ccataacctg	catctcatta	acatcatctc	cttaccagtg	cactgaccta	gtgagaaaag	82620
gaacaacaag	cattcagcga	ctcctgcggt	gctccagggg	aagttagaat	tgcttggctg	82680
gggcagaggc	ccctggtgat	ctggacctgc	gtgcccccat	ttgcccacct	tetgeeetge	82740
acaaccagtg	cccctgcctt	gccagccaga	ctgtttttca	ggctcctgca	cacctccctg	82800
tattgacacc	ctattttcct	tttattcaga	gtattaatcc	tgaggtctga	cctaggaaat	82860
tttcattggt	tcttcaagca	gtcacctttc	tgtgggcctt	ttctttcctc	tttgttctcg	82920
taacaccctg	ggcataactc	taccgaacca	gaactccttg	gtgtctctgc	agcgtgttct	82980
ttgtgttttg	ctcatggctt	aatctccaga	gcctaataca	gtgcctgatg	tgtattagat	83040
gctcaataga	tgctcattaa	gttaaagtag	aagacacctc	tcagcagagt	tctcttaagg	83100
tgttgtgaat	agcattggga	aagaacattt	attttttaat	tacattaaat	acaaacagat	83160
ataataaaat	aaatcatatg	cccagtgcta	tgtcttaatt	ttttaacata	tcaataaaga	83220
gactttaaaa	cacataacac	caccctctcc	cctccaaatt	tcctttccgg	gaaagtctcc	83280
ttttggaatc	ataggaagca	cttactaagt	tgatttattg	taaaaaaacc	aagatcctaa	83340
taaatctcag	aagatctcct	gttaacctaa	agagaccact	gatgtggatt	ctgtatttgg	83400
ttgtgctgac	aaaagtttcc	cagtaattgt	ttattttaat	t gg c gtagat	gtggtactgt	83460
acctaattta	aggcacttgt	ccctctgaga	gtagagacca	agctatagaa	aatcactggt	83520
gttgtaggga	aagcctttcc	ccaggatccc	tgcaaaaaag	gtcttgattt	ttattctgaa	83580
agatgccctc	attttttgtt	cagctataaa	agttcatata	ttgaaaggag	gtctaggaag	83640
tctcactgtg	taaaccactg	aaacttcaaa	tttactttag	agttttgttt	ctggaaatgt	83700
catttctgtt	taaaaataca	tctttgttat	agtattattt	tagatctttt	tattttctgt	83760
agtggggaat	tatacaggta	gactacattt	tataaaccag	atatttcaga	ggaatattct	83820
tcaattggcc	tgccttggtg	tatgtaacac	ttaccctgaa	aagctctgat	ttcaaagaca	83880
cagttagttc	tctagtatat	cttcccagcc	tcaacaacca	gacttaagaa	ggaagtgaag	83940
gattcatctt	tcccactttc	ctgcggccac	cctgagccat	cagtagttgt	gatgtttgtg	84000
gaaagagtgt	ggaccctģag	ctgggtggga	gaagcaggct	gatictcagcg	ctggcatggc	84060
ttagggctgc	acccatctca	gctcacatgg	ttaattaagg	gttttgtggt	ggttacagag	84120
gatctcgagg	gctatcccag	ccagcgggct	cctgggtctg	tcatccctgc	ctgtgctttg	84180
ttcagaaact	acagggattc	agtttcccat	ttgcacagca	gcacccagtc	tttgcttttc	84240
tgtttcttcg	tggcttttaa	atgttatcat	attaaccatc	tagagaggca	ccctgcaagg	84300
ttattcctct	cacctgcttt	tgctttcctt	gatttgatga	aatttacago	ttctttctct	84360
cttccattat	ctttcagcca	aaagaaacag	agaaaagaaa	tactgacact	tgcctccaat	84420
tatatttcta	ctctgatttt	taaaattgtt	tttttcttat	attattatto	tagttattag	84480
gtaacctgcc	: tcagtttagt	caaccaataa	ttagttatcg	tsgctctgct	ttaaccccag	84540
gacatcagac	tcttttttc	cccagcagct	tcaactctat	gaggaaggtg	agacagggct	84600
ggggttgctg	cteggeeget	tgccttggcd	ggtgccctcc	ctcttattct	gcagtctgta	84660
tagaagttgo	atccatttgc	cagccactct	aagaacaaaa	tatggccaga	actaggaagt	84720
aaccttgaca	a gagttcttga	actcctcaga	gggaaaaatg	ttctttattc	cattatcatg	84780
ttaaaaatca	gtaaacttgt	atttaacaav	v gtacttctgc	agttgtacag	g ctgttgtaca	84840
					a gagacacttt	84900
ttcacttact	ctatcttaat	ttctgatgct	ttatctataa	a aaatctttta	a gtgtgaccca	84960

taaaaacatg ttttagtatc tcctttaaaa cccaggagca ttcctggaaa aatagactaa 85020 taaaaccttt ttccctttcc cagtttaact tttgaagcat gtttgaattt tattttcaga 85080 85140 gtaaaacata atttttaatg tttatgtact tttatttgca atactgtctt gacaacactg tetgagatat caggetetta aaaatgaaat aaagtttgea atgtgggget atgteteeca 85200 85260 tggtttaact ttctgcattt ctgtgttttc tcagatatga tttttcttca gggaacagag 85320 gtcatattta aagtggcttt aagtctgttg ggaagccata agcccttgat tctgcagcat 85380 gaaaacctag aaaccatagt tgactttata aaaagcacgc tacccaacct tggcttggta 85440 cagatggaaa agaccatcaa tcaggtatga gtcagtccaa accttgcaaa tgcttaagcc 85500 atcotagata tgtagaaact taaatctctc ttgagcagga actgtttcct accactttgt 85560 gttctgaacg gcattctgca tgatgcctgg catggaggag gcatatcaca aacgtgtgga 85620 atgatcgtga gtgtgtgtgt tgtgagcgtc atggtgaaat gccacatgga aacatggtgg 85680 caatgtttag ctgtagaaac cagcacaggt tattagtagt ttcttacatt taagagactt 85740 cagetetagt agettgttet tetgaaacat atatatattt atgeaatgat geaatgtagg 85800 gttttgtaca ttgagtgctt tgatttgtgt gtgtgtatgt tgaatggttt taattggaat 85860 tttctccaaa taattctttg ataacaaagt tatgataggg aacatatatt ctatgaattt 85920 gtttcatgat gtgtgtgtgt ggtttttttt taactgaatt cagttcaata tctgtggctt 85980 cattacctct ggtttcagta tacaatagaa ccattatcct ctgaagtgtt agaggctgag 86040 aggtgagttt tactgggaat tacaactaaa ctagatggtg aatgccctgg gttgggcatg 86100 ggagcagatt ttgatcctgt acagtttaag gaagaacctg ccagtaatgg ctgctgatga 86160 tggaggacta tgctccttag tagagaaagg ggcatcattg gaagtgtcta gaggctgggt 86220 agctacgaaa gcactggaag ggatttcatc attgagtcac tgcagagtca gcagtcaagc 86280 ctctcataag cctagaatct gctccagata gtcatgtgtc acttaataat agcaatacgt 86340 tttgagaact gcatcattag gtgatttcat cattgtgcaa acttcataaa gtgtacttaa 86400 86460 acccaagtgg tatagectac tacacaceta cacatgttgt teetaggeee tacaaatetg tatgacatgt gactatactg aatactgttg gtagttgtaa cacaaggtgg agtatttgtg 86520 tatctaaaca taactaaaca tagaaaaggt acaataaaaa tatggtattc taatcttatg 86580 ggaccaccat catatatgca gttcgccatt gaccaaaatg ttgtaatgca acacacaact 86640 gtgtaacgaa agcatagagc aatcaggcaa aaacaaatgg tgaaataaag ctatttttga 86700 aaaatcctta ctctggagat ttctgtagtc caaaaggaat ccatgattcc agtggattgc 86760 atccacatgc agtgtttgtg attttcattt gcagccacac cttaggtgtt aagcacagaa 86820 aaagatgcaa atttggcctg caaaagaaag aggtttcata ccagttgtta actttagatt 86880 tetgtttgca cattgcatat gecettatga aagaacagtt ettgtetgtt etgeactcat 86940 ctttaattga gageetetee atetetttte etteeetgga acaetettet tgatgtggat 87000 agetttgget tgttggggee ttgetttttg ceaggttggg tgggeatega aatataegea 87060 87120 tgateceatt cagtgcagge egeacteetg agaggatata aagtggttea etgactgace 87180 cacactcact ctgctgcaaa gtggaaaggt aggggttcaa actcaagtcc ctcccacctc aaagtgettt ageagetete etacaetgee aagageetet ggaggteatt taatttagag 87240 tttttcccta ttttaccagg attctaatac tgacttctcc acccttttga ttctttgatt 87300 tetggeattt teatteatte tttettteat teattetett ettacagett ttgttgeatg 87360 87420 tacttactta catttacagc ttctagggca gacccccgag agccttggtt acctagactg 87480 agggetatat ceaetacetg acatgteace ttgeteetgt ceeteaggee atceeagetg

acattgttta	cctcctaagt	attgagcctc	agaaaaaaat	cccattgtct	cctattttct	87540
gtaaaaaaca	aaaaataaaa	cgtattgaga	atacttagga	tacatcaggt	gctgtttcag	87600
tgctggaaga	gtggaaatgg	acacagcatg	ggaagaaaat	agctgtgcgt	gtacctggtt	87660
tgtttcaagc	cgctatctgg	ctatttggaa	gttgccattc	atttttccac	tgacttttt	87720
ttttttt	tttttgagac	agagtettge	tctgctaccc	aggctggaat	gcggtggtgt	87780
gatettgget	cactgcaacc	tccaccttct	gggctcaagc	attctcgtgc	ctcagcttcc	87840
tgagtacctg	ggactatagg	catgtgccac	cacgctcagc	taatttttgt	atttttagta	87900
gaggtggggt	ttcgccatgt	tggccaggct	ggtcttgaac	tectggeete	aagtgatcca	87960
cctgcctcat	cctcccaaag	tgctgggatt	acatgcatga	gccactgtgc	ccgacctcca	88020
ctgacttaat	aactctagga	cataggtatt	ataattccta	tttttataga	tgaagctgag	88080
cagagagtaa	catgcccggc	cccctgtaga	aaggcagggt	ctgtgggagc	cagggctgtg	88140
aggttggagc	tgaggtgttt	gagtccagct	ggacttaaaa	gatgacctaa	gatcggctgg	88200
cagacatttt	ccaaagaagg	ccaggcagta	tatattttag	gcttcacggg	tcataacgtc	88260
acaactactc	acctctgcga	cagctactca	cctcagcctt	tgcagcacaa	acaatccatg	88320
acagcatgta	aggggatgat	gggtcatgtt	gcaaaaaact	ttattttaaa	aactgtgcag	88380
cgtgatgggc	ttggtccaca	ggtggtggtt	tgctgatccc	tgaactaaag	gatcacagca	88440
tgtgagaagg	tacaggaatg	agagcagaga	gcaattctca	gaacctgagg	tgttcaacat	88500
tttgggggta	tcgggagcac	aaaaatttgc	atttagggcc	cgggttttta	tcagtggtcc	88560
ttatagaaag	tagatctacc	cgcatctctc	ccctctttcc	ctctggtgtt	tctatctgaa	88620
cttgacatct	gagtgttctc	tgtcaggctt	tctgctttcc	cactgccccc	tcccatccag	88680
agggtgctgt	agtcttaagt	tctacacaga	aagcagacat	ggggtccagc	atgattcctc	88740
tgcagcctta	gagatecece	aggcccaaat	cttgggggtc	ttcagagtag	atgtaatgga	88800
tgcttccatt	gcagatggtg	gagcatactg	ccctgtgtac	agatgg ggtg	gggcaggaag	88860
tgagggcatc	tgggtggtcc	tctgtgatag	ctgtggttcc	atttcattaa	atgccctccc	88920
tgtatagata	gtctcagcct	gagcaactgg	gacacagctg	gatcctgggt	ggaagaggtg	88980
gccaaggaca	gggcţctgga	gctctggctc	aatgtggaca	ggctgaaagc	agccagagag	89040
ggcaattcca	aaggtgacag_	agcccaggac	agaagaccaa	gggtgtctga	agcctttgtg	89100
gcagtgtact	taccaaagga	cctgtccatt	gaataaacca	tctacatctt	ctgacccaag	89160
aagaatggaa	actttgggaa	taattagtaa	caaaggaaag	gaggtcagtg	ttgtattctt	89220
gtcacagtgg	gtgctctgtg	gctgtgaagt	ctcagctcag	tttaaggaag	aaaaaagga	89280
agggtggctg	tgggaggcag	ggcaaatagc	ctattttcag	catcctttag	gctccactca	89340
gagcatggcc	tcagcccagc	attgtcatca	tatcatgtca	gagcttgtta	gaaactcagc	89400
atctggggcc	acacccgaga	cctgctgaat	cagaatctgc	atttcagtga	gatcaccagg	89460
ggattcacat	acacacaaac	agctgagaaa	ccctgctgtg	ggcaactctg	ttagaaacac	89520
aatgaacaaa	ggageceetg	ttccagttga	gcttgtaggt	tagaaaccag	ggttcctgta	89580
ttcagaagac	acacctcaaa	tcaggggcaa	aggtgcctct	tctgcctgtg	ggggagccgt	89640
cacttcttgg	gcagtttgca	ccgtggaaaa	ggagtagttt	tgtacg ag ga	caactggtgc	89700
cataccagga	gggtggggcg	tggcggggag	aagtggttta	ccactggcgt	tgttgaaaat	89760
				gtgggtttga		89820
				caggccgact		89880
				gcagggggtt		89940
aagctctgtg	gcgagttctg	caaaccrtca	gggttcatga	ctttattaat	cagtgtccat	90000

ggactgtgaa gagaaatgct gagtctacaa tagcaaatga gccaagaaca taaacagaca 90060 attcaccgaa gaggagatat ctagtaaaca aatatctggg aaagtatttg gcttcatgtg 90120 taatttaaac ttatgtaaca tataatgctt tactctacta gataatagaa agacatttct 90180 tgatgccagt acccagcacc aagggtatac tgtatgcaga acattagcat gttgctgatg 90240 gcagtgcaca ttgattagtg gctgttggga gacaatttgg cgaaacatat cccaagccag 90300 taaaatattc ataccetttg actcagtcat cccgtttctt ggaatgtatc ctcaggaaat 90360 aatccaaaat atgagggaag ccatatgtat aaggatattc tcctagactt gtcacttata 90420 ataacagaaa cttggaacta gatgtctaac acttgatgac tggattaata tgatgatggt 90480 aggttgaget ggtagaatat catgaageca gttatatata gegacatgaa aaagetetta 90540 tttgatacaa tgttaagtaa aggaaaagtg ggataggaaa ttttatgttgʻgttatgttta 90600 gaactagaaa aacatgcttt taggaaatag gaaatatagc tagatataaa agttgtattt 90660 ggtgattttt cttttatttt tcaagcttcc aataatgtag ctctattgct tcagtaactt 90720 aaaatagttt tatcttttcg gcaaaacatc gaaagtatgg aaatagtcat tcctactttg 90780 gcaaacagaa gagaaatttt cttcagtacc aaaattctgg aacttgactg aaaactatga 90840 agaacctaag agccaggatg acaggaaggc tctagatccc cagtaattac aactctagtg 90900 gaattgetet gagatgggee ageaagaaag aagatgagag eeagteeeee ttgeagaggg 90960 gccaggtacc ttgcagcttt gtgtagtgac cagtgctcag ggaacggctt aggcaagacc 91020 ctgggggagg tgggcactgc acttgtccag cctcaggagt gactcagacc agaaatgaaa 91080 acaccttaaa gtgtatatat cttgttttcc tatcaacacc tagtttttaa tattcgtctg 91140 ttttattcat ctgagacaac ataccaaagg attgggtttt taatgttagg ccttcctgct 91200 ctttctctgg ctgagaactg ctcctggcag tggatcactt gtgctgtcta agtgtgcaag 91260 gacaggegee ceteecaatt etttette eccaagtaat tageecaagg getgaageee 91320 togtocagtg accagggtto tootttgaco accagootoa tattgocatg gtttggggta 91380 aattcagggg cataactgca gaatgaaggg cctaggagtc ttggcagtca ggagatcatc 91440 aggcaattaa gcagagatga ttgtgaccca gggtggttcc tagggattaa tggaggcctg 91500 gaagagttta tggetttggg tactgetgag agccattaac ttaacacaga acatcaatce 91560 gtaggaaaag ccagaggttt tgttcccagg ctttccaggt taggagatca cttaaatctt 91620 tgtgaaagaa aaaaagtaat agtgtacatg acatttattc agcaccatat ttataattat 91680 acatgagtgc caaacaatct cagttttaac atttgtggtt tttactgttc agactattca 91740 gaagttccat gacgtggcat ccattgatga tggttttgct gaggttgaaa tgtgagggtt 91800 ttgtggcagg tgtgtggaag cagatccctt agctaagaga gcgcctgctc aacctgccag 91860 gcgtgtgctg ggaacttcca ggagttgcct cgtttaattc tcacagccat cctgggaggt 91920 aggegetgte tgeeceaggt gggaggeeag gaaetgtgge tgagagaggt taagtaeega 91980 gctcaagtca ttcatagagt cgccagtgga gccgcaattt taaggctgac tcaaagcctc 92040 tgagccagtt accteggetg teaaatggag ataataatee etgetgaeet caeggteget 92100 ctgaaggaca aaagagaatc agttcaatcc agtaaacaat tctctctct cctcttactc 92160 ccctcgcaca catgcacaca caccccacag atataatgga ttttagtttt taggcatcaa 92220 atgacttttg tgccttatta atattatcca ctgaatcaaa aacagcaagc tgaaaaattc 92280 atctcaagga agagaaaata agattgttgg gaatggtgag aaaggaaaca tggtttttga 92340 aaattgattc cagggaagat aggctagttt gaatgccagt agggagccat cagaagaagt 92400 92460 tttctgccct tttaaagcct gacttcactt tctgaatgtg tgttctgatc tagcagggtt 92520

WO 00/08209

			56			
tttttttt	ttcttttaag	atggtcccag	cttgactgca	ttctcagatc	catcagataa	92580
acgttagggc	ttcactgctg	tgctgagagg	ccccagcccc	tggggttctc	tcatagaaac	92640
aactggaaag	aaaggaaatg	ccttgggcag	cagcagcagc	agctgtcttc	tgattctgct	92700
ttccgccctg	ccttccttac	caagagaaag	tacagacacg	gacggcttga	gtcacttagg	92760
cacttaggag	ttgtttttca	cacgtgtggt	gttttcgtca	ccattactat	tgtgggaaag	92820
aagacaactc	aggcatcgtt	tcgtattcac	tcatctgtgt	gggtgacatg	tgggttttgg	92880
ctcatttctg	catatttgtg	tgcaaaggag	agttttttag	taaacagtcc	cattacttag	92940
ctgttcttgt	aactctgaaa	acccaactga	actataatta	aactttgact	tggtgactct	93000
gcaaacaggc	tatgattctt	ttgtttcttt	tctcctttta	acccatagtt	gatgtatcta	93060
acctaacaga	attttcagag	aaaagaagtg	aaataagaac	taaaaataaa	tttttatgtc	93120
tttaaaaatg	agaggttttt	tttttttt	tggcttttgg	aaggtgagta	tcaaaaacct	93180
gtacttaatg	ttaccttgga	attatttcta	gatgtttctt	atatcctttt	gtcccaagta	93240
aaattattac	cttctcagtg	cgtagttttt	cttatttatt	acttctagta	ccaagtgtag	93300
agctaagcgt	agaggagacg	cttcacaggt	gcgcattgtc	gtgattgcag	acgcctgcct	93360
gtacttgtgg	ggtttttctc	agttttagta	cgtgatgact	tttctttcta	taacaggtat	93420
ttgaaatgga	catcgctaaa	cagttacaag	cttatgaagt	tgagtaccac	gtccttcaag	93480
aagaacttat	cgattcctct	cctctcagtg	acaaccaaag	aatggataaa	ttagagaaaa	93540
ccaacagcag	cttacgcaaa	cagaaccttg	acctccttga	acagttgcag	gtagagcata	93600
tttataaagc	agcttcctga	atcacaaata	tatggtagtt	cattaactca	ccaaaggcaa	93660
cagcaggctg						93720
ggatcaaagg						93780
				ttctgcactt		93840
ggggacaggc						93900
gccgttgtca						93960
				aataaatata		94020
				agatttgatt	ar.	94080
				tactcaggtg		94140
				cagagcaagc		94200
				acccccacat		94260
				ttgtcaatct		94320
_	_			cacaagtgtg		94380
_				tgtttttgag		94440
				gggctgtcct		94500
				atgagctttg		94560
				ctatctgaaa		94620
				gaaagagagg		94680
				tgaaatttaa		94740
				atacaattta		94800
				cctgtctggc		94860
				tgataggtcc		94920
				agtgtttagc		94980
aggcagttct	caaataaaag	geregerea	ccgggtaaat	gaccttgtag	cccccggcg	95040

WO 00/08209

•			•			•
acagagcata	gaaagtaatt	tcatgctgct	cctgtgctat	tgtttttgtg	aagacaggga	95100
gctgtgaaaa	actgcttagc	tacctacatt	cctcaataaa	ggcatcagac	agtaattggt	95160
gattacagat	gttctccctg	gaatggtcgt	tctcttgacc	aagtagtcct	acacttctgg	95220
aaggatcatt	cagaactgtg	gtctatgcca	acccaccagt	agttcctgag	tccctgcagt	95280
gctgagtgct	gggggccacc	aagttgagta	agacactgca	gctctcaaag	agttggatct	95340
aggattgtat	tgtatcgatt	tgtggtgttt	ggatatagtt	tttccatgat	cccctacgaa	95400
aatatgcctc	tcatatgtaa	gaatcatgcc	tcctccgtgt	acacttttca	gacactgaca	95460
aggaagggtg	ttcaatacag	tactgaattt	tcatatagct	tttctggggg	ggccaaaata	95520
ccaaaatcaa	cccatttcct	acctttattc	tgtccataaa	attgttagaa	atatcaaaat	95580
cccatttcat	tccctgttaa	atacatgtga	acgttgtcta	gacgctggag	agcaaattct	95640
accacctctt	ttgttcagca	gtacatcaga	cgattgcata	gacgtgccag	atggaaccaa	95700
ataataatgc	acatggattt	gtcataatcc	gtacaagtca	ttgacgccca	cactgagcca	95760
ggtgctgtgg	gagacaggac	catgtgtgaa	agagaagaca	tgcttgcttc	tataaaagca	95820
tcggtgttat	tgaggagacc	tgacattaat	gcagaatagc	aaatgaccat	gcaaattaat	95880
tcactactaa	ctaagctgca	ggttgcacct	cggaatgcag	aggggcttca	aagtgatgag	95940
ggtaggccct	gagccaggcc	ctgatgatgg	gtggattttg	aggatcagag	agtac agct t	96000
agagagatac	cccaagtggg	accacccctt	gcccagtagg	ctgacaaact	aaggctcttg	96060
gtcccttttt	catattttgg	gtgttctagt	ggcccagcca	gagctagact	tcgagtcatg	96120
aattttctgg	cacaagtgtt	gtcacattca	aaaaagtatt	ttctttgttt	gaaaaatgaa	96180
aaatatatat	atatgtgtgt	gtgtatatat	atatgtgtgt	gtgtgtgtat	atatgtgtgt	96240
gtgtatatat	acgt gt gtgt	gtatatgtgt	gtgtgtgtgt	gtgtgtgt	gtgtatatat	96300
gtgtatatat	ataccatttt	teccacetaa	aatggagcat	ggcaaatctg	gactggatta	96360
gtgagataga	ccaagtcaca	gagcactcca	ggatgcagct	gtgagctggg	gaacaggtca	96420
gaaaggcctc	agggacatca	gcatacatgt	tggagtttct	gcagttttct	tagggaaccc	96480
tttaatgtca	ctagagctaa	cacacttgtc	acctgggaag	caagcetgee	agagcaaatt	96540
agagagacga	gggacagttg	ctagaaagac	acacctggaa	gttctattta	actagcatta	96600
attatgtgct	aggtgctga'g	gatgtagact	gagtgagatc	ctcattcctc	ctctgtaggg	96660
tggaagagag	gatattgctt	gtctccatgg	ctcgtagtga	acagtcagtg	agaccaggca	96720
cataaacctc	ttagcagaac	acttggcctt	tctaaggact	ccatatgtgt	tccggggtaa	96780
atgcctgtgt	ttcttgacgt	agtgatgtct	tgttcctcta	gacatcacta	actttacaca	96840
gtagctttag	atggcgtgga	cgtgaataaa	tgcaacttag	gttttcttgt	tggtttcttt	96900
ttgag tatca	ttgtgtttgt	aaagaatttc	agattagagg	attgttacca	cgtgggcctt	96960
caggaggaaa	ctgttttgag	tttttgtcag	cccgaaatcg	atttgtgcgt	ttaagtatat	97020
gtgctcatca	aaacaggcca	ggctctgctg	cagtaacaaa	cttacaagtc	tccgaggctt	97080
ggcacaacgg	aagtctttgt	gtcactgacg	cccacttcag	ctttgtgttg	ctgaagcatt	97140
ctttggccct	atccgagctg	tccctctggt	ggtggtgcct	gggggtttgg	gttccctctg	97200
tgttgtgatc	caccatctca	gcatcggctt	ccacagcagc	catagcagga	gaagaaaatg	97260
ctgggggctc	tcagggtgct	tttaagggcc	tggccaccga	cctgcaaggg	gtgcgagttg	97320
ccttcctgtg	tggccagaac	tgatgataaa	ctgtagactc	atccctgctg	aaact cg gct	97380
				tcccacctgt		97440
				aaatggtagg		97500
ttgaggccac	cattgagaag	ctcctgagca	gtgagagcaa	gctgaagcag	gccatgctta	97560

THE RESIDENCE OF THE PROPERTY OF THE PROPERTY

-			58			•
ccttagaact	ggagcggtcg	gccctgctgc	agacggtgga	ggagctgcgg	cggcggagcg	97620
cagagcccag	cgaccgggag	cctgagtgca	cgcagcccga	geccaeggge	gactgacagc	97680
tctgcaggag	agattgcaac	accateccae	actgtccagg	ccttaactga	gagggacaga	97740
agacgctgga	aggagagaag	gaagcgggaa	gtgtgcttct	cagggaggaa	accggcttgc	97800
cagcaagtag	attettacga	actccaactt	gcaattcagg	gggcatgtcc	cagtgttttt	97860
tttgttgttt	ttagatacta	aatcgtccct	tctccagtcc	tgattactgt	acacagtagc	97920
tttagatggc	gtggacgtga	ataaatgcaa	cttatgtttt	cttgttggtt	cctttttgag	97980
tgtcactgtg	tttgtaaaga	gcattcacaa	tacggtggaa	tttcaaaagc	tggaagagct	98040
cgagatcatg	cctcaggcaa	aggcgtgggt	ccatcgttct	tccgagaggg	tttgtgtggc	98100
gactacaccc	tcagcgtccc	tggcaaggtg	cagttggctc	tcgcccattc	'ttgttatgga	98160
aacctaagat	gatcattggg	aagatcagtg	atcttgggtc	attgatccct	ggctcagagg	98220
atagcggttt	ccatcataaa	ccaagatgat	gagttcagcc	tttatccctc	gtggttccac	98280
tagatgtaac	ttaaaggagt	taacatttga	ggactttgtt	ctacatcaga	ttttactatt	98340
tgaatgttta	agatcacttt	attgaatttg	aagatcatca	aattaaataa	aatgatttat	98400
ttaatttgga	tatcctgatc	actgtcaagt	gaaatggatc	tctctctttg	gtatttaagg	98460
aagtttgtct	ttaaaaaaaa	aatagagtgt	tttcatacat	ttttgcttat	cccataagta	98520
cagttgatca	aagtcatagt	aggtaaatgc	tttatgggac	agctgacacc	ttttagaccc	98580
taccaggtat	tgctagcatg	tgagctgcag	ttgtggggtc	tgagatattt	ctttgtggta	98640
gtttcatacc	catactatag	agtcatgtat	ttatttttgc	ctgttgtgtg	atgtaatgca	98700
atcatgttcc	tttgagtctc	catcccttgg	aaatctgact	tcttgcagaa	ggagtaggca	98760
catcaagata	ttcaggggtg	ccccaagagt	ctgggacttt	caaaaaaaaa	agatcaggct	98820
nnaactgcag	tcagatttat	gacagctgac	agtttt tca g	aggtcgcaca	cagtgactct	98880
cctctctcag	gatgacgagg	acctgtgcct	tcaacaagca	aaatgctgct	cacggttgtc	98940
ctgcttgcag	ccagtcactg	tgtaaagcct	ctctgatgtg	cacttaagag	tgggttgctt	99000
tctcacaaag	atggggttct	gtgcagtcac	aggtcacttc	cttgacaaca	caatcatttc	99060
tgatctttat	cactgtaacc	acgtcttcta	ttccatagga	gtttcttttg	attctctcag	99120
ttgcgggggg	catctcttaa	tcctggggta	aaaggagaga	ttgccatact	tagactcact	99180
gtgagtctcc	ccggccattt	cacgaggaga	ccacagtgct	gccaccagtg	cctaaacagg	99240
tggctggcat	tcgagacttc	ctcctgttcc	ctgggtcaga	ggatagcggt	ttccatcata	99300
aaccaagatg	atgagttcag	cctttatccc	tcgtggttcc	gctagatgta	acttatagga	99360
gttaacattt	gaggactttg	ttctgcatca	gatcttacta	tttgaatgtt	tactgttgga	99420
ttttgggcat	cttattactg	ttactcaaaa	acattgactc	tgcatcaaga	aagaaacaag	99480
aaagcaataa	aacaagaaat	aattcatgct	cacattttta	tggtggtttt	tttttttt	99540
ttaactttgg	atttttgctt	ttcagcccag	gagtaaagga	atgccttatg	aacacctgtg	99600
gcctacgtgt	ggtcatgacc	caaccatcag	tgagattatt	tgagatattg	gtgtctgcat	99660
ccagtgttgt	tatctgagtg	tttattacgt	aagttgtaac	acctctacac	agggtgtgag	99720
	atgagaccag					99780
	gaaaggactg					99840
	taggcaagga					99900
ttaagccatg	ttttgggtta	tatttccccc	aacactcatt	tgtgcacttg	gtggtgtcaa	99960

WO 00/08209	59	PCT/IB99/01444
- 222	37	·
<211> 3983		
<212> DNA		
<213> Homo sapiens		
<220>		
<221> CDS		
<222> 1713725		
		_
<220>		
<221> polyA_signal		·
<222> 39423947		
<223> AATAAA		
<220>		
<221> misc_feature		
<222> 36		
<223> n=a, g, c or t		
<400> 3		
ccaggccgtc cccaggatgc cc		
agegegeege ggeacaggtt te	tgcatatg aagtgtgtaa a	aatagattgc ttgatccaaa 120
acagaaaaac agtgataact gt	tttgctga gttcccagac c	
		Met Glu
		1
cca ata aca ttc aca gca Pro Ile Thr Phe Thr Ala.		
5	10	15
gtg gat ttt ggc ctg cag		
Val Asp Phe Gly Leu Gln		
-	_	30
acc acc atg ccc atg ctg		
Thr Thr Met Pro Met Leu		
35 40	45	50
agc agg cag tcc acc aga	aag gaa cct gta acc a	aag caa gtc cgg ctt 368
Ser Arg Gln Ser Thr Arg	Lys Glu Pro Val Thr I	Lys Gln Val Arg Leu
55	60	65
tgc gtt tca ccc tct gga	ctg aga tgt gaa cct g	gag cca ggg aga agt 416
Cys Val Ser Pro Ser Gly	Leu Arg Cys Glu Pro G	Glu Pro Gly Arg Ser
70	7 5	80
caa cag tgg gat ccc ctg	atc tat tcc agc atc t	ttt gag tgc aag cct 464
Gln Gln Trp Asp Pro Leu	Ile Tyr Ser Ser Ile F	Phe Glu Cys Lys Pro
85	90	95

wo	00/08:	209]	PCT/IB9	99/01444
	-							•	50							•
cag	cgt	gtt	cac	aaa	ctg	att	cac	aac	agt	cat	gac	cca	agt	tac	ttt	512
Gln	Arg	Val	His	Lys	Leu	Ile	His	Asn	Ser	His	Asp	Pro	Ser	Tyr	Phe	
	100					105					110					
gct	tgt	ctg	att	aag	gaa	gac	gct	gtc	cac.	cgg	cag	agt	atc	tgc	tat	560
Ala	Cys	Leu	Ile	Lys	Glu	Asp	Ala	Val	His	Arg	Gln	Ser	Ile	Cys	Tyr	
115					120					125					130	
gtg	ttc	aaa	gcc	gat	gat	caa	aca	aaa	gtg	cct	gag	atc	atc	agc	tcc	608
Val	Phe	Lys	Ala	Asp	Asp	Gln	Thr	Lys	Val	Pro	Glu	Ile	Ile	Seŗ	Ser	
				135					140					145		
atc	cgt	cag	gcg	9 99	aag	atc	gcc	cgg	cag	gag	gag	ctg	cac	tgc	ccg	656
Ile	Arg	Gln	Ala	Gly	Lys	Ile	Ala	Arg	Gln	Glu	Glu	Leu	His	Cys	Pro	
			150					155					160			
tcc	gag	ttc	gac	gac	acg	ttt	tcc	aag	aag	ttc	gag	gtg	ctc	ttc	tgc	704
Ser	Glu	Phe	Asp	Asp	Thr	Phe	Ser	Lys	Lys	Phe	Glu	Val	Leu	Phe	Cys	
		165					170					175				
ggc	cgc	gtg	acg	gtg	gcg	cac	aag	aag	gct	ccg	ccg	gcc	ctg	atc	gac	752
Gly	Arg	Val	Thr	Val	Ala	His	Lys	Lys	Ala	Pro	Pro	Ala	Leu	Ile	Asp	
	180					185					190					
gag	tgc	atc	gag	aag	ttc	aat	cac	gtc	agc	ggc	agc	cgg	9 99	tcc	gag	800
Glu	Cys	Ile	Glu	Lys	Phe	Asn	His	Val	ser	Gly	Ser	Arg	Gly	Ser	Glu	
195					200					205					210	
agc	ccc	c gc	CCC	aac	ccg	ccc	cat	gcc	gcg	ccc	aca	ggg	agc	cag	gag	848
Ser	Pro	Arg	Pro	Asn	Pro	Pro	His	Ala	Ala	Pro	Thr	Gly	Ser	Gln	Glu	
				215					220					225		
cct	gtg	cgc	agg	ccc	atg	cgc	aag	tcc	ttc	tcc	cag	ccc	ggc	ctg	cgc	896
Pro	Val	Arg	Arg	Pro	Met	Arg	Lys	Ser	Phe	Ser	Gln	Pro	Gly	Leu	Arg	
			230	,	٠.			235					24 0			
tcg	ctg	gcc	ttt	agg	aag	gag	ctg	cag	gat	ggg	ggc	ctc	cga	agc	agc	944
Ser	Leu	Ala	Phe	Arg	Lys	Glu	Leu	Gln	Asp	Gly	Gly	Leu	Arg	Ser	Ser	
		245					250					255				
ggc	ttc	ttc	agc	tcc	ttc	gag	gag	agc	gac	att	gag	aac	cac	ctc	att	992
Gly	Phe	Phe	Ser	Ser	Phe	Glu	Glu	Ser	Asp	Ile	Glu	Asn	His	Leu	Ile	
	260					265					270					
agc	gga	cac	aat	att	gtg	cag	ccc	aca	gat	atc	gag	gaa	aat	cga	act	1040
Ser	Gly	His	Asn	Ile	Val	Gln	Pro	Thr	qaA	Ile	Glu	Glu	Asn	Arg	Thr	
275					280					285					290	
atg	ctc	ttc	acg	att	ggc	cag	tct	gaa	gtt	tac	ctc	atc	agt	cct	gac	1088
Met	Leu	Phe	Thr	Ile	Gly	Gln	Ser	Glu	Val	Tyr	Leu	Ile	Ser	Pro	Asp	
				295					300					305		
acc	aaa	aaa	ata	gca	ttg	gag	aaa	aat	ttt	aag	gag	ata	tcc	ttt	tgc	1136
Thr	Lys	Lys	Ile	Ala	Leu	Glu	Lys	Asn	Phe	Lys	Glu	Ile	Ser	Phe	Cys	
			310					315					320			

WO 00/08209

	61	1 C 1/1D 7 7/01444
tet cag gge ate aga ca	gtg gac cac ttt ggg tt	atc tgt cgg gag 1184
Ser Gln Gly Ile Arg His	Val Asp His Phe Gly Phe	e Ile Cys Arg Glu
325	330	335
tot too gga ggt ggc gg	ttt cat ttt gtc tgt tac	gtg ttt cag tgc 1232
Ser Ser Gly Gly Gly Gl	Phe His Phe Val Cys Ty	. Val Phe Gln Cys
340	345 350)
aca aat gag gct ctg gt	gat gaa att atg atg acc	c ctg aaa cag gcc 1280
Thr Asn Glu Ala Leu Va	Asp Glu Ile Met Met Th	: Leu Lys Glm Ala
355 366	365	370
ttc acg gtg gcc gca gtg	g cag cag aca gct aag gcg	g cca gcc cag ctg 1328
Phe Thr Val Ala Ala Va	Gln Gln Thr Ala Lys Ala	a Pro Ala Gln Leu
375	380	385
tgt gag ggc tgc ccc ctg	g caa age etg cae aag ete	tgt gag agg ata 1376
Cys Glu Gly Cys Pro Let	ı Gln Ser Leu His Lys Let	ı Cys Glu Arg Ile
390	395	400
gag gga atg aat tot to	c aaa aca aaa cta gaa cto	g caa aag cac ctg 1424
Glu Gly Met Asn Ser Ser	Lys Thr Lys Leu Glu Leu	ı Gln Lys His Leu
405	410	415
acg aca tta acc aat cag	g gag cag gcg act att tti	gaa gag gtt cag 1472
Thr Thr Leu Thr Asn Gli	Glu Gln Ala Thr Ile Phe	e Glu Glu Val Gln
420	425 436)
aaa ttg aga ccg aga aa	gag cag cga gag aat gaa	a ttg att att tct 1520
Lys Leu Arg Pro Arg Ass	n Glu Gln Arg Glu Asn Gla	Leu Ile Ile Ser
435 449		450
ttt ctg aga tgt tta ta	gaa gag aaa cag aaa gaa	a cac atc cat att 1568
	c Glu Glu Lys Gln Lys Glu	
455/ -	460	465
	a tog cag atg gca gca gag	
	r Ser Gln Met Ala Ala Gli	
470	475	480
	e act cga ttt agg cta ga	
	a Thr Arg Phe Arg Leu As	-
485	490	495
	a aca gag tct tta gaa ag	
	u Thr Glu Ser Leu Glu Se	-
500	505 51	
	c etg cag gaa cac tec at	
	y Leu Gln Glu His Ser Il	ser val Asp Leu 530
515 52		
	t aca tta agt aac acc ag r Thr Leu Ser Asn Thr Se	
		545
535	540	243

and the second of the second o

									62								
gtg	g tgt	gaa	aag	gag	gcc	ttg	ccc	atc	tct	gag	ago	tcc	ttt	aag	ctc	1856 -	
Va]	Cys	Glu	Lys	Glu	Ala	Leu	Pro	Ile	Ser	Glu	Ser	Ser	Phe	Lys	Leu		
			550	ı				555					560				
ctc	ggc	tcc	tcg	gag	gac	ctg	tcc	agt	gac	tcg	gag	agt	cat	cto	cca	1904	
Leu	Gly	Ser	Ser	Glu	Asp	Leu	Ser	Ser	Asp	Ser	Glu	Ser	His	Leu	Pro		
		565					570					575					
gaa	gag	cca	gct	ccg	ctg	tcg	CCC	cag	cag	gcc	ttc	agg	agg	cga	gca	1952	
Glu	Glu	Pro	Ala	Pro	Leu	Ser	Pro	Gln	Gln	Ala	Phe	Arg	Arg	Arg	Ala		
	580					585					590						
		ctg														2000	
Asn	Thr	Leu	Ser	His	Phe	Pro	Ile	Glu	Cys	Gln	Glu	Pro	Pro	Gln	Pro		
595					600					605					610		
		9 99														2048	
Ala	Arg	Gly	Ser	Pro	Gly	Val	Ser	Gln	Arg	Lys	Leu	Met	Arg	Tyr	His		
				615					620					625			
		agc														2096	
Ser	Val	Ser		Glu	Thr	Pro	His		Arg	Lys	Asp	Phe	Glu	Ser	Lys		
			630					635					640				
		cat														2144	
Ата	Asn	His	Leu	GIY	Asp	Ser		Gly	Thr	Pro	Val		Thr	Arg	Arg		
ant	+ ==	645					650					655					
		tgg													_	2192	
mis	660	Trp	Arg	GIII	GIII	665	Pne	Leu	Arg	vai		Thr	Pro	Gln	Lys		
aca		gat	tat	tcc	200		t = t	~ 22	~a+	+ - +	670	~~~					
		Asp											_			2240	
675	-1 -		,		æ80	••• 9	-7-	CIU	пор	685	. ;-		nen	GIY	690		
ctt	ccc	cca	саа	,	-	tta	αаа	cca	att				aaa	ccc		2200	
		Pro														2288	
			_	695					700	-7-			Cry	705	PIIC		
ggc	CCC	cca	cca	gag	gaa	aag	aaa	agg	aca	tct	cqt	qaq	ctc		gag	2336	
		Pro														2330	
			710					715			_		720				
ctg	tgg	caa	aag	gct	att	ctt	caa	cag	ata	ctg	ctg	ctt	aga	atg	gag	2384	
		Gln												_			
		725					730					735					
aag	gaa	aat	cag	aag	ctc	caa	gcc	tct	gaa	aat	gat	ttg	ctg	aac	aag	2432	
Lys	Glu	Asn	Gln	Lys	Leu	Gln	Ala	Ser	Glu	Asn	Asp	Leu	Leu	Asn	Lys		
	740					745					750						
cgc	ctg	aag	ctc	gat	tat	gaa	gaa	att	act	ccc	tgt	ctt	aaa	gaa	gta	2480	
Arg	Leu	Lys	Leu	Asp	Tyr	Glu	Glu	Ile	Thr	Pro	Cys	Leu	Lys	Glu	Val		
755					760					765					770		

WO 00	0/082	209												1	PCT/IB99/0	1444
	-								53							•
act a	aca	gtg	tgg	gaa	aag	atg	ctt	agc	act	cca	gga	aga	tca	aaa	att	2528
Thr I	Thr	Val	Trp	Glu	Lys	Met	Leu	Ser	Thr	Pro	Gly	Arg	Ser	Lys	Ile	
				775					780					785		
aag t	ttt	gac	atg	gaa	aaa	atg	cac	tcg	gct	gtt	aaa	caa	ggt	gtg	cca	2576
Lys F	Phe	Asp	Met	Glu	Lys	Met	His	Ser	Ala	Val	Gly	Gln	Gly	Val	Pro	
			790					795					800			
cgt c	cat	cac	cga	ggt	gaa	atc	tgg	aaa	ttt	cta	gct	gag	caa	ttc	cac	2624
Arg H	His	His	Arg	Gly	Glu	Ile	Trp	Lys	Phe	Leu	Ala	Glu	Gln	Phe	His	
		805					810					815				
ctt a	aaa	cac	cag	ttt	ccc	agc	aaa	cag	cag	cca	aag	gat	gtg	cca	tac	2672
Leu I	Lys	His	Gln	Phe	Pro	Ser	Lys	Gln	Gln	Pro	Lys	Asp	Val	Pro	Tyr	
8	320					825					830					
aaa g	gaa	ctc	tta	aag	cag	ctg	act	tcc	cag	cag	cat	gcg	att	ctt	att	2720
Lys G	31u	Leu	Leu	Lys	Gln	Leu	Thr	Ser	Gln	Gln	His	Ala	Ile	Leu	Ile	
835					840					845					850	
gac c	ctt	9 99	cga	acc	ttt	cct	aca	cac	cca	tac	ttc	tct	gcc	cag	ctt	2768
Asp L	Leu	Gly	Arg	Thr	Phe	Pro	Thr	His	Pro	Tyr	Phe	Ser	Ala	Gln	Leu	
				855					860					865		
gga g	gca	gga	cag	cta	tcg	ctt	tac	aac	att	ttg	aag	gcc	tac	tca	ctt	2816
Gly A	Ala	Gly	Gln	Leu	ser	Leu	Tyr	Asn	Ile	Leu	Lys	Ala	Tyr	Ser	Leu	
			870					875					880			
cta g	gac	cag	gaa	gtg	gga	tat	tgc	caa	ggt	ctc	agc	ttt	gta	gca	ggc	2864
Leu A	Asp	Gln	Glu	Val	Gly	Tyr	Cys	Gln	Gly	Leu	Ser	Phe	Val	Ala	Gly	
		885					890					895				
att t	ttg	ctt	ctt	cat	atg	agt	gag	gaa	gag	gcg	ttt	aaa	atg	ctc	aag	2912
Ile I	Leu	Leu	Leu	Hís	Met	Ser	Glu	Glu	Glu	Ala	Phe	Lys	Met	Leu	Lys	
9	900			£	-	905					91,0-					
ttt d	ctg	atg	ttt	gac	atg	9 99	ctg	cgg	aaa	cag	tat	cgg	cca	gac	atg	2960
Phe I	Leu	Met	Phe	Asp	Met	Gly	Leu	Arg	Lys	Gln	Tyr	Arg	Pro	Asp	Met	
915					920					925					930	
att a	att	tta	cag	atc	cag	atg	tac	cag	ctc	tcg	agg	ttg	ctt	cat	gat	3008
Ile 1	Ile	Leu	Gln	Ile	Gln	Met	Tyr	Gln	Leu	Ser	Arg	Leu	Leu	His	Asp	
				935					940					945		
tac o	cac	aga	gac	ctc	tac	aat	cac	ctg	gag	gag	cac	gag	atc	ggc	CCC	3056
Tyr H	His	Arg	Asp	Leu	Tyr	Asn	His	Leu	Glu	Glu	His	Glu	Ile	Gly	Pro	
			950					955					960			
agc o	ctc	tac	gct	gcc	CCC	tgg	ttc	ctc	acc	atg	ttt	gcc	tca	cag	ttc	3104
Ser I	Leu	Tyr	Ala	Ala	Pro	Trp	Phe	Leu	Thr	Met	Phe	Ala	Ser	Gln	Phe	
		965					970					975				
ccg	ctg	gga	ttc	gta	gcc	aga	gtc	ttt	gat	atg	att	ttt	ctt	cag	gga	3152
Pro 1	Leu	Gly	Phe	Val	Ala	Arg	Val	Phe	Asp	Met	Ile	Phe	Leu	Gln	Gly	
9	980					985					990					

wo	00/08	209													PCT/IB99/	01444
	•							(54							•
aca	gag	gtc	ata	ttt	aaa	gtg	gct	tta	agt	ctg	ttg	gga	agc	cat	aag	3200
Thr	Glu	Val	Ile	Phe	Lys	Val	Ala	Leu	Ser	Leu	Leu	Gly	Ser	His	Lys	
995					1000	כ				1009	5				1010	
ccc	ttg	att	ctg	cag	cat	gaa	aac	cta	gaa	acc	ata	gtt	gac	ttt	ata	3248
Pro	Leu	Ile	Leu	Gln	His	Glu	Asn	Leu	Glu	Thr	Ile	Val	Asp	Phe	Ile	
				1019	5				1020	0				102	5	
aaa	agc	acg	cta	ccc	aac	ctt	ggc	ttg	gta	cag	atg	gaa	aag	acc	atc	3296
Lys	Ser	Thr	Leu	Pro	Asn	Leu	Gly	Leu	Val	Gln	Met	Glu	Lys	Thr	Ile	
			1036)				1035	5				104	0		
aat	cag	gta	ttt	gaa	atg	gac	atc	gct	aaa	cag	tta	caa	gct.	tat	gaa	3344
Asn	Gln	Val	Phe	Glu	Met	Asp	Ile	Ala	Lys	Gln	Leu	Gln	Ala	Tyr	Glu	
		1045	5				1050)				105	5			
gtt	gag	tac	cac	gtc	ctt	caa	gaa	gaa	ctt	atc	gat	tcc	tct	cct	ctc	3392
Val	Glu	Tyr	His	Val	Leu	Gln	Glu	Glu	Leu	Ile	Asp	Ser	Ser	Pro	Leu	
	1060)				1069	5				1070)				
agt	gac	aac	caa	aga	atg	gat	aaa	tta	gag	a aa	acc	aac	agc	agc	tta	3440
Ser	Asp	Asn	Gln	Arg	Met	Asp	Lys	Leu	Glu	Lys	Thr	Asn	Ser	Ser	Leu	
107	5				1080)				1089	5				1090	
cgc	aaa	cag	aac	ctt	gac	ctc	ctt	gaa	cag	ttg	cag	gtg	gca	aat	ggt	3488
Arg	Lys	Gln	Asn	Leu	Asp	Leu	Leu	Glu	Gln	Leu	Gln	Val	Ala	Asn	Gly	
				109	5				1100	0				110	5	
agg	atc	caa	agc	ctt	gag	gcc	acc	att	gag	aag	ctc	ctg	agc	agt	gag	3536
Arg	Ile	Gln	Ser	Leu	Glu	Ala	Thr	Ile	Glu	Lys	Leu	Leu	Ser	Ser	Glu	
			1110	0				1119	5				112	0		
agc	aag	ctg	aag	cag	gcc	atg	ctt	acc	tta	gaa	ctg	gag	cgg	tcg	gcc	3584
Ser	Lys	Leu	Lys	(Gln	Ala	Met	Leu	Thr	Leu	Glu	Leu	Glu	Arg	Ser	Ala	
		1129	5		£ =		1130)			~ ;-	113	5			
ctg	ctg	cag	acg	gtg	gag	gag	ctg	cgg	cgg	cgg	agc	gca	gag	ccc	agc	3632
Leu	Leu	Gln	Thr	Val	Glu	Glu	Leu	Arg	Arg	Arg	Ser	Ala	Glu	Pro	Ser	
	114	0				1149	5				115	0				
gac	cgg	gag	cct	gag	tgc	acg	cag	ccc	gag	ccc	acg	ggc	gac	tga		3677
Asp	Arg	Glu	Pro	Glu	Cys	Thr	Gln	Pro	Glu	Pro	Thr	Gly	Asp	*		
115	5				116)				116	5					
cag	ctct	gca 🤄	ggag	agat	tg ca	aaca	ccat	c cc	acac	tgtc	cag	gccti	taa	ctga	gaggga	3737
cag	aaga	cgc 1	tgga	agga	ga ga	aagga	aagc	g g g:	aagt	gtgc	ttc	tcag	gga (ggaa	accggc	3797
ttg	ccag	caa	gtag	attc	tt a	cgaa	ctcca	a act	ttgc	aatt	cag	3 999	cat (gtcc	cagtgt	3857
ttt	tttt	gtt (gttt	ttag	at a	ctaa	atcg	t cc	cttc	tcca	gtc	ctga	tta ·	ctgt	acacag	3917
tag	cttt	aga 1	tggc	gtgg	ac g	tgaai	taaa	t gc	aact	tatg	ttt	taaa	aaa	aaaa	aaaaaa	39 7 7
aaa	aaa															3983

<210> 4 <211> 3988

WO 00/08209 PCT/IB99/01444 65 <212> DNA <213> Homo sapiens <220> <221> CDS <222> 176..3730 <220> <221> polyA_signal <222> 3947..3952 <223> AATAAA <220> <221> misc_feature <222> 1..458 <223> homology with Genset 5' EST in ref : A35235 <400> 4 ataataggca ctgaagacat gttaatggaa ggtggatttg tgattcagaa cctctagact 60 acctgggcga gtcttttaaa atgtttctgc atatgaagtg tgtaaaatag attgcttgat 120 ccaaaacaga aaaacagtga taactgtttt gctgagttcc cagacccttc ccaag atg 178 Met 1 gaa cca ata aca ttc aca gca agg aaa cat ctg ctt cct aac gag gtc 226 Glu Pro Ile Thr Phe Thr Ala Arg Lys His Leu Leu Pro Asn Glu Val 10 15 tog gtg gat ttt ggć ctg cag ctg gtg ggc tcc ctg cct gtg cat tcc 274

Ser Val Asp Phe Gly Leu Gln Leu Val Gly Ser Leu Pro Val His Ser 20 25 30 ctg acc acc atg ccc atg ctg ccc tgg gtt gtg gct gag gtg cga aga 322 Leu Thr Thr Met Pro Met Leu Pro Trp Val Val Ala Glu Val Arg Arg 35 40 ctc age agg cag tcc acc aga aag gaa cct gta acc aag caa gtc cgg 370 Leu Ser Arg Gln Ser Thr Arg Lys Glu Pro Val Thr Lys Gln Val Arg 55 ctt tgc gtt tca ccc tct gga ctg aga tgt gaa cct gag cca ggg aga 418 Leu Cys Val Ser Pro Ser Gly Leu Arg Cys Glu Pro Glu Pro Gly Arg 70 75 agt caa cag tgg gat ccc ctg atc tat tcc agc atc ttt gag tgc aag 466 Ser Gln Gln Trp Asp Pro Leu Ile Tyr Ser Ser Ile Phe Glu Cys Lys 85 90 95

cct cag cgt gtt cac aaa ctg att cac aac agt cat gac cca agt tac

514

-	•	-
h	r	۱

								•	,,,							•	
Pro	Gln	Arg	Val	His	Lys	Leu	Ile	His	Asn	Ser	His	Asp	Pro	Ser	Tyr		
		100					105					110					
ttt	gct	tgt	ctg	att	aag	gaa	gac	gct	gtc	cac	cgg	cag	agt	atc	tgc		562
Phe	Ala	Cys	Leu	Ile	Lys	Glu	Asp	Ala	Val	His	Arg	Gln	Ser	Ile	Cys		
	115					120					125						
tat	gtg	ttc	aaa	gcc	gat	gat	caa	aca	aaa	gtg	cct	gag	atc	atc	agc		610
Tyr	Val	Phe	Lys	Ala	Asp	Asp	Gln	Thr	Lys	Val	Pro	Glu	Ile	Ile	Ser		
130					135					140				-	145		
tcc	atc	cgt	cag	gcg	ggg	aag	atc	gcc	cgg	cag	gag	gag	ctg	cac	tgc		658
Ser	Ile	Arg	Gln	Ala	Gly	Lys	Ile	Ala	Arg	Gln	Glu	Glu	Leu	His	Cys		
				150	-				155					160			
ccg	tcc	gag	ttc	gac	gac	acg	ttt	tcc	aag	aag	ttc	gag	gtg	ctc	ttc		706
Pro	Ser	Glu	Phe	Asp	Asp	Thr	Phe	Ser	Lys	Lys	Phe	Glu	Val	Leu	Phe		
			165					170					175				
tgc	ggc	cgc	gtg	acg	gtg	gcg	cac	aag	aag	gct	ccg	ccg	gcc	ctg	atc		754
Cys	Gly	Arg	Val	Thr	Val	Ala	His	Lys	Lys	Ala	Pro	Pro	Ala	Leu	Ile		
		180					185					190					
gac	gag	tgc	atc	gag	aag	ttc	aat	cac	gtc	agc	ggc	agc	cgg	9 99	tcc		802
Asp	Glu	Cys	Ile	Glu	Lys	Phe	Asn	His	Val	Ser	Gly	Ser	Arg	Gly	Ser		
	195					200					205						
gag	agc	ccc	cgc	CCC	aac	ccg	ccc	cat	gcc	gcg	ccc	aca	9 99	agc	cag		850
Glu	Ser	Pro	Arg	Pro	Asn	Pro	Pro	His	Ala	Ala	Pro	Thr	Gly	Ser	Gln		
210					215					220					225		
gag	cct	gtg	cgc	agg	ccc	atg	cgc	aag	tcc	ttc	tcc	cag	ccc	ggc	ctg		898
Glu	Pro	Val	Arg	Arg	Pro	Met	Arg	Lys	Ser	Phe	Ser	Gln	Pro	Gly	Leu		
			•	230					235					240			
cgc	tcg	ctg	gcc	ttt.	ragg	aag	gag	ctg	cag	gat	aaa	ggc	ctc	cga	agc		946
Arg	Ser	Leu	Ala	Phe	Arg	Lys	Glu	Leu	Gln	Asp	Gly	Gly	Leu	Arg	Ser		
			245					250					255				
agc	ggc	t tc	ttc	agc	tcc	ttc	gag	gag	agc	gac	att	gag	aac	cac	ctc		994
Ser	Gly	Phe	Phe	Ser	Ser	Phe	Glu	Glu	Ser	Asp	Ile	Glu	Asn	His	Leu		
		260					265					270					
att	agc	g ga	cac	aat	att	gtg	cag	CCC	aca	gat	atc	gag	gaa	aat	cga	1	042
Ile	Ser	Gly	His	Asn	Ile	Val	Gln	Pro	Thr	Asp	Ile	Glu	Glu	Asn	Arg		
	275					280					285						
act	atg	ctc	ttc	acg	att	g gc	cag	tct	gaa	gtt	tac	ctc	atc	agt	cct	1	1090
Thr	Met	Leu	Phe	Thr	Ile	Gly	Gln	Ser	Glu	Val	Tyr	Leu	Ile	Ser	Pro		
290					295					300					305		
gac	acc	aaa	a aa	ata	gca	ttg	gag	a aa	aat	ttt	aag	gag	ata	tcc	ttt	1	1138
Asp	Thr	Lys	Lys	Ile	Ala	Leu	Glu	Lys	Asn	Phe	Lys	Glu	Ile	Ser	Phe		
				310					315					320			
tgc	tct	cag	ggc	atc	aga	cac	gtg	gac	cac	ttt	999	ttt	atc	tgt	cgg	3	1186

wo	00/082	209						6	7					•	CIAD	
Cvs	Ser	Gln	Gly	Ile	Arg	His	Val	Asp	His	Phe	Gly	Phe	Ile	Cys	Arg	
- 4			325					330					335			
gag	tct	tcc	gga	ggt	ggc	ggc	ttt	cat	ttt	gtc	tgt	tac	gtg	ttt	cag	1234
-	-			Gly												
		340					345					350				
tgc	aca	aat	gag	gct	ctg	gtt	gat	gaa	att	atg	atg	acc	ctg	aaa	cag	1282
Cys	Thr	Asn	Glu	Ala	Leu	Val	Asp	Glu	Ile	Met	Met	Thr	Leu	Lys	Gln	
	355					360					365			-		
gcc	ttc	acg	gtg	gcc	gca	gtg	cag	cag	aca	gct	aag	gcg	cca	gcc	cag	1330
Ala	Phe	Thr	Val	Ala	Ala	Val	Gln	Gln	Thr	Ala	Lys	Ala	Pro	Ala	Gln	
370					375					380					385	
ctg	t gt	gag	ggc	tgc	CCC	ctg	caa	agc	ctg	cac	aag	ctc	tgt	gag	agg	1378
Leu	Cys	Glu	Gly	Cys	Pro	Leu	Gln	Ser	Leu	His	Lys	Leu	Cys	Glu	Arg	
				390					395					400		
	-			aat												1426
Ile	Glu	Gly	Met	Asn	Ser	Ser	Lys	Thr	Lys	Leu	Glu	Leu		Lys	His	
			405					410					415			
_	-			acc												1474
Leu	Thr	Thr	Leu	Thr	Asn	Gln		Gln	Ala	Thr	Ile		Glu	GIu	Val	
		420					425					430			_ 4_ 4_	1500
_				ccg												1522
Gln			Arg	Pro	Arg		Giu	GIn	Arg	GIU		GIU	ьeu	116	TIE	
	435					440		~~~		a2a	445	C = 5	cac	ato	cat	1570
				tgt Cys												
		Leu	Arg	Cys		ıyı	Giu	Gia	цуз	460	ביים	014			465	
450				` aag	455	202	tca	cad	ato	_	gča.	aaa	aat	att	gga	1618
															Gly	
116	GIY	GIU	Mec	. дуз 470	0111	****	-	0_11	475					480		
agt	gaa	ı tta	cca		agt	acc	act	cga	ttt	agg	cta	gat	ato	cto	g aaa	1666
_															Lys	
50-			485					490					495			
aac	aaa	ı gca	aag	aga	tct	tta	aca	gag	tct	tta	gaa	agt	att	ttg	g tcc	1714
															ı Ser	
		500					505					510				
cgg	g ggt	aat	: aaa	a gcc	aga	ggc	cto	g cag	gaa	a cac	tac	ato	agt	gt	g gat	1762
Arg	g Gly	y Ası	ı Lys	s Ala	Arg	gly	Leu	ı Glr	Gli	ı His	s Ser	: Ile	e Ser	r Vai	l Asp	
	51	5				520	•				525	5				
															g cca	1810
Let	ı Ası	p Se	r Se	r Lei	ı Ser	Ser	Thi	. Lei	ı Sei	r Ası	n Thi	s Se	r Ly:	s Gl	u Pro	
530					535					54					545	
tc	gt	g tg	t ga	a aag	ggag	ggc	ttg	g cco	ate	c tc	t gag	g ag	c to	c tt	t aag	1858

	_							6	8							•
Ser	Val	Cys	Glu	Lys	Glu	Ala	Leu	Pro	Ile	Ser	Glu	Ser	Ser	Phe	Lys	
				550					555					560		
ctc	ctc	ggc	tcc	tcg	gag	gac	ctg	tcc	agt	gac	tcg	gag	agt	cat	ctc	1906
Leu	Leu	Gly	Ser	Ser	Glu	Asp	Leu	Ser	Ser	Asp	Ser	Glu	Ser	His	Leu	
			565					570					57 5			
cca	gaa	gag	cca	gct	ccg	ctg	tcg	ccc	cag	cag	gcc	ttc	agg	agg	cga	1954
Pro	Glu	Glu	Pro	Ala	Pro	Leu	Ser	Pro	Gln	Gln	Ala	Phe	Arg	Arg	Arg	
		580					585					590				
gca	aac	acc	ctg	agt	cac	ttc	ccc	atc	gaa	tgc	cag	gaa	cct	cca	caa	2002
Ala	Asn	Thr	Leu	Ser	His	Phe	Pro	Ile	Glu	Cys	Gln	Glu	Pro	Pro	Gln	
	595					600					605					
cct	gcc	cgg	ggg	tcc	ccg	ggg	gtt	tcg	caa	agg	aaa	ctt	atg	agg	tat	2050
Pro	Ala	Arg	Gly	Ser	Pro	Gly	Val	Ser	Gln	Arg	Lys	Leu	Met	Arg	Tyr	
610					615					620					625	
cac	tca	gtg	agc	aca	gag	acg	cct	cat	gaa	cga	aag	gac	ttt	gaa	tcc	2098
His	Ser	Val	Ser	Thr	Glu	Thr	Pro	His	Glu	Arg	Lys	Asp	Phe	Glu	Ser	
				630					635					640		
aaa	gca	aac	cat	ctt	ggt	gat	tct	ggt	999	act	cct	gtg	aag	acc	cgg	2146
Lys	Ala	Asn	His	Leu	Gly	Asp	ser	Gly	Gly	Thr	Pro	Val	Lys	Thr	Arg	
			645					650					655			
agg	cat	tcc	tgg	agg	cag	cag	ata	ttc	ctc	cga	gta	gcc	acc	ccg	cag	2194
Arg	His	Ser	Trp	Arg	Gln	Gln	Ile	Phe	Leu	Arg	Val	Ala	Thr	Pro	Gln	
		660					665					670				
aag	gcg	tgc	gat	tct	tcc	agc	aga	tat	gaa	gat	tat	tca	gag	ctg	gga	2242
Lys	Ala	Cys	Asp	Ser	ser	Ser	Arg	Tyr	Glu	Asp	Tyr	Ser	Glu	Leu	Gly	
	675		,			680					685					
												gaa				2290
Glu	Leu	Pro	Pro	Arg	Ser	Pro	Leu	Glu	Pro	Val	Cys	Glu	Asp	Gly	Pro	
690					695					700					705	
															cga	2338
Phe	Gly	Pro	Pro	Pro	Glu	Glu	Lys	Lys	Arg	Thr	Ser	Arg	Glu	Leu	Arg	
				710					715					720		
															atg	2386
Glu	Lev	Trp	Gln	Lys	Ala	Ile	Leu	Glr	Gln	Ile	Leu	Leu	Lev	Arg	Met	
			72 5					730					735			
															aac	2434
Glu	ı Lys	s Glu	ı Asn	Gln	Lys	Leu	Gln	Ala	Ser	Glu	ı Asr	ı Asp	Lei	ı Let	ı Asn	
		740					745					750				
															a gaa	2482
Lys	s Arg	Lev	ı Lys	Leu	ı Asp	туг	Glu	ı Glı	1 Ile	th:			Lei	ı Lys	s Glu	
	75!					760					769					0530
gta	a act	t aca	a gtg	g tgg	g gaa	aaq	g ato	g cti	ago	c act	cca	a gga	a aga	a tca	a aaa	2530

WO 00/08209 PCT/IB99/01444 69

									17							•
Val	Thr	Thr	Val	Trp	Glu	Lys	Met	Leu	Ser	Thr	Pro	Gly	Arg	Ser	Lys	
770					7 75					780					785	
att	aag	ttt	gac	atg	gaa	aaa	atg	cac	tcg	gct	gtt	g gg	caa	ggt	gtg	2578
Ile	Lys	Phe	Asp	Met	Glu	Lys	Met	His	ser	Ala	Val	Gly	Gln	Gly	Val	
				790					795					800		
cca	cgt	cat	cac	cga	ggt	gaa	atc	tgg	aaa	ttt	cta	gct	gag	caa	ttc	2626
Pro	Arg	His	His	Arg	Gly	Glu	Ile	Trp	Lys	Phe	Leu	Ala	Glu	Gln	Phe	
			805					810					815	_		
cac	ctt	aaa	cac	cag	ttt	ccc	agc	aaa	cag	cag	cca	aag	gat	gtg	cca	2674
His	Leu	Lys	His	Gln	Phe	Pro	Ser	Lys	Gln	Gln	Pro	Lys	Asp	Val	Pro	
		820					825					830				
tac	aaa	gaa	ctc	tta	aag	cag	ctg	act	tcc	cag	cag	cat	gcg	att	ctt	2722
Tyr	Lys	Glu	Leu	Leu	Lys	Gln	Leu	Thr	Ser	Gln	Gln	His	Ala	Ile	Leu	
	835					840					845					
att	gac	ctt	g gg	cga	acc	ttt	cct	aca	cac	cca	tac	ttc	tct	gcc	cag	2770
Ile	Asp	Leu	Gly	Arg	Thr	Phe	Pro	Thr	His	Pro	Tyr	Phe	Ser	Ala	Gln	
850					855					860					865	
ctt	gga	gca	gga	cag	cta	tcg	ctt	tac	aac	att	ttg	aag	gcc	tac	tca	2818
Leu	Gly	Ala	Gly	Gln	Leu	Ser	Leu	Tyr	Asn	Ile	Leu	Lys	Ala	Tyr	Ser	
				870					875					880		
ctt	cta	gac	cag	gaa	gtg	gga	tat	tgc	caa	ggt	ctc	agc	ttt	gta	gca	2866
Leu	Leu	Asp	Gln	Glu	Val	Gly	Tyr	Cys	Gln	Gly	Leu	Ser	Phe	Val	Ala	
			885					890					895			
ggc	att	ttg	ctt	ctt	cat	atg	agt	gag	gaa	gag	gcg	ttt	aaa	atg	ctc	2914
Gly	Ile	Leu	Leu	Leu	His	Met	Ser	Glu	Glu	Glu	Ala	Phe	Lys	Met	Leu	
		900	•				905					910				
			atg													2962
Lys	Phe	Leu	Met	Phe	Asp	Met	Gly	Leu	Arg	Lys	Gln	Tyr	Arg	Pro	Asp	
	915					920					925					
_			tta													3010
Met	Ile	Ile	Leu	Gln	Ile	Gln	Met	Tyr	Gln	Leu	Ser	Arg	Leu	Leu	His	
930					935					940					945	
-			aga													3058
Asp	Tyr	His	Arg	Asp	Leu	Tyr	Asn	His	Leu	Glu	Glu	His	Glu	Ile	Gly	
				950					955					960		
	_														cag	3106
Pro	Ser	Leu	Tyr	Ala	Ala	Pro	Trp	Phe	Leu	Thr	Met	Phe			Gln	
			965					970					975			
															cag	3154
Phe	Pro			Phe	Val	Ala			Phe	Asp	Met			Lev	Gln	
		980					985					990				2202
gga	aca	gag	gtc	ata	ttt	aaa	gtg	gct	tta	agt	ctg	ttg	gga	ago	cat	3202

wo	00/082	209						-	70						PCT/IB99	/01444
	-						>		70 -	.	.	T	03	0	***	-
Gly		Glu	Val	Ile	Phe			Ala	Leu	ser			GTĀ	ser	HIS	
	995					1000					1005					3350
_									cta -							3250
Lys	Pro	Leu	Ile	Leu	Gln	His	Glu	Asn	Leu			TTE	vaı	Asp		
1010					1015					1020					1025	2222
		_							ttg							3298
Ile	Lys	Ser	Thr	Leu	Pro	Asn	Leu	Gly	Leu		Gln	Met	GLu			
				1030					1035					104		
									gct							3346
Ile	Asn	Gln	Val	Phe	Glu	Met	Asp	Ile	Ala	Lys	Gln	Leu	Gln,	Ala	Tyr	
			1045					1050					105			
_	-								gaa							3394
Glu	Val	Glu	Tyr	His	Val	Leu	Gln	Glu	Glu	Leu	Ile	Asp	Ser	Ser	Pro	
		106	0				106	5				107	0			
ctc	agt	gac	aac	caa	aga	atg	gat	aaa	tta	gag	aaa	acc	aac	agc	agc	3442
Leu	Ser	Asp	Asn	Gln	Arg	Met	Asp	Lys	Leu	Glu	Lys	Thr	Asn	Ser	Ser	
	107	5				108	0				108	5				
tta	cgc	aaa	cag	aac	ctt	gac	ctc	ctt	gaa	cag	ttg	cag	gtg	gca	aat	3490
Leu	Arg	Lys	Gln	Asn	Leu	Asp	Leu	Leu	Glu	Gln	Leu	Gln	Val	Ala	Asn	
109	0				109	5				110	0				1105	
ggt	agg	atc	caa	agc	ctt	gag	gcc	acc	att	gag	aag	ctc	ctg	agc	agt	3538
Gly	Arg	Ile	Gln	Ser	Leu	Glu	Ala	Thr	Ile	Glu	Lys	Leu	Leu	Ser	Ser	
				111	0				111	5				112	0	
gag	agc	aag	ctg	aag	cag	gcc	atg	ctt	acc	tta	gaa	ctg	gag	cgg	tcg	3586
Glu	Ser	Lys	Leu	Lys	Gln	Ala	Met	Leu	Thr	Leu	Glu	Leu	Glu	Arg	Ser	
			112	5				113	0				113	5		
gcc	ctg	ctg	cag	acg	gtg	gag	gag	ctg	cgg	cgg	cgg	agc	gca	gag	ccc	3634
Ala	Leu	Leu	Gln	Thr	Val	Glu	Glu	Leu	Arg	Arg	Arg	Ser	Ala	Glu	Pro	
		114	0				114	5				115	0			
ago	gac	cgg	gag	cct	gag	tgo	acg	cag	ccc	gag	ccc	acg	ggc	gac	tga	3682
Ser	Asp	Arg	g Glu	Pro	Glu	Cys	Thr	Gln	Pro	Glu	Pro	Thr	Gly	/ Asp	*	
	115	5				116	0				116	5				
caç	jctct	gca	ggag	agat	tg c	aaca	ccat	c cc	acac	tgto	cag	gcct	taa	ctga	aga g gga	3742
cag	gaaga	cgc	tgga	agga	ga g	aagg	gaago	g gg	jaagt	gtgo	tto	tcag	gga	ggaa	accggc	3802
															cagtgt	3862
															cacacag	3922
															aaaaaa	3982
	aaaa	-														3988

<210> 5

<211> 1168

<212> PRT

<213> Homo sapiens

<400	> 5														
Met	Glu	Pro	Ile	Thr	Phe	Thr	Ala	Arg	Lys	His	Leu	Leu	Pro	Asn	Glu
1				5					10					15	
Val	Ser	Val	Asp	Phe	Gly	Leu	Gln	Leu	Val	Gly	Ser	Leu	Pro	Val	His
			20					25					30		
Ser	Leu	Thr	Thr	Met	Pro	Met	Leu	Pro	Trp	Val	Val	Ala	Glu	Val	Arg
		35					40					45			
Arg	Leu	Ser	Arg	Gln	Ser	Thr	Arg	Lys	Glu	Pro	Val	Thr	Lys	Gln	Val
	50					55					60				
Arg	Leu	Cys	Val	Ser	Pro	Ser	Gly	Leu	Arg	Cys	Glu	Pro	Glu	Pro	Gly
65					70					75					80
Arg	Ser	Gln	Gln	Trp	Asp	Pro	Leu	Ile	Tyr	Ser	Ser	Ile	Phe	Glu	Cys
				85					90					95	
Lys	Pro	Gln	Arg	Val	His	Lys	Leu	Ile	His	Asn	Ser	His	Asp	Pro	Ser
			100					105					110		
Tyr	Phe	Ala	Cys	Leu	Ile	Lys	Glu	Asp	Ala	Val	His	Arg	Gln	Ser	Ile
		115					120					125			
Cys	Tyr	Val	Phe	Lys	Ala	Asp	Asp	Gln	Thr	Lys	Val	Pro	Glu	Ile	Ile
	130					135					140				
Ser	Ser	Ile	Arg	Gln	Ala	Gly	Lys	Ile	Ala	Arg	Gln	Glu	Glu	Leu	His
145					150					155					160
Cys	Pro	Ser	Glu	Phe	Asp	Asp	Thr	Phe	Ser	Lys	Lys	Phe	Glu	Val	Leu
				165					170					175	
Phe	Cys	Gly	Arg	\ Val	Thr	Val	Ala	His	Lys	Lys	Ala	Pro	Pro	Ala	Leu
			180		f -			185			ą i-		190		
Ile	Asp	Glu	Cys	Ile	Glu	Lys	Phe	Asn	His	Val	Ser	Gly	Ser	Arg	Gly
		195					200					205			
Ser	Glu	Ser	Pro	Arg	Pro	Asn	Pro	Pro	His	Ala	Ala	Pro	Thr	Gly	Ser
	210					215					220				
Gln	Glu	Pro	Val	Arg	Arg	Pro	Met	Arg	Lys	Ser	Phe	Ser	Gln	Pro	Gly
225					230					235					240
Leu	Arg	Ser	Leu	Ala	Phe	Arg	Lys	Glu	Leu	Gln	Asp	Gly	Gly	Leu	Arg
				245					250					255	
Ser	Ser	Gly	Phe	Phe	Ser	Ser	Phe	Glu	Glu	Ser	Asp	Ile	Glu	Asn	His
			260					265					270		
Leu	Ile	Ser	Gly	His	Asn	Ile	· Val	Gln	Pro	Thr	Asp	Ile	Glu	Glu	Asn
		275					280					285			
Arg	Thr	Met	Leu	Phe	Thr	Ile	Gly	Gln	Ser	Glu	Val	Tyr	Leu	Ile	Ser
	290					295					300				
Pro	Asp	Thr	Lys	Lys	Ile	Ala	Lev	Glu	Lys	Asn	Phe	Lys	Glu	Ile	Ser

									2						
305					310					315					320
Phe	Cys	Ser	Gln	Gly	Ile	Arg	His	Val	Asp	His	Phe	Gly	Phe	Ile	Cys
•				325					330					335	
Arg	Gľu	Ser	Ser	Gly	Gly	Gly	Gly	Phe	His	Phe	Val	Cys	Tyr	Val	Phe
J			340	-	_	_		345					350		
Gln	Cvs	ፐ ክዮ		Glu	Ala	Leu	Val	Asp	Glu	Ile	Met	Met	Thr	Leu	Lys
0	O, E	355					360					365			*
Gln	λla		Thr	Va l	αlα	Δla		Gln	Gln	Thr	sία		Ala	Pro	Ala
GIII		FIIC	LILL	Vai	riiu	375	vuı	J	0111	****	380				
01	370	~	<i>C</i> 1	~1··	Cura		Len	Cln.	Ser	Leu		Lare	T.633	Cve	Glu
	Leu	Cys	GIU.	GIŸ		PIO	пец	Gili	261	395	HITS	БуЗ	пси	Cys	400
385		~3	~3		390	a	G	T	m}		T	~ 1	τ	01-	
Arg	He	G1u	GIY		Asn	ser	ser	гÀг		Lys	ьeu	GIU	Leu		гÀг
				405	_			-	410					415	
His	Leu	Thr	Thr	Leu	Thr	Asn	Gln		GIn	Ala	Thr	11e		GIu	Glu
			420					425					430		
Val	Gln	Lys	Leu	Arg	Pro	Arg	Asn	Glu	Gln	Arg	Glu	Asn	Glu	Leu	Ile
		435					440					445			
Ile	Ser	Phe	Leu	Arg	Cys	Leu	Tyr	Glu	Glu	Lys	Gln	Lys	Glu	His	Ile
	450					455					460				
His	Ile	Gly	Glu	Met	Lys	Gln	Thr	Ser	Gln	Met	Ala	Ala	Glu	Asn	Ile
465					470					475					480
Gly	Ser	Glu	Leu	Pro	Pro	Ser	Ala	Thr	Arg	Phe	Arg	Leu	Asp	Met	Leu
				485					490					495	
Lys	Asn	Lys	Ala	Lys	Arg	Ser	Leu	Thr	Glu	Ser	Leu	Glu	Ser	Ile	Leu
			500					505					510		
Ser	Arg	Gly	Asn	Lys	Ala	Arg	Gly	Leu	Gln	Glu	His	Ser	Ile	Ser	Val
		515			<i>f</i> -		520				- 	_525		,-	
Asp	Leu	Asp	Ser	Ser		Ser	Ser	Thr	Leu	Ser	Asn	Thr	Ser	Lys	Glu
	530					535					540				
Pro	Ser	Val	Cys	Glu	Lys	Glu	Ala	Leu	Pro	Ile	Ser	Glu	Ser	Ser	Phe
545					550					555					560
Lys	Leu	Leu	Gly	Ser	Ser	Glu	Asp	Leu	Ser	Ser	Asp	Ser	Glu	Ser	His
				565					570					575	
Leu	Pro	Glu	Glu	Pro	Ala	Pro	Leu	Ser	Pro	Gln	Gln	Ala	Phe	Arg	Arg
			580					585					590		
Arg	Ala	Asn	Thr	Leu	Ser	His	Phe	Pro	Ile	Glu	Cys	Gln	Glu	Pro	Pro
		595					600					605			
Gln	Pro	Ala	Arg	Gly	Ser	Pro	Gly	Val	Ser	Gln	Arg	Lys	Leu	Met	Arg
	610		J	•		615					620				
Tvr			Val	Ser	Thr	Glu	Thr	Pro	His	Glu	Arg	Lys	Asp	Phe	Glu
625					630					635		-			640
		Ala	Asn	His			Asp	Ser	Gly	Gly	Thr	Pro	Val	Lys	Thr
	- 1 -					-	_		_	-					

								7	3						
				645					650					655	
Arg .	Arg	His	Ser	Trp	Arg	Gln	Gln	Ile	Phe	Leu	Arg	Val	Ala	Thr	Pro
			660					665					670		
Gln	Lys	Ala	Cys	Asp	Ser	Ser	Ser	Arg	Tyr	Glu	Asp	Tyr	Ser	Glu	Leu
		675					680					685			
Gly	Glu	Leu	Pro	Pro	Arg	Ser	Pro	Leu	Glu	Pro	Val	Cys	Glu	Asp	Gly
-	690					695					700				
Pro	Phe	Gly	Pro	Pro	Pro	Glu	Glu	Lys	Lys	Arg	Thr	Ser	Arg	Glu	Leu
705					710					715					720
Arq	Glu	Leu	Trp	Gln	Lys	Ala	Ile	Leu	Gln	Gln	Ile	Leu	Leu	Leu	Arg
- · · · · ·			-	725	_				730					735	
Met	Glu	Lys	Glu	Asn	Gln	Lys	Leu	Gln	Ala	Ser	Glu	Asn	Asp	Leu	Leu
		-	740			_		745					750		
Asn	Lvs	Ara	Leu	Lvs	Leu	Asp	Tyr	Glu	Glu	Ile	Thr	Pro	Cys	Leu	Lys
	- <i>1</i> -	755		-		-	760					765			
Glu	Val		Thr	Val	Trp	Glu	Lys	Met	Leu	Ser	Thr	Pro	Gly	Arg	Ser
-	770				-	7 75	_				780				
Lvs		Lvs	Phe	Asp	Met	Glu	Lys	Met	His	Ser	Ala	Val	Gly	Gln	Gly
785		-1			790		•			795					800
	Pro	Ara	His	His		Gly	Glu	Ile	Trp	Lys	Phe	Leu	Ala	Glu	Gln
·		3		805	_	-			810					815	
Phe	His	Leu	Lys	His	Gln	Phe	Pro	Ser	Lys	Gln	Gln	Pro	Lys	Asp	Val
			820					825					830		
Pro	Tvr	Lys	Glu	Leu	Leu	Lys	Gln	Leu	Thr	Ser	Gln	Gln	His	Ala	Ile
		835				_	840					845			
Leu	Ile		Leu	Gly	Arg	Thr	Phe	Pro	Thr	His	Pro	Tyr	Phe	Ser	Ala
	850	-			f -	855					860		-	-	
Gln	Leu	Gly	Ala	Gly	Gln	Leu	Ser	Leu	Tyr	Asn	Ile	Leu	Lys	Ala	Tyr
865		_			870					875					880
Ser	Leu	Leu	Asp	Gln	Glu	Val	Gly	Tyr	Cys	Gln	Gly	Leu	Ser	Phe	Val
				885					890					895	
Ala	Gly	Ile	Leu	Leu	Leu	His	Met	Ser	Glu	Glu	Glu	Ala	Phe	Lys	Met
	-		900					905					910		
Leu	Lys	Phe	Leu	Met	Phe	Asp	Met	Gly	Leu	Arg	Lys	Gln	Tyr	Arg	Pro
	•	915					920					925			
qzA	Met	Ile	· Ile	Leu	Gln	Ile	Gln	Met	Tyr	Gln	Leu	Ser	Arg	Leu	Leu
-15	930					935					940				
His			His	Arc	, Asp	Leu	і Туг	Asn	His	Leu	Glu	Glu	His	: Glu	Ile
945	-	-			950					955					960
	Pro	Ser	Leu	туг	Ala	Ala	Pro	Trp	Phe	Leu	Thr	Met	Phe	e Ala	Ser
-				965					970					979	
Gln	Phe	Pro	Leu	ı Gly	/ Phe	val	L Ala	a Arg	y Val	Phe	Asp	Met	Ile	e Phe	Leu

74 980 985 990 Gln Gly Thr Glu Val Ile Phe Lys Val Ala Leu Ser Leu Leu Gly Ser 1000 995 His Lys Pro Leu Ile Leu Gln His Glu Asn Leu Glu Thr Ile Val Asp 1015 1020 Phe Ile Lys Ser Thr Leu Pro Asn Leu Gly Leu Val Gln Met Glu Lys 1030 1035 1025 Thr Ile Asn Gln Val Phe Glu Met Asp Ile Ala Lys Gln Leu Gln Ala 1045 1050 Tyr Glu Val Glu Tyr His Val Leu Gln Glu Glu Leu Ile Asp Ser Ser 1060 1065 1070 Pro Leu Ser Asp Asn Gln Arg Met Asp Lys Leu Glu Lys Thr Asn Ser 1080 Ser Leu Arg Lys Gln Asn Leu Asp Leu Glu Gln Leu Gln Val Ala 1090 1095 1100 Asn Gly Arg Ile Gln Ser Leu Glu Ala Thr Ile Glu Lys Leu Leu Ser 1110 1115 Ser Glu Ser Lys Leu Lys Gln Ala Met Leu Thr Leu Glu Leu Glu Arg 1130 1125 1135 Ser Ala Leu Leu Gln Thr Val Glu Glu Leu Arg Arg Arg Ser Ala Glu 1140 1145 Pro Ser Asp Arg Glu Pro Glu Cys Thr Gln Pro Glu Pro Thr Gly Asp 1160 1165 1155 <210> 6 <211> 18 <u>.</u> :-<212> DNA 1. <213> Artificial Sequence <220> <221> misc binding <222> 1..18 <223> sequencing oligonucleotide PrimerPU <400> 6 tgtaaaacga cggccagt

18

<210> 7
<211> 18
<212> DNA
<213> Artificial Sequence

WO 00/08209		PCT/IB99/01444
-	75	

<220>

<221> misc_binding

<222> 1..18

<223> sequencing oligonucleotide PrimerRP

<400> 7

caggaaacag ctatgacc

18